

## Computing Optimal Ate Pairings on Elliptic Curves with Embedding Degree 9, 15 and 27

Emmanuel Fouotsa · Nadia El Mrabet · Aminatou Pecha

10 November 2018

**Abstract** Much attention has been given to efficient computation of pairings on elliptic curves with even embedding degree since the advent of pairing-based cryptography. The existing few works in the case of odd embedding degrees require some improvements. This paper considers the computation of optimal ate pairings on elliptic curves of embedding degrees  $k = 9, 15$  and  $27$  which have twists of order three. Mainly, we provide a detailed arithmetic and cost estimation of operations in the tower extensions field of the corresponding extension fields. A good selection of parameters enables us to improve the theoretical cost for the Miller step and the final exponentiation using the lattice-based method comparatively to the previous few works that exist in these cases. In particular for  $k = 15$  and  $k = 27$  we obtained an improvement, in terms of operations in the base field, of up to 25% and 29% respectively in the computation of the final exponentiation. Also, we obtained that elliptic curves with embedding degree  $k = 15$  present faster results than BN12 curves at the 128-bit security levels. We provided a MAGMA implementation in each case to ensure the correctness of the formulas used in this work.

**Keywords** Elliptic Curves · Optimal Pairings · Miller's algorithm · Extension fields arithmetic · Final exponentiation

---

E.Fouotsa  
Department of Mathematics, Higher Teacher Training College,  
University of Bamenda  
E-mail: emmanuel Fouotsa@yahoo.fr

N. El Mrabet  
LIASD, Université Paris 8,  
93526 Saint Denis Cedex (France)  
*SAS-CMP Gardanne, France*

A. Pecha  
LIASD, Université Paris 8,  
93526 Saint Denis Cedex (France)  
*Institut Supérieur du Sahel, Université de Maroua P.O.Box 46 Maroua, Cameroon*

**Mathematics Subject Classification (2000)** 14H52 · 1990S

## 1 Introduction

Pairings are bilinear maps defined on the group of rational points of elliptic or hyper elliptic curves [40]. They enable to realise many cryptographic protocols such as the Identity-Based Cryptosystem [10], Identity-Based Encryption [12], the Identity-Based Undeniable Signature [30], Short Signatures [11] or Broadcast Encryption [19]. A survey of some applications of pairings can be found in [16], [9, Chapter X]. These many applications justify the research on the efficient computation of pairings. Generally, let  $E$  be an ordinary elliptic curve defined over a finite field  $\mathbb{F}_p$  and  $r$  a large prime divisor of the order of the group  $E(\mathbb{F}_p)$ . The embedding degree of  $E$  with respect to  $r$  and the prime number  $p$  is the smallest integer  $k$  such that  $r \mid p^k - 1$ . The  $\rho$ -value of the elliptic curve  $E$  is the value  $\log p / \log r$  measuring the size of the base field relatively to the size  $r$  of a subgroup of  $E(\mathbb{F}_p)$ . The Tate pairing and its variants are the most used in cryptography. They map two linearly independent points of the subgroup of order  $r$  of  $E(\mathbb{F}_{p^k})$  to the group of  $r$ -th roots of unity in the finite field  $\mathbb{F}_{p^k}$ . The computation of the Tate pairing and its variants consists of an application of the Miller algorithm [35] and a final exponentiation. Efficient computation of pairings requires construction of pairing-friendly elliptic curves over  $\mathbb{F}_p$  with prescribed embedding degree  $k$  (see for example [8] or [17]) and efficient arithmetic in the towered fields associated to  $\mathbb{F}_{p^k}$  (see [27], [20], [25], [14]). A lot of work has been done for shortening the Miller loop leading to the concept of pairing lattices [22], or the optimal pairing described by Vercauteren which can be computed with the smallest number of iterations in the Miller algorithm [39]. Due to these advances, the final exponentiation step has become a serious task. In this work, we concentrate on elliptic curves  $E$  over  $\mathbb{F}_p$  with embedding degree 9, 15 and 27. These curves admit twists of degree three which enable computations

Security level	Bit length of $r$	Bit length of $p^k$	$k$ $\rho \approx 1$	$k$ $\rho \approx 2$
80	160	960 – 1280	6 – 8	3 – 4
128	256	3000 – 5000	12 – 20	6 – 10
192	384	8000 – 10000	20 – 26	10 – 13
256	512	14000 – 18000	28 – 36	14 – 18

**Table 1** Bit sizes of curves parameters and corresponding embedding degrees to obtain commonly desired levels of security.

to be done in subfields and also lead to the denominator elimination technique. To our knowledge just few works ([29], [36] and [41]) exist in these cases and much attention have been given only to elliptic curves with even embedding degree (see for example [1],[18]). Also, another motivation to our work is the recent results on the resolution of discrete logarithm problem [26].

Indeed according to the first analysis of this article, as for instance in [21],[4] the security level for elliptic curves with friable embedding degree should be taken greater than those presented in Table 1. The main consequence is that elliptic curves with embedding degree 12 or 18 may not be the one assuring a nice ratio between the security level and the arithmetic. Elliptic curves with odd embedding degree could become interesting and more efficient than elliptic curves with even embedding degree. Also it is also noticed that elliptic curves with odd embedding degree especially with  $k = 27$  may be suitable for computing product of pairings [41]. In this work we consider the following parameter's sizes ( $p \approx 2^{343}$ ,  $r \approx 2^{257}$ ,  $\rho = 1.33$ ), ( $p \approx 2^{575}$ ,  $r \approx 2^{385}$ ,  $\rho = 1.5$ ), ( $p \approx 2^{579}$ ,  $r \approx 2^{514}$ ,  $\rho = 1.12$ ) for curves with  $k = 9, 15, 27$  respectively. This corresponds to the 128, 192 and 256-bit security levels respectively according to recommendations in Table 1 [17]. However, considering the recent recommendations based on the advances on Discrete Logarithm computation with the Number Field Sieve (NFS) algorithm and its variants ([26], [5], [34], [4]) the security level provided by the above parameters may reduce. Indeed, let  $2^{-d} \exp((c + o(1))(\log Q)^{1/3} (\log \log Q)^{2/3})$  where  $d$  and  $c$  are constants, be the running time of the NFS algorithm, with  $Q = p^k$ . The base-two logarithm of this runtime with ( $o(1) = 0$ ) gives  $S(Q, c, d) = c(\lg e)(\log Q)^{1/3}(\log \log Q)^{2/3} - d$ . We then know that the constants  $c$  and  $d$ , the embedding degree  $k$  and the security level  $l$  must satisfy the following Pollard-Rho security and variants of NFS security constraints  $\log Q/k \geq 2\rho l$  and  $S(Q, c, d) \geq l$ . Therefore the previous parameters provide a security level of 109, 168, 214 bits instead of 128, 192, 256-bit respectively for curves with  $k = 9, 15, 27$ . But for now we still consider in Section 4, 5 and 6 recommendation from Table 1 in order to make a fair comparison with previous works. Later in section 8, we consider advances in discrete logarithm computation and provide tentative updated parameters at the 128, 192, 256 bits respectively for curves with  $k = 9, 15, 27$ . So we proposed a detailed arithmetic in the towering fields associated to the fields  $\mathbb{F}_{p^9}$ ,  $\mathbb{F}_{p^{15}}$  and  $\mathbb{F}_{p^{27}}$ . The lattice-based method explained by Fuentes *et al.* [18] is applied to compute the final exponentiation in the cases  $k = 9, 15$ . We also find a simple expression and explicit cost evaluation for the optimal pairing in the cases  $k = 9$  and  $k = 15$  comparatively to the work in [36]. The results obtained are an improvement with respect to previous works [29], [36] and [41] respectively for  $k = 9, 15$  and 27. Precisely, our contributions (see Table 3 and subsection 8.4 for comparison) in this work are:

1. Determination of an explicit cost of the computation of the optimal pairing for elliptic curves stated above. This includes a good selection of parameters for a shorter Miller loop and an efficient exponentiation. In particular, we saved one inversion in  $\mathbb{F}_{p^{27}}$  for the computation of the Miller loop in the case  $k = 27$ .
2. Details on the arithmetic in the tower of subfields of  $\mathbb{F}_{p^9}$ ,  $\mathbb{F}_{p^{15}}$  and  $\mathbb{F}_{p^{27}}$ . Especially, we give the cost of the computation of Frobenius maps and Inversions in the cyclotomic subgroups of  $\mathbb{F}_{p^9}^*$ ,  $\mathbb{F}_{p^{15}}^*$  and  $\mathbb{F}_{p^{27}}^*$ , (see Appendices A, B and C).

3. Improvement of the costs of the final exponentiation by saving  $828M_1 + 145S_1$ ,  $1170M_1 + 7767S_1$  and  $8676M_1 + 32136S_1$  operations for elliptic curves of embedding degrees 9, 15 and 27 respectively, comparatively to previous works in these cases; where  $M_k, S_k$  represent the costs of multiplication and squaring in the finite field  $\mathbb{F}_{p^k}$ .
4. In Section 8 we look for new parameters considering the advances in Discrete Logarithm computation to update the cost of the optimal ate pairings on the studied curves at the 128, 192 and 256-bit security levels. We then compare our results with known curves such as BN and BLS curves. In particular we obtained that elliptic curves with embedding degree  $k = 15$  present faster results than BN12 curves at the 128-bit security levels.

We also provide a MAGMA implementation in each case to ensure the correctness of the formulas used in this work. The code is available in [?].

The rest of this paper is organised as follows: In Section 2 we briefly present the Tate and ate pairings together with the Miller algorithm for their efficient computation, we also recall the concept of optimal ate pairing and the lattice-based method for computing the final exponentiation. Sections 4, 5 and 6 present arithmetic in subfields, and costs estimation of the Miller step and the final exponentiation when considering the embedding degrees  $k = 9, 15$  and 27 respectively. Each of these sections includes a comparative analysis with previous work. Section 7 presents a general comparison of the results obtained in this work and the previous results in the literature. In Section 8 we look for new parameters considering the advances in Discrete Logarithm computation to update the cost of the optimal ate pairings on the studied curves at the 128, 192 and 256-bit security levels. We then compare our results with known curves such as BN and BLS curves. We conclude the work in Section 9 in which we suggest as future work the search for parameters to have *subgroup secure* ordinary curves [6] and to ensure protection against *small-subgroup attacks*[32].

### Notations

The following notations are used in this work.

$M_k, S_k, I_k$  : Cost of multiplication, squaring and inversion in the field  $\mathbb{F}_{p^k}$ , for any integer  $k$ .

## 2 Background and previous works

### 2.1 Pairings and the Miller Algorithm

Let  $E$  be an elliptic curve defined over  $\mathbb{F}_p$ , a finite field of characteristic  $p > 3$ . Let  $r$  be a large prime factor of the group order of the elliptic curve. Let  $m \in \mathbb{Z}$  and  $P \in E(\mathbb{F}_p)[r]$  a point of  $E$  of order  $r$  with coordinates in  $\mathbb{F}_p$ . Let  $f_{m,P}$  a function with divisor  $\text{Div}(f_{m,P}) = m(P) - ([m]P) - (m-1)(\mathcal{O})$  where  $\mathcal{O}$  denotes the identity element of the group of points of the elliptic curve. We denote  $k$  the smallest integer such that  $r$  divides  $p^k - 1$ , also called



## 2.2 Use of Twists

Twists of elliptic curves enable to efficiently compute pairings. Indeed, in the Miller algorithm the doubling of a point (line 3) and the addition of points (lines 6 and 8) are done in the extension field  $\mathbb{F}_{p^k}$  in the case of the ate pairing. The use of twists enables to perform these operations rather in a subfield of  $\mathbb{F}_{p^k}$  and also leads to the denominator elimination. More precisely, a twist of an elliptic curve  $E$  defined over a finite field  $\mathbb{F}_p$  is an elliptic curve  $E'$  defined over  $\mathbb{F}_p$  which is isomorphic to  $E$  over an algebraic closure of  $\mathbb{F}_p$ . The smallest integer  $d$  such that  $E$  and  $E'$  are isomorphic over  $\mathbb{F}_{p^d}$  is called the degree or the order of the twist. Elliptic curves of embedding degree  $k = 9, 15$  and  $k = 27$  admit twists of order three. Explicit constructions of such curves can be found in [33], [15] and [7]. The general equation of these curves is given by  $E : y^2 = x^3 + b$ . The equation defining the twist  $E'$  has the form  $y^2 = x^3 + b\omega^6$  where  $\{1, \omega, \omega^2\}$  is the basis of the  $\mathbb{F}_{p^{k/3}}$ -vector space  $\mathbb{F}_{p^k}$  and the isomorphism between  $E'$  and  $E$  is  $\psi : E' \rightarrow E; (x', y') \mapsto (x'/\omega^2, y'/\omega^3)$ . Using this isomorphism, points  $Q$  in  $\mathbb{G}_2$  can be instead taken as  $(x\omega^{-2}, y\omega^{-3})$  where  $(x, y) \in E'(\mathbb{F}_{p^{k/3}})$ . The function  $h_{R,S}$  is defined by  $h_{R,S}(x, y) = \frac{y + \lambda(x_R - x) - y_R}{x - x_{R+S}}$  where  $\lambda$  is the slope of the line passing through  $R$  and  $S$ . Observe that using the equation of the curve  $y^2 = x^3 + b$  one has  $x - x_{R+S} = \frac{y^2 - y_{R+S}^2}{x^2 + x_{R+S}x + x_{R+S}^2}$ . In the present case of ate pairing, the addition  $R+S$  is performed in the extension field  $\mathbb{F}_{p^k}$  and the function  $h_{R,S}$  is evaluated at a point  $(x_P, y_P) \in E(\mathbb{F}_p)$ . So using the twist, the points  $R, S$  and  $R+S$  are taken in the form  $(x\omega^{-2}, y\omega^{-3})$  where  $(x, y) \in E'(\mathbb{F}_{p^{k/3}})$ . Therefore we have  $h_{R,S}(x_P, y_P) = \frac{(y_P\omega^5 + \lambda(x_R\omega^2 - x_P\omega^4) - y_R\omega^2)(x_P^2\omega^4 + x_{R+S}x_P\omega^2 + x_{R+S}^2)}{y_P^2\omega^9 - y_{R+S}^2\omega^3}$ . We observe that the denominator is an element of the subfield  $\mathbb{F}_{p^{k/3}}$  and so will be sent to 1 during the final exponentiation (line 10 in the Miller algorithm) since  $p^{k/3} - 1$  is a factor of  $p^k - 1$ . Consequently we simply ignore that denominator in the Miller algorithm for an efficient computation. More details on twists can be found in [13].

## 2.3 Optimal Pairings

The reduction of Miller's loop length is an important way to improve the computation of pairings. The latest work is a generalized method to find the shortest loop, which leads to the concept of optimal pairings due to Vercauteren [39]. Let  $\lambda = mr$  be a multiple of  $r$  such that  $r \nmid m$  and write  $\lambda = \sum_{i=0}^l c_i p^i = h(p)$ , ( $h(z) \in \mathbb{Z}[z]$ ). Recall that  $h_{R,S}$  is the Miller function defined in section 2.1. For  $i = 0, \dots, l$  set  $s_i = \sum_{j=i}^l c_j p^j$ ; then the map

$$e_o : \mathbb{G}_2 \times \mathbb{G}_1 \rightarrow \mu_r \quad (1)$$

$$(Q, P) \mapsto \left( \prod_{i=0}^l f_{c_i, Q}^{p^i}(P) \cdot \prod_{i=0}^{l-1} h_{[s_{i+1}]Q, [c_i p^i]Q}(P) \right)^{\frac{p^k - 1}{r}}$$

defines a bilinear pairing and non degenerate if

$$mkp^k \neq ((p^k - 1)/r) \cdot \sum_{i=0}^l ic_i p^{i-1} \pmod r.$$

The coefficients  $c_i : i = 0, \dots, l$  can be obtained from the short vectors obtained from the lattice

$$L = \begin{pmatrix} r & 0 & 0 & \dots & 0 \\ -p & 1 & 0 & \dots & 0 \\ -p^2 & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ -p^{\phi(k)-1} & 0 & 0 & \dots & 1 \end{pmatrix} \quad (2)$$

#### 2.4 Final Exponentiation and the Lattice-Based Method for its Computation

The result of the Miller loop's step is raised to the power  $\frac{p^k-1}{r}$ . This step is called the final exponentiation (line 10 in Miller's algorithm). The efficient computation of the final exponentiation has become a serious task. Observe that this exponent can be divided into two parts as follows:

$$\frac{p^k - 1}{r} = \left[ \frac{p^k - 1}{\phi_k(p)} \right] \cdot \left[ \frac{\phi_k(p)}{r} \right]$$

where  $\phi_k(x)$  is the  $k$ -th cyclotomic polynomial. The final exponentiation is

therefore computed as  $f^{\frac{p^k-1}{r}} = \left[ f^{\frac{p^k-1}{\phi_k(p)}} \right]^{\frac{\phi_k(p)}{r}}$ . The computation of the first

part  $A = f^{\frac{p^k-1}{\phi_k(p)}}$  is generally inexpensive as it consists of few multiplications, inversion and  $p$ -th powering in  $\mathbb{F}_{p^k}$ . The second part  $A^{\frac{\phi_k(p)}{r}}$  is considered more difficult and is called the hard part. An efficient method to compute the hard part is described by Scott *et al.* [37]. They suggested to write  $d = \frac{\phi_k(p)}{r}$  in base  $p$  as  $d = d_0 + d_1p + \dots + d_{\phi(k)-1}p^{\phi(k)-1}$  and find a short vectorial addition chains to compute  $A^d$  much more efficiently than the naive method. In [18], based on the fact that a fixed power of a pairing is still a pairing, Fuentes *et al.* [18] suggested to apply Scott *et al.*'s method with a power of any multiple  $d'$  of  $d$  with  $r$  not dividing  $d'$ . This could lead to a more efficient exponentiation than computing  $A^d$  directly. Their idea of finding the polynomial  $d'(x)$  is to apply the *LLL*-algorithm to the matrix formed by  $\mathbb{Q}$ -linear combinations of the elements  $d(x), xd(x), \dots, x^{degr-1}d(x)$ . They successfully applied the method in the case of elliptic curves of embedding degrees 8, 12 and 18 [18]. In Sections 4 and 5 we apply this method to improve the computation of the final exponentiation for elliptic curves of embedding degree  $k = 9$  and 15. A clever method was used by Zhang *et al.* [41] to compute the final exponentiation in the case  $k = 27$ .

### 3 Arithmetic in the tower of subfields of $\mathbb{F}_{p^9}$ , $\mathbb{F}_{p^{27}}$ and $\mathbb{F}_{p^{15}}$

A pairing is computed as an element of the extension field  $\mathbb{F}_{p^k}$ . But its efficient computation depends on the arithmetic of subfields of  $\mathbb{F}_{p^k}$  which is generally organised as tower of subfields extensions. In this section we recall the tower extension of finite fields  $\mathbb{F}_{p^9}$ ,  $\mathbb{F}_{p^{27}}$  and  $\mathbb{F}_{p^{15}}$ . We also give explicit cost of the arithmetic operations. For extension-field arithmetic in  $\mathbb{F}_{p^9}$  and  $\mathbb{F}_{p^{27}}$  we consider  $p \equiv 1 \pmod{3}$  motivated by the work of Barreto et al.[7] on the construction of elliptic curves of embedding degree 9 and 27. This implies that  $\mathbb{F}_{p^k}$  can be represented as  $\mathbb{F}_{p^{k/3}}[X]/(X^3 - \alpha)$ , for  $k = 3^i, i = 1, 2, 3$  where  $\alpha$  is a cubic non residue modulo  $p$ . Since  $p \equiv 1 \pmod{3}$  we have that  $X^3 - 7$  irreducible over  $\mathbb{F}_p$ . Therefore cubic extensions will be constructed using the polynomials  $X^3 - \alpha_i$  where  $\alpha_i = 7^{1/3^{i-1}}$ . A tower extension for  $\mathbb{F}_{p^{27}}$  together with the one for  $\mathbb{F}_{p^9}$  are then given by:

$$\begin{aligned}\mathbb{F}_{p^3} &= \mathbb{F}_p[u] \quad \text{with } u^3 = 7 \\ \mathbb{F}_{p^9} &= \mathbb{F}_{p^3}[v] \quad \text{with } v^3 = 7^{1/3} \\ \mathbb{F}_{p^{27}} &= \mathbb{F}_{p^9}[w] \quad \text{with } w^3 = 7^{1/9}\end{aligned}$$

The costs of the computation of the Frobenius maps and cyclotomic inversions are given in Lemma 1 for the extension  $\mathbb{F}_{p^9}$ . The proof of this Lemma is given in Appendix A.

**Lemma 1** *In the finite field  $\mathbb{F}_{p^9}$ ,*

1. *The computation of the  $p^3; p^6$ -Frobenius maps costs  $6M_1$*
2. *The computation of the  $p; p^2; p^4; p^5; p^7; p^8$ -Frobenius maps costs  $8M_1$*
3. *The inverse of an element  $\alpha$  of the  $G_{\phi_3(p^3)}$ -order cyclotomic subgroup is computed as  $\alpha^{-1} = \alpha^{p^3} \cdot \alpha^{p^6}$  and the cost is  $36S_1$*

Similarly, in the finite field  $\mathbb{F}_{p^{27}}$  the Lemma 2 gives the costs of the computation of the Frobenius maps and cyclotomic inversions. The proof of this Lemma is given in Appendix B.

**Lemma 2** *In the finite field  $\mathbb{F}_{p^{27}}$ ,*

1. *The computation of the  $p^3; p^6; p^9$ -Frobenius maps costs  $18M_1$*
2. *The computation of the  $p; p^2; p^4; p^5; p^7; p^8$ -Frobenius maps costs  $26M_1$*
3. *The inverse of an element  $\alpha$  of the  $G_{\phi_3(p^9)}$ -order cyclotomic subgroup is computed as  $\alpha^{-1} = \alpha^{p^9} \cdot \alpha^{p^{18}}$  and the cost is  $216S_1$*

In the case of  $\mathbb{F}_{p^{15}}$ , we consider pairing friendly curves over  $\mathbb{F}_p$  where  $p \equiv 1 \pmod{5}$  [15]. According to [31, Theorem 3.75] the polynomial  $X^5 - \alpha$  is irreducible over  $\mathbb{F}_p[X]$  if and only if  $\alpha$  is neither a cubic root nor a fifth root in  $\mathbb{F}_p$ . A tower extension for  $\mathbb{F}_{p^{15}}$  can be constructed as follows:

$$\begin{aligned}\mathbb{F}_{p^5} &= \mathbb{F}_p[u] \quad \text{with } u^5 = 7 \\ \mathbb{F}_{p^{15}} &= \mathbb{F}_{p^5}[v] \quad \text{with } v^3 = u. \text{ where } u \in \mathbb{F}_{p^5}\end{aligned}$$

Fields	Operations	Costs
$\mathbb{F}_{p^3}$	Multiplication $M_3$ Squaring $S_3$ Inversion $I_3$	$6M_1$ $5S_1$ $I_1 + 9M_1 + 2S_1$
$\mathbb{F}_{p^9}$	Multiplication $M_9$ Squaring $S_9$ Inversion $I_9$ Frobenius $p^3; p^6$ Frobenius $p; p^2; p^4; p^5; p^7; p^8$ Inversion in $G_{\phi_3(p^3)}$	$36M_1$ $25S_1$ $I_1 + 63M_1 + 12S_1$ $6M_1$ $8M_1$ $18M_1 + 15S_1$
$\mathbb{F}_{p^{27}}$	Multiplication $M_{27}$ Squaring $S_{27}$ Inversion $I_{27}$ Frobenius $p^3; p^6; p^9$ Frobenius $p; p^2; p^4; p^5; p^7; p^8$ Inversion in $G_{\phi_3(p^9)}$	$216M_1$ $125S_1$ $I_1 + 387M_1 + 62S_1$ $18M_1$ $26M_1$ $108M_1 + 75S_1$
$\mathbb{F}_{p^5}$	Multiplication $M_5$ Squaring $S_5$ Inversion $I_5$	$9M_1$ $9S_1$ $I_1 + 45M_1 + 5S_1$
$\mathbb{F}_{p^{15}}$	Multiplication $M_{15}$ Squaring $S_{15}$ Inversion $I_{15}$  Frobenius $p^5; p^{10}$ Frobenius $p; p^2; p^3; p^4; p^6; p^7; p^8; p^9$ Inversion in $G_{\phi_3(p^5)}$	$45M_1$ $45S_1$ $I_1 + 126M_1 +$ $23S_1$ $10M_1$ $14M_1$ $27M_1 + 27S_1$

**Table 2** Cost of operations in extension fields from [29], [36] and [41] and this work (see Appendix A, B and C)

Our main contribution in this section is the computation of Frobenius maps and the inversions in the  $\phi_n(\cdot)$ -order cyclotomic subgroup of  $\mathbb{F}_{p^k}^*$ . The costs of the computation of the Frobenius maps and cyclotomic inversions are given in Lemma 3.

**Lemma 3** *In the finite field  $\mathbb{F}_{p^{15}}$ ,*

1. *The computation of the  $p^5; p^{10}$ -Frobenius maps costs  $10M_1$*
2. *The computation of the  $p; p^2; p^3; p^4; p^6; p^7; p^8; p^9$ -Frobenius maps costs  $14M_1$*
3. *The inverse of an element  $\alpha$  of the  $G_{\phi_3(p^5)}$ -order cyclotomic subgroup is computed as  $\alpha^{-1} = \alpha^{p^5} \cdot \alpha^{p^{10}}$  and the cost is  $54S_1$*

*Proof* The proof is given in Appendix C.

In Table 2 we summarise the overall cost of operations in the tower of subfields described above. The costs for squaring, multiplication and inversion are from [29], [36] and [41] respectively for  $k = 9, 15$  and  $27$ . Explicit details of the cost of Frobenius maps and inversions in the cyclotomic subgroups are given in Appendix A, B and C.

## 4 Elliptic Curves with Embedding Degree 9

This section describes the computation of the optimal ate pairing (Miller step and the final exponentiation) on the parameterized elliptic curve defined in [33]. The correctness of the results can be verified with the MAGMA code available in [?]. This family of elliptic curves has embedding degree 9 and a  $\rho$ -value 1.33 and is parameterized by :

$$\begin{aligned} p &= ((x+1)^2 + ((x-1)^2(2x^3+1)^2)/3)/4 \\ r &= (x^6 + x^3 + 1)/3 \\ t &= x + 1 \end{aligned} \quad (3)$$

### 4.1 Optimal ate pairing

Based on the general framework described by Vercauteren in [39], the short vector obtained from the lattice  $L$  defined by equation (2) gives the optimal function  $h(z) = \sum_{i=0}^5 c_i z^i = x - z \in \mathbb{Z}[z]$ . A straightforward application of formula (1) yields the optimal pairing

$$\begin{aligned} e_o : \mathbb{G}_2 \times \mathbb{G}_1 &\longrightarrow \mu_r \\ (Q, P) &\longmapsto f_{x,Q}(P)^{\frac{p^9-1}{r}} \end{aligned}$$

### 4.2 Cost of the execution of the Miller loop

The Miller loop consists of the doubling steps (line 3 in the Miller algorithm) and addition steps (line 6 or 8 in the Miller algorithm). These steps use the Miller function  $h_{R,S}$  either in affine coordinates or in projective coordinates. The work of Zhang *et al.* [41, Section 3] presents the currently fastest formulas in projective coordinates. The doubling step costs  $9M_1 + 3M_3 + 9S_3$  and the cost of the addition step is  $9M_1 + 12M_3 + 5S_3$ . For an explicit cost of the computation of  $f_{x,Q}(P)$ , we wrote a Pari/GP code to find a suitable  $x$  with low Hamming weight and minimal number of bits for the 128 bit-security level according to Table 1. The best value we were able to find is  $x = 2^{43} + 2^{37} + 2^7 + 1$  which gives an  $r(x)$  prime of 257 bits and  $p(x)$  a prime of 343 bits. The values  $p$  and  $x$  are both congruent to 1 modulo 6 so that the corresponding elliptic curve is  $y^2 = x^3 + 1$  [29]. The computation of  $f_{x,Q}(P)$  therefore costs 43 doubling steps, 3 additions, 42 squaring and 45 multiplications in  $\mathbb{F}_{p^9}$ . Thus the total cost for the computation of the Miller loop for the optimal pairing on elliptic curves of embedding degree 9 is  $43(9M_1 + 3M_3 + 9S_3) + 3(9M_1 + 12M_3 + 5S_3) + 42S_9 + 45M_9$ . This is equal to  $45M_9 + 165M_3 + 414M_1 + 42S_9 + 402S_3$ . Using the arithmetic in Table 2, the overall cost is  $3024M_1 + 3060S_1$ . To our knowledge, no other explicit cost with a specific value of  $x$  is reported in the literature.

### 4.3 Cost of the computation of the final exponentiation

As explained in Section 2, the final exponentiation in this case can be divided as  $f^{(p^9-1)/r} = \left(f^{p^3-1}\right)^{(p^6+p^3+1)/r} = \left(f^{p^3-1}\right)^d$ . We then used the lattice method described by Fuentes *et al.* [18] that we briefly explained in Section 2.4. It is applied to the following matrix in which the coefficient 243 is used to obtain integer entries as  $d = (p^6 + p^3 + 1)/r$  is a polynomial with rational coefficients

$$M = \begin{pmatrix} 243d(x) \\ 243xd(x) \\ 243x^2d(x) \\ 243x^3d(x) \\ 243x^4d(x) \\ 243x^5d(x) \end{pmatrix} \quad (4)$$

We obtain the following multiple of  $d$ :  $d' = x^3d = k_0 + k_1p + k_2p^2 + k_3p^3 + k_4p^4 + k_5p^5$  where the polynomials  $k_i, i = 0, \dots, 5$  are as follows

$$\begin{aligned} k_0 &= -x^4 + 2x^3 - x^2, & k_1 &= -x^3 + 2x^2 - x, & k_2 &= -x^2 + 2x - 1, \\ k_3 &= x^7 - 2x^6 + x^5 + 3, & k_4 &= x^6 - 2x^5 + x^4, & k_5 &= x^5 - 2x^4 + x^3 \end{aligned}$$

They verify the relations (see the code in [?] for verification)

$$k_2 = -(x-1)^2, \quad k_1 = xk_2, \quad k_0 = xk_1, \quad k_5 = -xk_0, \quad k_4 = xk_5, \quad k_3 = xk_4 + 3$$

If we set  $A = f^{p^3-1}$  then

- The cost for the computation of  $A$  is 1  $p^3$ -Frobenius, 1 Inversion in  $\mathbb{F}_{p^9}$  and 1 multiplication in  $\mathbb{F}_{p^9}$ .
- The cost of the computation of  $A^{k_0}, A^{k_1}$  and  $A^{k_4}$  is 3 exponentiations by  $x$ ,
- The cost of the computation of  $A^{k_5}$  is one inversion in the cyclotomic subgroup and one exponentiation by  $x$ .
- The cost of the computation of  $A^{k_2}$  is one inversion in the cyclotomic subgroup and two exponentiations by  $(x-1)$ .
- The cost of the computation of  $A^{k_3}$  is 2 multiplications, one squaring and one exponentiation by  $x$ .

Note that the inversion in the cyclotomic subgroup  $G_{\phi_3(p^3)}$  of order  $p^6 + p^3 + 1$  is computed as  $A^{-1} = A^{p^3} \cdot A^{p^6}$  (see Appendix A for details and cost). The cost for the hard part  $A^{d'}$  is then 2 exponentiations by  $x-1$ , 5 exponentiations by  $x$ , 7 multiplications in  $\mathbb{F}_{p^9}$ , one squaring in  $\mathbb{F}_{p^9}$ , two cyclotomic inversions  $I_{G_{\phi_3(p^3)}}$  and  $p, p^2, p^3, p^4, p^5$ -Frobenius maps. Using the value of  $x$  given above, one exponentiation by  $x$  costs  $43S_9 + 3M_9$  whereas one exponentiation by  $x-1$  costs  $43S_9 + 2M_9$ . Finally the hard part costs  $2(43S_9 + 2M_9) + 5(43S_9 + 3M_9) + 7M_9 + 1S_9 + 2I_{G_{\phi_3(p^3)}} = 302S_9 + 26M_9 + 2I_{G_{\phi_3(p^3)}}$  and  $p, p^2, p^3, p^4, p^5$ -Frobenius maps. The total cost of the final exponentiation is  $1I_9 + 27M_9 + 302S_9 + 2I_{G_{\phi_3(p^3)}}$  and  $p, p^2, 2 * p^3, p^4, p^5$ -Frobenius maps.

#### 4.4 Improvement and comparison with previous work

From the results in [29], the hard part costs  $309S_9 + 50M_9$  and  $p, p^2, p^3, p^4, p^5$ -Frobenius maps. If we include the cost  $1I_9 + 1M_9$  and  $p^3$ -Frobenius for the easy part and using the arithmetic in Table 2, the overall cost is  $I_1 + 1115M_1 + 7592S_1$  for this work and  $I_1 + 1943M_1 + 7737S_1$  for Le et al. [29]. We therefore save  $828M_1 + 145S_1$  comparatively to their work. Although the sizes of  $p$  are 343 bits in this work and 348 bits in [29], the cost of a multiplication is almost the same in the two corresponding fields (see section 8.4).

### 5 Elliptic Curves with Embedding Degree 15

In this section we give explicit formulas together with their cost for the Miller loop in the computation of the optimal ate pairing. We then compute the cost of the final exponentiation on the parameterized elliptic curve defined in [15]. The correctness of the results can be verified in [?]. This family of elliptic curves has embedding degree 15 and a  $\rho$ -value 1.5 and is parameterized by :

$$\begin{aligned} p &= (x^{12} - 2x^{11} + x^{10} + x^7 - 2x^6 + x^5 + x^2 + x + 1)/3 \\ r &= x^8 - x^7 + x^5 - x^4 + x^3 - x + 1 \\ t &= x + 1 \end{aligned} \tag{5}$$

#### 5.1 Optimal ate pairing

The Vercauteren approach described in [39] enabled us to obtain the short vector from the lattice  $L$  defined by equation (2) which lead to the optimal function  $h(z) = \sum_{i=0}^5 c_i z^i = x - z \in \mathbb{Z}[z]$ . A straightforward application of formula (1) yields the optimal pairing

$$\begin{aligned} e_o : \mathbb{G}_2 \times \mathbb{G}_1 &\longrightarrow \mu_r \\ (Q, P) &\longmapsto f_{x,Q}(P)^{\frac{p^{15}-1}{r}} \end{aligned}$$

#### 5.2 Cost of the computation of the Miller loop

In this section, we consider the Miller function given in affine coordinates, following the analysis of Lauter et al. [28] who suggested to use affine coordinates at higher security level. Miller function used for the computation of  $f_{x,Q}(P)$  in this case is described in [41, Table 2] with the fastest cost to date. At 192 bits security level on elliptic curves with  $k = 15$ , the best value of  $x$  we were able to find with a Pari/GP code is  $x = 2^{48} + 2^{41} + 2^9 + 2^8 + 1$ . This value gives an  $r(x)$  prime of 385 bits and  $p(x)$  of 575 bits which correspond to parameters for 192-bit security level according to Table 1. The value of  $p$  is congruent to 1 modulo 5 and a curve equation can be  $y^2 = x^3 + 1$ . The Miller

loop consists here of computing  $f_{x,Q}$  which costs 48 doubling steps, 4 additions steps, 47 squaring and 51 multiplications in  $\mathbb{F}_{p^{15}}$ . Considering the currently fastest cost for doubling and addition step in [41, Table 2], the Miller loop costs  $48(15M_1 + 3M_5 + 2S_5 + I_5) + 4(15M_1 + 3M_5 + 2S_5 + I_5) + 47S_{15} + 51M_{15}$ , which is  $51M_{15} + 156M_5 + 780M_1 + 47S_{15} + 104S_5 + 52I_5$ . Using the arithmetic in Table 2, the overall cost is  $52I_1 + 6819M_1 + 3311S_1$ . To our knowledge no explicit cost is reported in the literature in the case  $k = 15$  with a specific value of  $x$ .

### 5.3 Cost of the computation of the final exponentiation

The final exponentiation in this case is written in a different way as  $f^{(p^{15}-1)/r} = \left(f^{p^5-1}\right)^{(p^{10}+p^5+1)/r} = \left(f^{p^5-1}\right)^d$ . This decomposition is used instead of  $\frac{p^{15}-1}{r} = \left[\frac{p^{15}-1}{\phi_{15}(p)}\right] \cdot \left[\frac{\phi_{15}(p)}{r}\right]$  as usually done, for efficiency reasons in the computation. Observe that  $\frac{p^{15}-1}{\phi_{15}(p)} = p^7 + p^6 + p^5 - p^2 - p - 1$  and  $\phi_{15}(p) = p^8 - p^7 + p^5 - p^4 + p^3 - p + 1$  will lead to several multiplications and Frobenius maps operations. Thus the lattice method described by Fuentes *et al.* [18] that we briefly explained in Section 2.4 is applied to the following matrix  $M$ . In the matrix  $M$  the coefficient  $\frac{59049}{19683}$  is used to obtain integer entries as  $d = (p^{10} + p^5 + 1)/r$  is a polynomial with rational coefficients.

$$M = \begin{pmatrix} \frac{59049}{19683}d(x) \\ \frac{59049}{19683}xd(x) \\ \frac{59049}{19683}x^2d(x) \\ \cdot \\ \cdot \\ \frac{59049}{19683}x^7d(x) \end{pmatrix} \quad (6)$$

We then obtained the following multiple of  $d$ :  $d' = 3x^3d = k_0 + k_1p + \dots k_9p^9$  where the polynomials  $k_i, i = 0, \dots, 9$  are defined as follows

$$\begin{aligned} k_0 &= -x^6 + x^5 + x^3 - x^2, & k_1 &= -x^5 + x^4 + x^2 - x \\ k_2 &= -x^4 + x^3 + x - 1 \\ k_3 &= x^{11} - 2x^{10} + x^9 + x^6 - 2x^5 + x^4 - x^3 + x^2 + x + 2 \\ k_5 &= x^{11} - x^{10} - x^8 + x^7 + 3 \\ k_4 &= x^{11} - x^{10} - x^9 + x^8 + x^6 - x^5 - x^4 + x^3 - x^2 + 2x + 2 \\ k_6 &= x^{10} - x^9 - x^7 + x^6 & k_7 &= x^9 - x^8 - x^6 + x^5 \\ k_8 &= x^8 - x^7 - x^5 + x^4 & k_9 &= x^7 - x^6 - x^4 + x^3 \end{aligned}$$

The polynomials  $k_i : i = 0, \dots, 9$  verify the relations (see the code in [?] for verification)

$$\begin{aligned} k_2 &= -(x-1)^2(x^2+x+1), & k_1 &= xk_2, & k_0 &= xk_1 \\ k_9 &= -xk_0, & k_8 &= xk_9, & k_7 &= xk_8 \\ k_6 &= xk_7, & k_5 &= xk_6 + 3, & k_4 &= M - (k_1 + k_7) \\ k_3 &= M - (k_0 + k_6 + k_9) \text{ where } M &= (k_2 + k_5 + k_8) \end{aligned}$$

Set  $A = f^{p^5-1}$  then

- The cost for the computation of  $A$  is 1  $p^5$ -Frobenius, 1 Inversion in  $\mathbb{F}_{p^{15}}$  and 1 multiplication in  $\mathbb{F}_{p^{15}}$  and .
- The computation of  $A^{k_2}$  is 2 exponentiations by  $x$ , 2 exponentiations by  $x - 1$ , 2 multiplications and 1 cyclotomic inversion,
- The cost of the computation of the  $A^{k_0}, A^{k_1}, A^{k_6}, A^{k_7}$  is 5 exponentiations by  $x$ , the computation of  $A^{k_9}$  costs 1 exponentiation by  $x$  and 1 cyclotomic inversion,
- The computation of  $A^{k_5}$  is 1 exponentiation by  $x$ , 2 multiplications and 1 squaring in  $\mathbb{F}_{p^{15}}$ ,
- The computation of  $A^{k_4}$  costs 4 multiplications in  $\mathbb{F}_{p^{15}}$  and 1 cyclotomic inversion,
- The computation of  $A^{k_3}$  costs 3 multiplications in  $\mathbb{F}_{p^{15}}$  and 1 cyclotomic inversion.

Therefore, the cost of the computation of  $A^{d'}$  is 2 exponentiations by  $x - 1$ , 9 exponentiations by  $x$ , 20 multiplications, one squaring in  $\mathbb{F}_{p^{15}}$ , four inversions in the cyclotomic subgroup  $G_{\phi_3(p^5)}$  of order  $p^{10} + p^5 + 1$  (note that  $A^{-1} = A^{p^5} \cdot A^{p^{10}}$  see Appendix C for details) and  $p, p^2, p^3, p^4, p^5, p^6, p^7, p^8, p^9$ -Frobenius maps. Using the value of  $x$  given above, the cost of the hard part is  $2(48S_{15} + 3M_{15}) + 9(48S_{15} + 4M_{15}) + 20M_{15} + 1S_{15} + 4I_{G_{\phi_3(p^5)}} = 529S_{15} + 62M_{15} + 4I_{G_{\phi_3(p^5)}}$  and  $p, p^2, p^3, p^4, p^5, p^6, p^7, p^8, p^9$ -Frobenius maps. The total cost of the final exponentiation in this work is therefore  $1I_{15} + 529S_{15} + 63M_{15} + 4I_{G_{\phi_3(p^5)}}$  and  $p, p^2, p^3, p^4, 2 * p^5, p^6, p^7, p^8, p^9$ -Frobenius maps.

*Remark 1* The cost given by Le *et al.* [29] for the hard part is 11 exponentiations by  $x$ , 22 multiplications, 2 inversions in  $\mathbb{F}_{p^{15}}$  and 9 Frobenius maps. The authors said that the cost of an inversion in  $\mathbb{F}_{p^{15}}$  is free with a reference to a similar computation but on elliptic curves with even embedding degree, unfortunately we do not see how this is possible. Also, they considered an  $x$  of 64 bits and hamming weight 7 and claimed that the cost is  $88M_{15} + 528S_{15}$  instead of  $11(6M_{15} + 64S_{15}) = 88M_{15} + 704S_{15}$ . Therefore if we count the 2 inversions in  $\mathbb{F}_{p^{15}}$  (these inverses are in fact in the cyclotomic subgroup  $G_{\phi_3(p^5)}$ ), then their final cost is  $88M_{15} + 704S_{15} + 2I_{G_{\phi_3(p^5)}}$  and 11 Frobenius maps, whereas our cost is  $529S_{15} + 62M_{15} + 4I_{G_{\phi_3(p^5)}}$ .

#### 5.4 Improvement and comparison with previous work

Considering the previous remark, the cost of the final exponentiation in [29] is  $1I_{15} + 704S_{15} + 89M_{15} + 2I_{G_{\phi_3(p^5)}}$  and  $p, p^2, p^3, p^4, 2 * p^5, p^6, p^7, p^8, p^9$ -Frobenius maps. We observe that we have improved the results by saving  $26M_{15} + 175S_{15} - 2I_{G_{\phi_3(p^5)}}$ . Using the arithmetic in Table 2, the overall cost is  $I_1 + 3066M_1 + 24071S_1$  for this work and  $I_1 + 4236M_1 + 31838S_1$  for Le *et al.* [29].

We therefore save  $26M_{15} + 175S_{15} - 2I_{G_{\phi_3(p^5)}} = 1170M_1 + 7767S_1$  comparatively to their work. A MAGMA code for the implementation to ensure the correctness of the decomposition of the final exponentiation and the Miller function is available in [?].

## 6 Elliptic Curves with Embedding Degree 27

The parameterized elliptic curves with embedding degree 27 is defined in [7]. This family has a  $\rho$ -value  $10/9$  and is parameterized by the following polynomials:

$$\begin{aligned} p &= 1/3(x-1)^2(x^{18} + x^9 + 1) + x \\ r &= 1/3(x^{18} + x^9 + 1) \\ t &= x + 1 \end{aligned} \tag{7}$$

### 6.1 The Miller loop and the final exponentiation

The Miller loop and the final exponentiation has been studied in [41]. They found the optimal function  $h(z) = \sum_{i=0}^{17} c_i z^i = x - z \in \mathbb{Z}[z]$  and the optimal pairing is given by

$$e_o : \mathbb{G}_2 \times \mathbb{G}_1 \longrightarrow \mu_r; (Q, P) \longmapsto f_{x,Q}(P)^{\frac{p^{27}-1}{r}}$$

The authors in [41] used the parameter  $x = 2^{28} + 2^{27} + 2^{25} + 2^8 - 2^3$  for their computation at 256-bit security level. The cost of the Miller step that they obtained is therefore  $28(3M_9 + 2S_9 + 1I_9 + 9M_1) + 4(3M_9 + 2S_9 + 1I_9 + 9M_1) + 27(6S_9) + 30(6M_9) + 1I_{27} = 276M_9 + 226S_9 + 32I_9 + 288M_1 + I_{27}$  operations. The computation of the final exponentiation in [41] requires  $1I_{27} + 11M_{27}$ , 17 powers of  $x$ , 2 powers of  $x - 1$  and  $p, p^2, p^3, p^4, p^5, p^6, p^7, p^8, 2 * p^9$ -Frobenius maps. Therefore the explicit cost of the final exponentiation is  $1I_{27} + 17(4(6M_9) + 28(6S_9) + 36M_1) + 2(5(6M_9) + 28(6S_9) + 36M_1) + 11(6M_9) + 228M_1 = 1I_{27} + 648M_9 + 3192S_9 + 912M_1$ .

Then the explicit cost for the computation of the Miller loop and the final exponentiation given in that work is  $12627M_1 + 8670S_1 + 33I_1$  and  $24627M_1 + 114998S_1 + 1I_1$  respectively (see [41] for details).

*Remark 2* The negative coefficient in the value of  $x$  affects the efficiency since one full inversion in  $\mathbb{F}_{p^{27}}$  is required in the Miller algorithm (line 8) and also 19 inversions in the cyclotomic subgroup are required when raising to the power of  $x$  during the final exponentiation.

In the next section we explain the choice of another parameter to avoid these additional operations.

## 6.2 Improvement and comparison with previous work

We use the arithmetic (especially the computation of inversion in the cyclotomic subgroup) and a specific value of  $x$  to improve the costs in [41]. Precisely, a careful search with a Pari/GP code enabled us to find the value  $x = 2^{29} + 2^{19} + 2^{17} + 2^{14}$  so that  $r$  has a prime factor of 514 bits length and the prime  $p$  has a bit length of 579 for 256-bit security level according to Table 1. An adequate elliptic curve has the equation  $y^2 = x^3 - 2$ . Although the corresponding base field is a bit larger (579 bits instead of 573 in [41], but  $m_{579} \approx m_{573}$  see section 8.4 for notations) and we have an extra doubling step, we avoid the full inversion in  $\mathbb{F}_{p^{27}}$  and 17 inversions in the cyclotomic subgroup  $\mathbb{G}_{\phi_3(p^9)}$  when raising to power  $x$ . We perform 2 inversions in the cyclotomic subgroup only when raising to power  $x - 1$ . The cost of the Miller loop now becomes  $29(3M_9 + 2S_9 + 1I_9 + 9M_1) + 3(3M_9 + 2S_9 + 1I_9 + 9M_1) + 27(6S_9) + 30(6M_9) = 276M_9 + 226S_9 + 32I_9 + 288M_1$ . Using the arithmetic in Table 2, the overall cost for the Miller loop is  $32I_1 + 12240M_1 + 6034S_1$  for this work where we saved at least one inversion in  $\mathbb{F}_{p^{27}}$ .

Our cost for the final exponentiation is  $1I_{27} + 17(3(6M_9) + 29(6S_9)) + 2(4(6M_9) + 29(6S_9)) + 2I_{\mathbb{G}_{\phi_3(p^9)}} + 11(6M_9) = 1I_{27} + 420M_9 + 3306S_9 + 2I_{\mathbb{G}_{\phi_3(p^9)}}$  and  $p, p^2, p^3, p^4, p^5, p^6, p^7, p^8, 2 * p^9$ -Frobenius maps. Using the arithmetic in Table 2, the overall cost is  $I_1 + 15951M_1 + 82862S_1$  for this work. The implementation of this pairing is available in [?].

## 7 General Comparison

In this section, we summarize the different costs obtained in this work and compare our results with previous works.

Curves	References	Miller loop	Final Exponentiation
$k=9$ 128-bit	Previous work [29]	<i>No specific cost reported</i>	$I_1 + 1943M_1 + 7737S_1$
	This work	$3024M_1 + 3060S_1$	$I_1 + 1115M_1 + 7592S_1$
$k=15$ 192-bit	Previous work [36]	<i>No specific cost reported</i>	$I_1 + 4263M_1 + 31811S_1$
	This work	$52I_1 + 6819M_1 + 3311S_1$	$I_1 + 3093M_1 + 24044S_1$
$k=27$ 256-bit	Previous work [41]	$33I_1 + 12627M_1 + 8670S_1$	$I_1 + 24627M_1 + 114998S_1$
	This work	$32I_1 + 12240M_1 + 6034S_1$	$I_1 + 15951M_1 + 82862S_1$

**Table 3** Comparison of the cost of the Miller loop and the final exponentiation.

If we assume that the cost of a squaring is the same as the cost of a multiplication then the cost of the final exponentiation is  $I_1 + 27137M_1$  and  $I_1 + 36074M_1$  for this work and previous work [29] respectively for  $k = 15$ . The theoretical improvement obtained in this work is therefore up to 25%. A similar analysis with  $k = 9$  yields an improvement of 8%. In the case of curves with  $k = 27$ , our results present an improvement of 14% and 29% for the Miller loop and final exponentiation respectively compare to the work in [41].

## 8 New Parameters for Optimal Ate pairing on Elliptic Curves with embedding degrees 9, 15 and 27

In this section we consider new parameters for parameterized curves of embedding degrees 9, 15 and 27 at the 128, 192 and 256-bit security levels. We consider recent advances on the computation of the discrete logarithm thanks to the Number Field Sieve (NFS) algorithm and its variants described in some papers as mentioned in the introduction. Mostly, the paper of Barbulescu and Duquesne [4] presents a detailed approach for computing new parameters for pairings. Based on their analysis, Scott and Guillevic [38] have proposed tentative general recommended parameters for classical security level and that we reproduce here in Table 4.

DL Algorithm	AES-128	AES-192	AES-256
NFS	3072	7680	15360
exTNFS	3618	9241	18480
SexTNFS	5004	12871	27410

**Table 4** Recommended extension fields size ( $p^k$ ) to obtain desired levels of security [38].

Following the Table 4, we searched for new parameters that will ensure resistance to SexTNFS algorithm at the various security levels for curves of embedding degrees 9, 15 and 27.

### 8.1 New parameters and costs for optimal ate pairing at the 128-bit security level for $k = 9$ and $k = 15$

- Case of  $k = 9$ . Following the recommendation from Table 4, we found the value  $x = 2^{70} + 2^{59} + 2^{46} + 2^{41} + 1$ . This gives a prime  $p$  of 559 bits and a prime  $r$  of 419 bits. We proceed as described in Section 4 to obtain the cost of the Miller loop and the final exponentiation. The Miller loop costs in this case  $70(9M_1 + 3M_3 + 9S_3) + 4(9M_1 + 12M_3 + 5S_3) + 69S_9 + 73M_9$ . This is equal to  $73M_9 + 258M_3 + 666M_1 + 69S_9 + 650S_3$ . Using the arithmetic in Table 2, the overall cost is  $4842M_1 + 4975S_1$ . Using the value of  $x$  given above, the hard part of the final exponentiation costs  $2(70S_9 + 3M_9) + 5(70S_9 + 4M_9) + 7M_9 + 1S_9 + 2I_{G_{\phi_3(p^3)}} = 491S_9 + 33M_9 + 2I_{G_{\phi_3(p^3)}}$  and  $p, p^2, p^3, p^4, p^5$ -Frobenius maps. The total cost of the final exponentiation is  $1I_9 + 34M_9 + 491S_9 + 2I_{G_{\phi_3(p^3)}}$  and  $p, p^2, 2 * p^3, p^4, p^5$ -Frobenius maps for a total cost of  $I_1 + 1367M_1 + 12317S_1$ .
- Case of  $k = 15$ . Following the recommendation from Table 4, we found the value  $x = 2^{31} + 2^{19} + 2^5 + 2^2$ . This gives a prime  $p$  of 371 bits and a prime  $r$  of 249 bits. We proceed as described in section 5 to obtain the cost of the Miller loop and the final exponentiation. The Miller loop costs in this case  $3(15M_1 + 13M_5 + 3S_5) + 31(15M_1 + 6M_5 + 7S_5) + 30S_{15} + 33M_{15}$ . This is equal to  $33M_{15} + 225M_5 + 510M_1 + 30S_{15} + 226S_5$ . Using the arithmetic in Table

2, the overall cost is  $4020M_1 + 3384S_1$ . Using the value of  $x$  given above, the hard part of the final exponentiation costs  $2(31S_{15} + 4M_{15} + 1I_{G_{\phi_3(p^5)}}) + 9(31S_{15} + 3M_{15}) + 20M_{15} + 1S_{15} + 4I_{G_{\phi_3(p^5)}} = 55M_{15} + 342S_{15} + 6I_{G_{\phi_3(p^5)}}$  and  $p, p^2, p^3, p^4, p^5, p^6, p^7, p^8, p^9$ -Frobenius maps. The total cost of the final exponentiation is  $I_{15} + 56M_{15} + 342S_{15} + 6I_{G_{\phi_3(p^5)}}$  and  $p, p^2, p^3, p^4, 2 * p^5, p^6, p^7, p^8, p^9$ -Frobenius maps for a total cost of  $I_1 + 2940M_1 + 15575S_1$ .

The following table 5 compares our results with previous results at the 128-bit security level.

Curves-Ref.	Miller loop	Final Exp.	Size of $p$	Total( $S_1 = M_1$ )
$k = 15$ (This work)	$4020M_1 + 3384S_1$	$I_1 + 2940M_1$ $+15575S_1$	371	$I_1 + 25919M_1$
KSS16 [4]	$7534M_1$	$I_1 + 18542M_1$	340	$I_1 + 26076M_1$
BLS12 [4]	$7708M_1$	$I_1 + 8295M_1$	461	$I_1 + 16003M_1$
BN12 [4,2]	$12068M_1$	$I_1 + 7485M_1$	461	$I_1 + 19553M_1$
$k = 9$ (This work)	$4842M_1 + 4975S_1$	$I_1 + 1367M_1$ $+12317S_1$	559	$I_1 + 23501M_1$

**Table 5** Comparison of the cost of the Miller loop and the final exponentiation at 128-bit security level.

## 8.2 New parameters and costs for pairings at the 192- bit security levels for $k = 15$ and $k = 27$

- Case of  $k = 15$ . Following the recommendation from Table 4, we found the value  $x = 2^{72} + 2^{40} + 2^9 + 2^5 + 1$ . This gives a prime  $p$  of 863 bits and a prime  $r$  of 577 bits. We proceed as described in section 5 to obtain the cost of the Miller loop and the final exponentiation. The Miller loop costs in this case  $4(15M_1 + 13M_5 + 3S_5) + 72(15M_1 + 6M_5 + 7S_5) + 71S_{15} + 75M_{15}$ . This is equal to  $75M_{15} + 484M_5 + 1140M_1 + 71S_{15} + 516S_5$ . Using the arithmetic in Table 2, the overall cost is  $8871M_1 + 7839S_1$ . Using the value of  $x$  given above, the hard part of the final exponentiation costs  $2(72S_{15} + 3M_{15}) + 9(72S_{15} + 4M_{15}) + 20M_{15} + 1S_{15} + 4I_{G_{\phi_3(p^5)}} = 62M_{15} + 793S_{15} + 4I_{G_{\phi_3(p^5)}}$  and  $p, p^2, p^3, p^4, p^5, p^6, p^7, p^8, p^9$ -Frobenius maps. The total cost of the final exponentiation is  $I_{15} + 63M_{15} + 793S_{15} + 4I_{G_{\phi_3(p^5)}}$  and  $p, p^2, p^3, p^4, 2 * p^5, p^6, p^7, p^8, p^9$ -Frobenius maps for a total cost of  $I_1 + 3201M_1 + 35816S_1$ .
- Case of  $k = 27$ . Following the recommendation from Table 4, we found the value  $x = 2^{25} + 2^{14} + 2^{17} + 2^4 + 1$ . This gives a prime  $p$  of 511 bits and a prime factor of  $r$  of 410 bits. We proceed as described in section 6 to obtain the cost of the Miller loop and the final exponentiation. The Miller loop costs in this case  $4(9M_1 + 1I_9 + 2S_9 + 3M_9) + 25(9M_1 + 1I_9 + 2S_9 + 3M_9) + 24S_{27} + 27M_{27}$ . This is equal to  $29I_9 + 27M_{27} + 87M_9 +$

$261M_1 + 24S_{27} + 58S_9$ . Using the arithmetic in Table 2, the overall cost is  $29I_1 + 11052M_1 + 4798S_1$ . Using the value of  $x$  given above, the final exponentiation costs  $I_{27} + 2(25S_{27} + 3M_{27}) + 17(25S_{27} + 4M_{27}) + 11M_{27} + 2I_{G_{\phi_3(p^9)}}$  and  $p, p^2, p^3, p^4, p^5, p^6, p^7, p^8, 2*p^9$ -Frobenius maps for a total cost of the final exponentiation  $I_1 + 19191M_1 + 59587S_1$ .

The cost for the case  $k = 24$  are obtained with the parameter given in [4] and the formulas from [2]

The following table 6 compares our results with previous results at the 192-bit security level.

Curves-Ref.	Miller loop	Final Exp.	Size of $p$	Total( $S_1 = M_1$ )
$k = 15$ (This work)	$8871M_1 + 7839S_1$	$I_1 + 3201M_1$ $+35816S_1$	863	$I_1 + 55727M_1$
BLS27 (This work)	$29I_1 + 11052M_1$ $+4798S_1$	$I_1 + 19191M_1$ $+59587S_1$	511	$30I_1 + 94628M_1$
KSS18 [4]	$15270M_1 + 2590S_1$	$8I_1 + 7977M_1$ $+18310S_1$	677	$8I_1 + 44147M_1$
BLS24 [4, 2]	$15495M_1$	$10I_1 + 27914M_1$	554	$10I_1 + 43409M_1$

**Table 6** Comparison of the cost of the Miller loop and the final exponentiation at 192-bit security level.

### 8.3 New parameters and costs for pairings at the 256- bits security levels for $k = 27$ and $k = 24$

- Case of  $k = 27$ . Following the recommendation from Table 4, we found the value  $x = 2^{51} + 2^{42} + 2^{28} + 2^9 + 1$ . This gives a prime  $p$  of 1019 bits and a prime factor of  $r$  of 883 bits. We proceed as described in section 6 to obtain the cost of the Miller loop and the final exponentiation. The Miller loop costs in this case  $4(9M_1 + 1I_9 + 2S_9 + 3M_9) + 51(9M_1 + 1I_9 + 2S_9 + 3M_9) + 50S_{27} + 53M_{27}$ . This is equal to  $55I_9 + 53M_{27} + 165M_9 + 495M_1 + 50S_{27} + 110S_9$ . Using the arithmetic in Table 2, the overall cost is  $55I_1 + 21348M_1 + 9660S_1$ . Using the value of  $x$  given above, the final exponentiation costs  $I_{27} + 2(51S_{27} + 3M_{27}) + 17(51S_{27} + 4M_{27}) + 11M_{27} + 2I_{G_{\phi_3(p^9)}}$  and  $p, p^2, p^3, p^4, p^5, p^6, p^7, p^8, 2*p^9$ -Frobenius maps for a total cost of the final exponentiation  $I_1 + 19191M_1 + 122337S_1$ .

The cost for the case  $k = 24$  are obtained with the parameter given in [4] and the formulas from [2]

The following table 7 compares our results with previous results at the 256-bit security level.

Curves-Ref.	Miller loop	Final Exp.	Size of $p$	Total( $S_1 = M_1$ )
$k = 27$ (This work)	$55I_1 + 21348M_1$ $+9660S_1$	$I_1 + 19191M_1$ $+122337S_1$	1019	$56I_1 + 172536M_1$
BLS24[4,2]	$18812M_1$	$10I_1 + 43142M_1$	1029	$10I_1 + 61954M_1$
KSS18[4]	$32238M_1 + 2620S_1$	$8I_1 + 7977M_1$ $+39520S_1$	1495	$8I_1 + 82355M_1$

**Table 7** Comparison of the cost of the Miller loop and the final exponentiation at 256-bit security level.

## 8.4 Comparison

To make a fair comparison of the results in tables 5, 6 and 7, we take note of the size of the base field. We consider implementations on 64-bit platform. Then following [2], a  $\mathbb{F}_p$  element is represented with  $\ell = 1 + \log_2(p)$  binary coefficients packed in  $n_{64} = \lceil \frac{\ell}{64} \rceil$  64 bits processor words and a  $\mathbb{F}_p$  multiplication can be implemented with approximately  $2n_{64}^2 + n_{64}$  operations. We denote  $m_c$  the cost of a multiplication in the finite field  $\mathbb{F}_p$  where  $p$  is of  $c$  bits. For the Table 5 we have that  $m_{461} \approx 1.35m_{371}$ . From this we see that, at the 128-bit security level, the total cost of computing the optimal ate pairing for elliptic curves with  $k = 15$  is  $19199m_{461}$  making these curves faster than the well known BN curves but slower than the KSS16 curves found in [4] as the best one at the 128-bit security level. From Table 6 we have that the cost for computing the optimal ate pairing for curves with  $k = 15$  is  $166067m_{511}$  as  $m_{863} \approx 2.98m_{511}$ . We conclude that at the 192-bit security level computing the optimal ate pairing is faster on elliptic curves with embedding degree  $k = 27$  than on curves with  $k = 15$  and in this case the BLS24 curves remain faster. At the 256-bit security level, we have that the BLS24 curves are the faster.

## 9 Conclusion

In this work we provided details and important improvements in the computation of the Miller loop and the final exponentiation for the optimal ate pairing on elliptic curves admitting cubic twists. An explicit cost evaluation is given for the Miller loop in the case of elliptic curves of embedding degree 9 and 15. In particular for  $k = 15$  and  $k = 27$  we obtained an improvement, in terms of operations in the base field, of up to 25% and 29% respectively in the computation of the final exponentiation. Also, we obtained that elliptic curves with embedding degree  $k = 15$  present faster results than BN12 curves at the 128-bit security levels. In comparison with curves having even embedding degrees we find that a lot of improvements are still required in the case of curves with odd embedding degree. One could try to compute compressed squarings in this case. To ensure the correctness of all the formulas used in this work, a MAGMA code for the implementation of the three pairings is given. Furthermore, a brief look at the parameters used in this work reveals that the curves with odd embedding degrees studied in this work are not *sub-*

group secure ordinary curves [6] and are not protected against *small-subgroup attacks*[32]. However this is not a particular case of elliptic curves of odd embedding degree but it appears from [6] that most of such parameters that have been found for curves with even embedding degree such as BN12 curves [8], KSS16 curves [24] or BLS12 curves [7] do not satisfied these security properties. As future work we could search for parameters to fulfill this security issue.

## References

1. Diego F. Aranha, Laura Fuentes-Castañeda, Edward Knapp, Alfred Menezes, and Francisco Rodríguez-Henríquez. Implementing pairings at the 192-bit security level. In *Pairing-Based Cryptography - Pairing 2012 - 5th International Conference, Cologne, Germany, May 16-18, 2012, Revised Selected Papers*, pages 177–195, 2012.
2. Diego F. Aranha, Laura Fuentes-Castañeda, Edward Knapp, Alfred Menezes, and Francisco Rodríguez-Henríquez. Implementing pairings at the 192-bit security level. In *Pairing-Based Cryptography - Pairing 2012 - 5th International Conference, Cologne, Germany, May 16-18, 2012, Revised Selected Papers*, pages 177–195, 2012.
3. R. Avanzi, H. Cohen, C. Doche, G. Frey, T. Lange, K. Nguyen, and F. Vercauteren. *Handbook of Elliptic and Hyperelliptic curve Cryptography*. Discrete Math. Appli. Chapman and Hall, 2006.
4. Razvan Barbulescu and Sylvain Duquesne. Updating key size estimations for pairings. *Journal of Cryptology*, 2018.
5. Razvan Barbulescu, Pierrick Gaudry, Aurore Guillevic, and François Morain. Improving NFS for the discrete logarithm problem in non-prime finite fields. In *Advances in Cryptology - EUROCRYPT 2015 - 34th Annual International Conference on the Theory and Applications of Cryptographic Techniques, Sofia, Bulgaria, April 26-30, 2015, Proceedings, Part I*, pages 129–155, 2015.
6. Paulo S. L. M. Barreto, Craig Costello, Rafael Misoczki, Michael Naehrig, Geovandro C. C. F. Pereira, and Gustavo Zanon. Subgroup security in pairing-based cryptography. In *Progress in Cryptology - LATINCRYPT 2015 - 4th International Conference on Cryptology and Information Security in Latin America, Guadalajara, Mexico, August 23-26, 2015, Proceedings*, pages 245–265, 2015.
7. Paulo S. L. M. Barreto, Ben Lynn, and Michael Scott. Constructing elliptic curves with prescribed embedding degrees. In *Security in Communication Networks, Third International Conference, SCN 2002, Amalfi, Italy, September 11-13, 2002. Revised Papers*, pages 257–267, 2002.
8. Paulo S. L. M. Barreto and Michael Naehrig. Pairing-friendly elliptic curves of prime order. In *Selected Areas in Cryptography, 12th International Workshop, SAC 2005, Kingston, ON, Canada, August 11-12, 2005, Revised Selected Papers*, pages 319–331, 2005.
9. I.F. Blake, G. Seroussi, and N.P. Smart. *Advances in Elliptic Curves in Cryptography*. London Mathematic Society, Cambridge University Press, (2005).
10. Dan Boneh and Matthew K. Franklin. Identity-based encryption from the Weil pairing. In *Advances in Cryptology - CRYPTO 2001, 21st Annual International Cryptology Conference, Santa Barbara, California, USA, August 19-23, 2001, Proceedings*, pages 213–229, 2001.
11. Dan Boneh, Ben Lynn, and Hovav Shacham. Short signatures from the weil pairing. *J. Cryptology*, 17(4):297–319, 2004.
12. Clifford Cocks. An identity based encryption scheme based on quadratic residues. In *Cryptography and Coding, 8th IMA International Conference, Cirencester, UK, December 17-19, 2001, Proceedings*, pages 360–363, 2001.
13. Craig Costello, Tanja Lange, and Michael Naehrig. Faster pairing computations on curves with high-degree twists. In *Public Key Cryptography - PKC 2010, 13th International Conference on Practice and Theory in Public Key Cryptography, Paris, France, May 26-28, 2010. Proceedings*, pages 224–242, 2010.

14. Augusto Jun Devegili, Michael Scott, and Ricardo Dahab. Implementing cryptographic pairings over barreto-naehrig curves. In *Pairing-Based Cryptography - Pairing 2007, First International Conference, Tokyo, Japan, July 2-4, 2007, Proceedings*, pages 197–207, 2007.
15. Pu Duan, Shi Cui, and Choong Wah Chan. Special polynomial families for generating more suitable elliptic curves for pairing-based cryptosystems. *IACR Cryptology ePrint Archive*, 2005:342, 2005.
16. Ratna Dutta, Rana Barua, and Palash Sarkar. Pairing-based cryptographic protocols : A survey. *IACR Cryptology ePrint Archive*, 2004:64, 2004.
17. David Freeman, Michael Scott, and Edlyn Teske. A taxonomy of pairing-friendly elliptic curves. *J. Cryptology*, 23(2):224–280, 2010.
18. Laura Fuentes-Castañeda, Edward Knapp, and Francisco Rodríguez-Henríquez. Faster hashing to  $\mathbb{G}_2$ . In *Selected Areas in Cryptography - 18th International Workshop, SAC 2011, Toronto, ON, Canada, August 11-12, 2011, Revised Selected Papers*, pages 412–430, 2011.
19. Vipul Goyal, Omkant Pandey, Amit Sahai, and Brent Waters. Attribute-based encryption for fine-grained access control of encrypted data. In *Proceedings of the 13th ACM Conference on Computer and Communications Security, CCS 2006, Alexandria, VA, USA, Ioctober 30 - November 3, 2006*, pages 89–98, 2006.
20. Robert Granger and Michael Scott. Faster squaring in the cyclotomic subgroup of sixth degree extensions. In *Public Key Cryptography - PKC 2010, 13th International Conference on Practice and Theory in Public Key Cryptography, Paris, France, May 26-28, 2010. Proceedings*, pages 209–223, 2010.
21. Aurore Guillevic. Kim-barbulescu variant of the number field sieve to compute discrete logarithms in finite fields. In <https://ellipticnews.wordpress.com/2016/05/02/kim-barbulescu-variant-of-the-number-field-sieve-to-compute-discrete-logarithms-in-finite-fields>, ellipticnews, May 2016.
22. Florian Hess. Pairing lattices. In *Pairing-Based Cryptography - Pairing 2008, Second International Conference, Egham, UK, September 1-3, 2008. Proceedings*, pages 18–38, 2008.
23. Florian Hess, Nigel P. Smart, and Frederik Vercauteren. The eta pairing revisited. *IEEE Transactions on Information Theory*, 52(10):4595–4602, 2006.
24. Ezekiel J. Kachisa, Edward F. Schaefer, and Michael Scott. Constructing brezing-weng pairing friendly elliptic curves using elements in the cyclotomic field. *IACR Cryptology ePrint Archive*, 2007:452, 2007.
25. Koray Karabina. Squaring in cyclotomic subgroups. *Math. Comput.*, 82(281), 2013.
26. Taechan Kim and Razvan Barbulescu. Extended tower number field sieve: A new complexity for the medium prime case. In *CRYPTO (1)*, volume 9814 of *Lecture Notes in Computer Science*, pages 543–571. Springer, 2016.
27. Neal Koblitz and Alfred Menezes. Pairing-based cryptography at high security levels. In *Cryptography and Coding, 10th IMA International Conference, Cirencester, UK, December 19-21, 2005, Proceedings*, pages 13–36, 2005.
28. Kristin E. Lauter, Peter L. Montgomery, and Michael Naehrig. An analysis of affine coordinates for pairing computation. In *Pairing-Based Cryptography - Pairing 2010 - 4th International Conference, Yamanaka Hot Spring, Japan, December 2010. Proceedings*, pages 1–20, 2010.
29. Duc-Phong Le and Chik How Tan. Speeding up ate pairing computation in affine coordinates. In *Information Security and Cryptology - ICISC 2012 - 15th International Conference, Seoul, Korea, November 28-30, 2012, Revised Selected Papers*, pages 262–277, 2012.
30. Benoît Libert and Jean-Jacques Quisquater. Identity based undeniable signatures. In *Topics in Cryptology - CT-RSA 2004, The Cryptographers' Track at the RSA Conference 2004, San Francisco, CA, USA, February 23-27, 2004, Proceedings*, pages 112–125, 2004.
31. R Lidl and H Niederreiter. *Finite Fields*. Cambridge university Press, Cambridge, 1997.
32. Chae Hoon Lim and Pil Joong Lee. A key recovery attack on discrete log-based schemes using a prime order subgroup. In *Advances in Cryptology - CRYPTO '97, 17th Annual International Cryptology Conference, Santa Barbara, California, USA, August 17-21, 1997, Proceedings*, pages 249–263, 1997.

33. Xibin Lin, Changan Zhao, Fangguo Zhang, and Yanming Wang. Computing the ate pairing on elliptic curves with embedding degree  $k = 9$ . *IEICE Transactions*, 91-A(9):2387–2393, 2008.
34. Alfred Menezes, Palash Sarkar, and Shashank Singh. Challenges with assessing the impact of nfs advances on the security of pairing-based cryptography. *Cryptology ePrint Archive*, Report 2016/1102, 2016.
35. Victor S. Miller. The weil pairing, and its efficient calculation. *J. Cryptology*, 17(4):235–261, 2004.
36. Nadia El Mrabet, Nicolas Guillermine, and Sorina Ionica. A study of pairing computation for elliptic curves with embedding degree 15. *IACR Cryptology ePrint Archive*, 2009:370, 2009.
37. Michael Scott, Naomi Benger, Manuel Charlemagne, Luis J. Dominguez Perez, and Ezekiel J. Kachisa. Fast hashing to  $\mathbb{G}_2$  on pairing-friendly curves. In *Pairing-Based Cryptography - Pairing 2009, Third International Conference, Palo Alto, CA, USA, August 12-14, 2009, Proceedings*, pages 102–113, 2009.
38. Michael Scott and Aurore Guillevic. A new family of pairing-friendly elliptic curves. *Cryptology ePrint Archive*, Report 2018/193, 2018.
39. Frederik Vercauteren. Optimal pairings. *IEEE Transactions on Information Theory*, 56(1):455–461, 2010.
40. L.C. Washington. *Elliptic Curves, Number Theory and Cryptography*. Discrete Math .Appli, Chapman and Hall, 2008.
41. Xusheng Zhang and Dongdai Lin. Analysis of optimum pairing products at high security levels. In *Progress in Cryptology - INDOCRYPT 2012, 13th International Conference on Cryptology in India, Kolkata, India, December 9-12, 2012. Proceedings*, pages 412–430, 2012.

## A Arithmetic in $\mathbb{F}_{p^9}$

Let  $a = a_0 + a_1v + a_2v^2 \in \mathbb{F}_{p^9}$  with  $a_i \in \mathbb{F}_{p^3}$ .

### A.1 Cyclotomic inversion

We assume that  $a$  lies in the cyclotomic subgroup  $G_{\phi_3(p^3)}$ , so that  $a^{p^6+p^3+1} = 1$  i.e.  $a^{-1} = a^{p^6}a^{p^3}$ . In order to compute  $a^{p^6}a^{p^3}$ , we need the values of  $v^{p^3}$  and  $v^{p^6}$ . But  $v^{p^3} = v^{3(p^3-1)/3+1} = v^{3(p^3-1)/3}v = (v^3)^{(p^3-1)/3}v = (7^{1/3})^{(p^3-1)/3}v$  since  $v^3 = 7^{1/3}$ .

Let  $\mu = (7^{1/3})^{(p^3-1)/3}$ , we have  $\mu \neq 1$  and  $\mu^3 = 1$  so that  $\mu$  is a primitive cubic root of unity in  $\mathbb{F}_{p^3}$ . We obtain  $v^{p^3} = \mu v$  and  $v^{p^6} = (v^{p^3})^{p^3} = (\mu v)^{p^3} = \mu(v)^{p^3} = \mu\mu v = \mu^2v$ . We then have  $a^{p^3} = a_0^{p^3} + a_1^{p^3}v^{p^3} + a_2^{p^3}(v^2)^{p^3} = a_0 + a_1v^{p^3} + a_2(v^2)^{p^3} = a_0 + a_1\mu v + a_2\mu^2v^2$  and  $a^{p^6} = (a^{p^3})^{p^3} = a_0 + a_1(\mu v)^{p^3} + a_2(\mu^2v^2)^{p^3} = a_0 + a_1\mu^2v + a_2\mu^4v^2$ . So that when using  $v^3 = 7^{1/3}$  and  $\phi_3(\mu) = \mu^2 + \mu + 1 = 0$ , we finally have:

$$a^{p^6}a^{p^3} = (a_0^2 - a_1a_27^{1/3}) + (a_2^27^{1/3} - a_0a_1)v + (a_1^2 - a_0a_2)v^2$$

This costs  $3M_3 + 3S_3 = 18M_1 + 15S_1$  with additional additions.

### A.2 Frobenius operators

The  $p^i$ -Frobenius is the map  $\pi^i: \mathbb{F}_{p^9} \rightarrow \mathbb{F}_{p^9}, a \mapsto a^{p^i}$ . Let  $a \in \mathbb{F}_{p^9}$ ,  $a = a_0 + a_1v + a_2v^2$  with  $a_i \in \mathbb{F}_{p^3}$  then  $\pi(a) = a_0^p + a_1^pv^p + a_2^p(v^2)^p$ . Now  $a_0 \in \mathbb{F}_{p^3}$  can be written as  $a_0 = g_0 + g_1u + g_2u^2$ ,  $g_i \in \mathbb{F}_p$  so that  $a_0^p = g_0 + g_1u^p + g_2(u^2)^p$ .

We have  $u^p = u^{3(p-1)/3+1} = (u^3)^{(p-1)/3}u = 7^{(p-1)/3}u$  and since 7 is not a cube in  $\mathbb{F}_p$ ,  $7^{(p-1)/3} \neq 1$ . Let  $\alpha = 7^{(p-1)/3}$  then  $\alpha \neq 1$  and  $\alpha^3 = 1$ ; it means that  $\alpha$  is a primitive cubic root of unity in  $\mathbb{F}_p$  and  $u^p = \alpha u$ . Therefore  $a_0^p = g_0 + g_1 u^p + g_2 (u^2)^p = g_0 + g_1 \alpha u + g_2 \alpha^2 u^2$  and similarly  $a_1^p = g_3 + g_4 u^p + g_5 (u^2)^p = g_3 + g_4 \alpha u + g_5 \alpha^2 u^2$  and  $a_2^p = g_6 + g_7 u^p + g_8 (u^2)^p = g_6 + g_7 \alpha u + g_8 \alpha^2 u^2$ . Now for the computation of  $v^p$  observe that  $v^p = v^{3(p-1)/3+1} = (v^3)^{(p-1)/3}v = (7^{1/3})^{(p-1)/3}v = 7^{(p-1)/9}v$  so that if  $\beta = 7^{(p-1)/9}$  then we have  $\beta \neq 1$ ,  $\beta^3 = 7^{(p-1)/3} = \alpha \neq 1$ ,  $\beta^9 = 1$ . Thus  $\beta$  is a primitive ninth root of unity in  $\mathbb{F}_p$  and  $v^p = \beta v$ .

Finally  $a^p = g_0 + g_1 \alpha u + g_2 \alpha^2 u^2 + (g_3 \beta + g_4 \alpha \beta u + g_5 \alpha^2 \beta u^2)v + (g_6 \beta^2 + g_7 \alpha \beta^2 u + g_8 \alpha^2 \beta^2 u^2)v^2$  and the following algebraic relations:  $\alpha = \beta^3$ ,  $\alpha \beta = \beta^4$ ,  $\alpha \beta^2 = \beta^5$ ,  $\alpha^2 \beta = \beta^7$ ,  $\alpha^2 \beta^2 = \beta^8$  yield to  $a^p = (g_0 + g_1 \beta^3 u + g_2 \beta^6 u^2) + (g_3 \beta + g_4 \beta^4 u + g_5 \beta^7 u^2)v + (g_6 \beta^2 + g_7 \beta^5 u + g_8 \beta^8 u^2)v^2$ . The cost of  $p$ -Frobenius:  $8M_1$ . This is the same as the cost of  $p^2, p^4, p^5, p^7$  and  $p^8$ -Frobenius.

For the  $p^3$ -Frobenius operator, observe from A.1 that  $v^{p^3} = \mu v$ . Then

$$a^{p^3} = a_0 + a_1 \mu v + a_2 \mu^2 v^2 = (g_0 + g_1 u + g_2 u^2) + (g_3 + g_4 u + g_5 u^2)\mu v + (g_6 + g_7 u + g_8 u^2)\mu^2 v^2.$$

As  $t = \mu^2$  is precomputed; we finally have

$$a^{p^3} = (g_0 + g_1 u + g_2 u^2) + (g_3 \mu + g_4 \mu u + g_5 \mu u^2)v + (g_6 t + g_7 t u + g_8 t u^2)v^2.$$

The cost of  $p^3$ -Frobenius:  $6M_1$ . This is the same as the cost of  $p^6$ -Frobenius.

## B Arithmetic in $\mathbb{F}_{p^{27}}$

### B.1 Cyclotomic inversion

We follow the same procedure as in A.1. The element  $a = a_0 + a_1 w + a_2 w^2 \in \mathbb{F}_{p^{27}}$  with  $a_i \in \mathbb{F}_{p^9}$  in the cyclotomic subgroup  $G_{\phi_3(\mathbb{F}_{p^9})}$  satisfies  $a^{p^{18}+p^9+1} = 1$  so that  $a^{-1} = a^{p^{18}} a^{p^9}$ .

In order to compute  $a^{p^{18}} a^{p^9}$ , we need the values of  $w^{p^9}$  and  $w^{p^{18}}$ . We have

$$w^{p^9} = w^{3(p^9-1)/3+1} = w^3(w^9-1)/3w = (w^3)^{(p^9-1)/3}w = (7^{1/9})^{(p^9-1)/3}w \text{ since } w^3 = 7^{1/9}.$$

Let  $\sigma = (7^{1/9})^{(p^9-1)/3}$  then  $\sigma \neq 1$  and  $\sigma^3 = 1$ . Hence  $\sigma$  is a primitive cubic root of unity in  $\mathbb{F}_{p^9}$  i.e.  $\phi_3(\sigma) = 0$ . We obtain  $w^{p^9} = \sigma w$  and we now compute  $w^{p^{18}}$  as  $w^{p^{18}} = (w^{p^9})^{p^9} = (\sigma w)^{p^9} = \sigma(w)^{p^9} = \sigma \sigma w = \sigma^2 w$ .

$$a^{p^9} = a_0 + a_1 w^{p^9} + a_2 (w^2)^{p^9} = a_0 + a_1 \sigma w + a_2 \sigma^2 w^2 \text{ and}$$

$$a^{p^{18}} = (a^{p^9})^{p^9} = a_0 + a_1 (\sigma w)^{p^9} + a_2 (\sigma^2 w^2)^{p^9} = a_0 + a_1 \sigma^2 w + a_2 \sigma^4 w^2. \text{ After expanding and reducing using } w^3 = 7^{1/9} \text{ and } \phi_3(\sigma) = \sigma^2 + \sigma + 1 = 0 \text{ we obtain}$$

$$a^{p^{18}} a^{p^9} = (a_0^2 - a_1 a_2 7^{1/9}) + (a_2^2 7^{1/9} - a_0 a_1)w + (a_1^2 - a_0 a_2)w^2$$

The computation costs  $3(36M_1) + 3(25S_1) = 108M_1 + 75S_1$ .

### B.2 Frobenius operators

The  $p^i$ -Frobenius is the map  $\pi^i: \mathbb{F}_{p^{27}} \rightarrow \mathbb{F}_{p^{27}}, a \mapsto a^{p^i}$ . Let  $a = a_0 + a_1 w + a_2 w^2$  with  $a_i \in \mathbb{F}_{p^9}$  an element of  $\mathbb{F}_{p^{27}}$ .

$\pi(a) = a^p = (a_0 + a_1 w + a_2 w^2)^p = a_0^p + a_1^p w^p + a_2^p (w^2)^p$ . The element  $a_0 \in \mathbb{F}_{p^9}$  can be written as  $a_0 = (h_0 + h_1 u + h_2 u^2) + (h_3 + h_4 u + h_5 u^2)v + (h_6 + h_7 u + h_8 u^2)v^2$ ,  $h_i \in \mathbb{F}_p$ .

We have  $a_0^p = (h_0 + h_1 u + h_2 u^2) + (h_3 + h_4 u + h_5 u^2)v + (h_6 + h_7 u + h_8 u^2)v^2$ ,  $h_i^p = h_i$ .

$u^p = u^{3(p-1)/3+1} = (u^3)^{(p-1)/3}u = 7^{(p-1)/3}u$ . Since 7 is not a cube in  $\mathbb{F}_p$ , we have  $\alpha = 7^{(p-1)/3}$   $\alpha \neq 1$  and  $\alpha^3 = 1$ . It means that  $\alpha$  is a primitive cubic root of unity in  $\mathbb{F}_p$  and  $u^p = \alpha u$ .  $v^p = v^{3(p-1)/3+1} = (v^3)^{(p-1)/3}v = (7^{1/3})^{(p-1)/3}v = 7^{(p-1)/9}v$ .

We have  $\beta = 7^{(p-1)/9} \neq 1$  and  $\beta^9 = 1$ . Thus  $\beta$  is a primitive ninth root of unity in  $\mathbb{F}_p$  and  $v^p = \beta v$ . Also  $w^p = w^{3(p-1)/3+1} = (w^3)^{(p-1)/3}w = (7^{1/9})^{(p-1)/3}w = 7^{(p-1)/27}w$ . We also observe that  $\gamma = 7^{(p-1)/27} \neq 1$ ,  $\gamma^3 = 7^{(p-1)/9} = \beta \neq 1$ ,  $\gamma^9 = 7^{(p-1)/3} = \alpha \neq 1$ ,  $\gamma^{27} = 1$ .

Thus  $\gamma$  is a primitive twenty-seventh root of unity in  $\mathbb{F}_p$  and  $w^p = \gamma w$ .

$$\begin{aligned}
a_0^p &= ((h_0 + h_1u + h_2u^2) + (h_3 + h_4u + h_5u^2)v + (h_6 + h_7u + h_8u^2)v^2)^p \\
&= (h_0 + h_1u^p + h_2(u^2)^p) + (h_3 + h_4u^p + h_5(u^2)^p)v^p + (h_6 + h_7u^p + h_8(u^2)^p)(v^2)^p \\
&= (h_0 + h_1\alpha u + h_2\alpha^2 u^2) + (h_3 + h_4\alpha u + h_5\alpha^2 u^2)\beta v + (h_6 + h_7\alpha u + h_8\alpha^2 u^2)\beta^2 v^2 \\
&= (h_0 + h_1\alpha u + h_2\alpha^2 u^2) + (h_3\beta + h_4\alpha\beta u + h_5\alpha^2\beta u^2)v + (h_6\beta^2 + h_7\alpha\beta^2 u + h_8\alpha^2\beta^2 u^2)v^2. \\
a_1^p &= (h_9 + h_{10}u + h_{11}u^2) + (h_{12} + h_{13}u + h_{14}u^2)v + (h_{15} + h_{16}u + h_{17}u^2)v^2 \\
&= (h_9 + h_{10}u^p + h_{11}(u^2)^p) + (h_{12} + h_{13}u^p + h_{14}(u^2)^p)v^p + (h_{15} + h_{16}u^p + h_{17}(u^2)^p)(v^2)^p \\
&= (h_9 + h_{10}\alpha u + h_{11}\alpha^2 u^2) + (h_{12} + h_{13}\alpha u + h_{14}\alpha^2 u^2)\beta v + (h_{15} + h_{16}\alpha u + h_{17}\alpha^2 u^2)\beta^2 v^2 \\
&= (h_9 + h_{10}\alpha u + h_{11}\alpha^2 u^2) + (h_{12}\beta + h_{13}\alpha\beta u + h_{14}\alpha^2\beta u^2)v + (h_{15}\beta^2 + h_{16}\alpha\beta^2 u + h_{17}\alpha^2\beta^2 u^2)v^2. \\
a_2^p &= (h_{18} + h_{19}u + h_{20}u^2) + (h_{21} + h_{22}u + h_{23}u^2)v + (h_{24} + h_{25}u + h_{26}u^2)v^2 \\
&= (h_{18} + h_{19}u^p + h_{20}(u^2)^p) + (h_{21} + h_{22}u^p + h_{23}(u^2)^p)v^p + (h_{24} + h_{25}u^p + h_{26}(u^2)^p)(v^2)^p \\
&= (h_{18} + h_{19}\alpha u + h_{20}\alpha^2 u^2) + (h_{21} + h_{22}\alpha u + h_{23}\alpha^2 u^2)\beta v + (h_{24} + h_{25}\alpha u + h_{26}\alpha^2 u^2)\beta^2 v^2 \\
&= (h_{18} + h_{19}\alpha u + h_{20}\alpha^2 u^2) + (h_{21}\beta + h_{22}\alpha\beta u + h_{23}\alpha^2\beta u^2)v + (h_{24}\beta^2 + h_{25}\alpha\beta^2 u + h_{26}\alpha^2\beta^2 u^2)v^2. \\
\pi(a) &= (a_0 + a_1w + a_2w^2)^p = a_0^p + a_1^p w^p + a_2^p (w^2)^p = a_0^p + a_1^p \gamma w + a_2^p \gamma^2 w^2 \\
&= (h_0 + h_1\alpha u + h_2\alpha^2 u^2) + (h_3\beta + h_4\alpha\beta u + h_5\alpha^2\beta u^2)v + (h_6\beta^2 + h_7\alpha\beta^2 u + h_8\alpha^2\beta^2 u^2)v^2 + \\
&\quad ((h_9 + h_{10}\alpha u + h_{11}\alpha^2 u^2) + (h_{12}\beta + h_{13}\alpha\beta u + h_{14}\alpha^2\beta u^2)v + (h_{15}\beta^2 + h_{16}\alpha\beta^2 u + h_{17}\alpha^2\beta^2 u^2)v^2)\gamma w + \\
&\quad ((h_{18} + h_{19}\alpha u + h_{20}\alpha^2 u^2) + (h_{21}\beta + h_{22}\alpha\beta u + h_{23}\alpha^2\beta u^2)v + (h_{24}\beta^2 + h_{25}\alpha\beta^2 u + h_{26}\alpha^2\beta^2 u^2)v^2)\gamma^2 w^2.
\end{aligned}$$

We have these following algebraic relations:  $\alpha = \beta^3$ ,  $\alpha\beta = \beta^4$ ,  $\alpha\beta^2 = \beta^5$ ,  $\alpha^2\beta = \beta^7$  and  $\alpha^2\beta^2 = \beta^8$ . Therefore

$$\begin{aligned}
\pi(a) &= ((h_0 + h_1\beta^3 u + h_2\beta^6 u^2) + (h_3\beta + h_4\beta^4 u + h_5\beta^7 u^2)v + (h_6\beta^2 + h_7\beta^5 u + h_8\beta^8 u^2)v^2) + \\
&\quad ((h_9\gamma + h_{10}\beta^3\gamma u + h_{11}\beta^6\gamma u^2) + (h_{12}\beta\gamma + h_{13}\beta^4\gamma u + h_{14}\beta^7\gamma u^2)v + (h_{15}\beta^2\gamma + h_{16}\beta^5\gamma u + h_{17}\beta^8\gamma u^2)v^2)w + \\
&\quad ((h_{18}\gamma^2 + h_{19}\beta^3\gamma^2 u + h_{20}\beta^6\gamma^2 u^2) + (h_{21}\beta\gamma^2 + h_{22}\beta^4\gamma^2 u + h_{23}\beta^7\gamma^2 u^2)v + \\
&\quad (h_{24}\beta^2\gamma^2 + h_{25}\beta^5\gamma^2 u + h_{26}\beta^8\gamma^2 u^2)v^2)w^2.
\end{aligned}$$

The following values are precomputed:  $\lambda_0 = \beta^2$ ,  $\lambda_1 = \beta^3$ ,  $\lambda_2 = \beta^4$ ,  $\lambda_3 = \beta^5$ ,  $\lambda_4 = \beta^6$ ,  $\lambda_5 = \beta^7$ ,  $\lambda_6 = \beta^8$ ,  $\lambda_7 = \gamma^2$ ,  $\lambda_8 = \beta\gamma$ ,  $\lambda_9 = \lambda_0\gamma$ ,  $\lambda_{10} = \lambda_1\gamma$ ,  $\lambda_{11} = \lambda_2\gamma$ ,  $\lambda_{12} = \lambda_3\gamma$ ,  $\lambda_{13} = \lambda_4\gamma$ ,  $\lambda_{14} = \lambda_5\gamma$ ,  $\lambda_{15} = \lambda_6\gamma$ ,  $\lambda_{16} = \lambda_0\lambda_7$ ,  $\lambda_{17} = \lambda_1\lambda_7$ ,  $\lambda_{18} = \lambda_2\lambda_7$ ,  $\lambda_{19} = \lambda_3\lambda_7$ ,  $\lambda_{20} = \lambda_4\lambda_7$ ,  $\lambda_{21} = \lambda_5\lambda_7$ ,  $\lambda_{22} = \lambda_6\lambda_7$ .  $\lambda_{23} = \beta\lambda_7$ .

$$\begin{aligned}
\text{Thus } \pi(a) &= ((h_0 + h_1\lambda_1 u + h_2\lambda_4 u^2) + (h_3\beta + h_4\lambda_2 u + h_5\lambda_5 u^2)v + (h_6\lambda_0 + h_7\lambda_3 u + h_8\lambda_6 u^2)v^2) + \\
&\quad (h_9\lambda_7 + h_{10}\lambda_1\lambda_7 u + h_{11}\lambda_4\lambda_7 u^2) + (h_{12}\lambda_8 + h_{13}\lambda_{11} u + h_{14}\lambda_{14} u^2)v + \\
&\quad (h_{15}\lambda_9 + h_{16}\lambda_{12} u + h_{17}\lambda_{15} u^2)v^2)w + ((h_{18}\lambda_7 + h_{19}\lambda_{17} u + h_{20}\lambda_{20} u^2) + (h_{21}\lambda_{23} + h_{22}\lambda_{18} u + h_{23}\lambda_{21} u^2)v + \\
&\quad (h_{24}\lambda_{16} + h_{25}\lambda_{19} u + h_{26}\lambda_{22} u^2)v^2)w^2.
\end{aligned}$$

The cost of  $p$ -Frobenius:  $26M_1 + 18a$ . This is also equal to the cost of  $p^2, p^4, p^5, p^7, p^8$  Frobenius

For the  $p^9$  Frobenius operator, observe from B.1 that  $w^{p^9} = \sigma w$ . Then

$$\begin{aligned}
a^{p^9} &= a_0 + a_1\sigma w + a_2\sigma^2 w^2 = ((h_0 + h_1u + h_2u^2) + (h_3 + h_4u + h_5u^2)v + (h_6 + h_7u + h_8u^2)v^2) + \\
&\quad ((h_9 + h_{10}u + h_{11}u^2) + (h_{12} + h_{13}u + h_{14}u^2)v + (h_{15} + h_{16}u + h_{17}u^2)v^2)\sigma w + \\
&\quad ((h_{18} + h_{19}u + h_{20}u^2) + (h_{21} + h_{22}u + h_{23}u^2)v + (h_{24} + h_{25}u + h_{26}u^2)v^2)\sigma^2 w^2.
\end{aligned}$$

$$\begin{aligned}
\text{We then have } a^{p^9} &= ((h_0 + h_1u + h_2u^2) + (h_3 + h_4u + h_5u^2)v + (h_6 + h_7u + h_8u^2)v^2) + \\
&\quad ((h_9\sigma + h_{10}\sigma u + h_{11}\sigma u^2) + (h_{12}\sigma + h_{13}\sigma u + h_{14}\sigma u^2)v + (h_{15}\sigma + h_{16}\sigma u + h_{17}\sigma u^2)v^2)w + \\
&\quad ((h_{18}\sigma^2 + h_{19}\sigma^2 u + h_{20}\sigma^2 u^2) + (h_{21}\sigma^2 + h_{22}\sigma^2 u + h_{23}\sigma^2 u^2)v + (h_{24}\sigma^2 + h_{25}\sigma^2 u + h_{26}\sigma^2 u^2)v^2)w^2.
\end{aligned}$$

As  $s = \sigma^2$  is precomputed; we have  $a^{p^9} = ((h_0 + h_1u + h_2u^2) + (h_3 + h_4u + h_5u^2)v + (h_6 + h_7u + h_8u^2)v^2) + ((h_9\sigma + h_{10}\sigma u + h_{11}\sigma u^2) + (h_{12}\sigma + h_{13}\sigma u + h_{14}\sigma u^2)v + (h_{15}\sigma + h_{16}\sigma u + h_{17}\sigma u^2)v^2)w + ((h_{18}s + h_{19}s u + h_{20}s u^2) + (h_{21}s + h_{22}s u + h_{23}s u^2)v + (h_{24}s + h_{25}s u + h_{26}s u^2)v^2)w^2$ .

The cost of  $p^9$ -Frobenius:  $18M_1$ . This is the same as the cost of  $p^3$  and  $p^6$  Frobenius.

## C Arithmetic in $\mathbb{F}_{p^{15}}$

The arithmetic of the extension field  $F_{p^5}$  is studied in [36]. In this section we only consider inversion in cyclotomic subgroup and Frobenius operators.

### C.1 Cyclotomic inversion

An element  $a = a_0 + a_1v + a_2v^2 \in \mathbb{F}_{p^{15}}$  with  $a_i \in \mathbb{F}_{p^5}$  in the cyclotomic subgroup  $G_{\phi_3(p^5)}$  satisfies  $a^{p^{10}+p^5+1} = 1$  so that  $a^{-1} = a^{p^{10}} a^{p^5}$ .  
 $v^{p^5} = v^{5(p^5-1)/5+1} = v^{5(p^5-1)/5}v = (v^5)^{(p^5-1)/5}v = (7^{1/3})^{(p^5-1)/5}v$  since  $v^5 = 7^{1/3}$ .

Let  $\omega = (7^{1/3})^{(p^5-1)/5}$ . We have  $\omega \neq 1$  and  $\omega^5 = 1$ . Hence  $\omega$  is a primitive fifth root of unity in  $\mathbb{F}_{p^5}$ . We obtain  $v^{p^5} = \omega v$ , and  $v^{p^{10}} = (v^{p^5})^{p^5} = (\omega v)^{p^5} = \omega(v)^{p^5} = \omega\omega v = \omega^2v$ .

$$a^{p^5} = (a_0 + a_1v + a_2v^2)^{p^5} = a_0^{p^5} + a_1^{p^5}v^{p^5} + a_2^{p^5}(v^2)^{p^5} = a_0 + a_1v^{p^5} + a_2(v^2)^{p^5} = a_0 + a_1\omega v + a_2\omega^2v^2.$$

$$a^{p^{10}} = (a^{p^5})^{p^5} = a_0 + a_1(\omega v)^{p^5} + a_2(\omega^2v^2)^{p^5} = a_0 + a_1\omega^2v + a_2\omega^4v^2.$$

$$a^{p^{10}} a^{p^5} = (a_0 + a_1\omega^2v + a_2\omega^4v^2)(a_0 + a_1\omega v + a_2\omega^2v^2)$$

After expanding and reducing using  $v^3 = u$  and  $\phi_5(\omega) = 0$  we obtain

$$a^{p^{10}} a^{p^5} = (a_0^2 + (1 + \omega^4)a_1a_2u) + \omega(a_2^2u + (1 + \omega)a_0a_1)v + \omega^2(a_1^2\omega + (1 + \omega^2)a_0a_2)v^2$$

This costs  $3(9M_1) + 3(9S_1) = 27M_1 + 27S_1$

### C.2 Frobenius operators

The  $p^i$ -Frobenius is the map  $\pi^i: \mathbb{F}_{p^{15}} \rightarrow \mathbb{F}_{p^{15}}, a \mapsto a^{p^i}$ . Let  $a \in \mathbb{F}_{p^{15}}; a = a_0 + a_1v + a_2v^2$  with  $a_i \in \mathbb{F}_{p^5}$ .  $\pi(a) = a^p = (a_0 + a_1v + a_2v^2)^p = a_0^p + a_1^p v^p + a_2^p (v^2)^p$ . As  $a_0 \in \mathbb{F}_{p^5}$  i.e.  $a_0 = g_0 + g_1u + g_2u^2 + g_3u^3 + g_4u^4$ ,  $g_i \in \mathbb{F}_p$ , we have  $a_0^p = (g_0 + g_1u + g_2u^2 + g_3u^3 + g_4u^4)^p = g_0 + g_1u^p + g_2(u^2)^p + g_3(u^3)^p + g_4(u^4)^p$ , since  $g_i^p = g_i$ .

Now we have  $u^p = u^{5(p-1)/5+1} = (u^5)^{(p-1)/5}u = 7^{(p-1)/5}u$  and as 7 is not a fifth power in  $\mathbb{F}_p$ ; so  $7^{(p-1)/5} \neq 1$ . Let  $\theta = 7^{(p-1)/5}$ ,  $\theta \neq 1$  and  $\theta^5 = 1$ . It means that  $\theta$  is a primitive fifth root of unity in  $\mathbb{F}_p$  and  $u^p = \theta u$ .  $a_0^p = g_0 + g_1u^p + g_2(u^2)^p + g_3(u^3)^p + g_4(u^4)^p = g_0 + g_1\theta u + g_2\theta^2u^2 + g_3\theta^3u^3 + g_4\theta^4u^4$ .

$$a_1^p = g_5 + g_6u^p + g_7(u^2)^p + g_8(u^3)^p + g_9(u^4)^p = g_5 + g_6\theta u + g_7\theta^2u^2 + g_8\theta^3u^3 + g_9\theta^4u^4.$$

$$a_2^p = g_{10} + g_{11}u^p + g_{12}(u^2)^p + g_{13}(u^3)^p + g_{14}(u^4)^p = g_{10} + g_{11}\theta u + g_{12}\theta^2u^2 + g_{13}\theta^3u^3 + g_{14}\theta^4u^4.$$

$v^p = v^{5(p-1)/5+1} = (v^5)^{(p-1)/5}v = (7^{1/3})^{(p-1)/5}v = (7^{1/3})^{(p-1)/5}v$ .  $7^{1/3}$  is not a fifth power in  $\mathbb{F}_p$ ; so  $(7^{1/3})^{(p-1)/5} \neq 1$ .

Set  $\beta = (7^{1/3})^{(p-1)/5}$ , we have  $\beta \neq 1$ ;  $\beta^5 = 1$ . Thus  $\beta$  is a primitive fifth root of unity in  $\mathbb{F}_p$  and  $v^p = \beta v$ .  $a^p = (a_0 + a_1v + a_2v^2)^p = a_0^p + a_1^p v^p + a_2^p (v^2)^p = (g_0 + g_1\theta u + g_2\theta^2u^2 + g_3\theta^3u^3 + g_4\theta^4u^4) + (g_5 + g_6\theta u + g_7\theta^2u^2 + g_8\theta^3u^3 + g_9\theta^4u^4)v^p + (g_{10} + g_{11}\theta u + g_{12}\theta^2u^2 + g_{13}\theta^3u^3 + g_{14}\theta^4u^4)(v^2)^p = (g_0 + g_1\theta u + g_2\theta^2u^2 + g_3\theta^3u^3 + g_4\theta^4u^4) + (g_5\beta + g_6\theta\beta u + g_7\theta^2\beta u^2 + g_8\theta^3\beta u^3 + g_9\theta^4\beta u^4)v + (g_{10}\beta^2 + g_{11}\theta\beta^2u + g_{12}\theta^2\beta^2u^2 + g_{13}\theta^3\beta^2u^3 + g_{14}\theta^4\beta^2u^4)v^2$ . We precomputed these following values:  $c_0 = \theta^2$ ,  $c_1 = \theta^3$ ,  $c_2 = \theta^4$ ,  $c_3 = \beta^2$ ,  $c_4 = \theta\beta$ ,  $c_5 = c_0\beta$ ,  $c_6 = c_1\beta$ ,  $c_7 = c_2\beta$ ,  $c_8 = \theta c_3$ ,  $c_9 = c_0c_3$ ,  $c_{10} = c_1c_3$ ,  $c_{11} = c_2c_3$ . So  $\pi(a) = (g_0 + g_1\theta u + g_2c_0u^2 + g_3c_1u^3 + g_4c_2u^4) + (g_5\beta + g_6c_4u + g_7c_5u^2 + g_8c_6u^3 + g_9c_7u^4)v + (g_{10}c_3 + g_{11}c_8u + g_{12}c_9u^2 + g_{13}c_{10}u^3 + g_{14}c_{11}u^4)v^2$ .

The cost of  $p$ -Frobenius is  $14M_1$ . This is the same cost as computing  $p^2, p^3, p^4, p^6, p^7, p^8, p^9$  Frobenius.

For the  $p^5$  Frobenius operator, observe from C.1 that  $v^{p^5} = \omega v$  so that  $a^{p^5} = (g_0 + g_1u + g_2u^2 + g_3u^3 + g_4u^4) + (g_5 + g_6u + g_7u^2 + g_8u^3 + g_9u^4)v^{p^5} + (g_{10} + g_{11}u + g_{12}u^2 + g_{13}u^3 + g_{14}u^4)(v^{p^5})^2$ .

$$= (g_0 + g_1u + g_2u^2 + g_3u^3 + g_4u^4) + (g_5\omega + g_6\omega u + g_7\omega u^2 + g_8\omega u^3 + g_9\omega u^4)v$$

$$+(g_{10}\omega^2 + g_{11}\omega^2u + g_{12}\omega^2u^2 + g_{13}\omega^2u^3 + g_{14}\omega^2u^4)v^2.$$

We precomputed  $d = \omega^2$ .

$\pi^5(a) = a^{p^5} = (g_0 + g_1u + g_2u^2 + g_3u^3 + g_4u^4) + (g_5\omega + g_6\omega u + g_7\omega u^2 + g_8\omega u^3 + g_9\omega u^4)v + (g_{10}d + g_{11}du + g_{12}du^2 + g_{13}du^3 + g_{14}du^4)v^2$ . The cost of  $p^5$ -Frobenius is  $10M_1$ . This is the same as the cost of  $p^{10}$  Frobenius.