

Secret sharing and duality

Laszlo Csirmaz*

Dedicated to the memory of Frantisek Matúš

Abstract

Secret sharing is an important building block in cryptography. All explicitly defined secret sharing schemes with known exact complexity bounds are multi-linear, thus are closely related to linear codes. The dual of such a linear scheme, in the sense of duality of linear codes, gives another scheme for the dual access structure. These schemes have the same complexity, namely the largest share size relative to the secret size is the same. It is a long-standing open problem whether this fact is true in general: the complexity of any access structure is the same as the complexity of its dual. We give an almost answer to this question. An almost perfect scheme allows negligible errors, both in the recovery and in the independence. There exists an almost perfect ideal scheme on 174 participants whose complexity is strictly smaller than that of its dual.

Key words: secret sharing; ideal access structure; matroid; duality; matroid ports, almost entropic polymatroid.

AMS subject classification: 05B35, 94A15, 06D50, 94A62.

1 Introduction

The construction of a secret sharing scheme whose complexity is smaller than that of its dual is a *tour de force* connecting several different pieces of earlier results. The final construction is actually weaker than would be ideal; filling the gap is an interesting research work. Theorems 19 and 20 state the equivalence of secret sharing conjectures and matroid representation problems.

We assume familiarity with secret sharing schemes, for an overview consult [1]. A significant portion of matroid and polymatroid theory is used. The standard textbook for matroids is [17], for polymatroids see [9] and works of F. Matúš [11, 12]. Nevertheless, most of the theorems and claims are proved – a notable exception is F. Matúš result from [13].

Following the usual practice, sets and their subsets are denoted by capital letters, their elements by lower case letters. The union sign \cup is frequently omitted as well as the curly brackets around singletons. Thus asP denotes the

*Central European University, Budapest

set $\{a, s\} \cup P$. The set difference operator has lower priority than the union, thus $aA - bB$ is $(\{a\} \cup A) - (\{b\} \cup B)$.

The paper is organized as follows. Section 2 introduces polymatroids, secret sharing, complexity measures, duality, and concludes with conjectures on the complexity of dual structures. Section 3 presents two questions on matroid representability and proves that they are equivalent to the conjectures. Section 4 gives a detailed account of Tarik Kaced's result on almost entropic matroids [7], completing the tour. Two proofs are postponed to the Appendix: the first is on matroid circuits used in Claim 5, the second is the the MMRV entropy inequality used in the proof of Theorem 21.

2 Preliminaries

2.1 Polymatroids

A *polymatroid* $\mathcal{M} = (f, M)$ is a non-negative, monotone and submodular function f defined on the collection of non-empty subsets of the finite set M . Here M is the *ground set*, and f is the *rank function*. If f takes non-negative integer values only, then \mathcal{M} is *integer*; an integer polymatroid is a *matroid* if the rank of singletons are either zero or one. Polymatroids can be identified to vectors in the $(2^{|M|} - 1)$ -dimensional Euclidean space where the coordinates are indexed by subsets of M . The collection of polymatroids with ground set M is a full-dimensional pointed polyhedral cone denoted by Γ_M .

For a discrete random variable ξ its information content is measured by the Shannon entropy $\mathbf{H}(\xi)$, see [19]. Let $\xi = \langle \xi_i : i \in M \rangle$ be a collection of discrete random variables with some joint distribution. For a subset $A \subseteq M$, the subcollection $\langle \xi_i : i \in A \rangle$ is denoted by ξ_A . The *conditional entropy* of random variables ξ_A and ξ_B is $\mathbf{H}(\xi_A | \xi_B) = \mathbf{H}(\xi_{A \cup B}) - \mathbf{H}(\xi_B)$ with value between zero and $\mathbf{H}(\xi_A)$. The value is zero if and only if ξ_A is determined completely by ξ_B , and equals $\mathbf{H}(\xi_A)$ if and only if the random variables ξ_A and ξ_B are independent.

As observed by Fujishige [4], the function $A \mapsto \mathbf{H}(\xi_A)$ is a rank function of a polymatroid which we denote by \mathcal{M}_ξ . The polymatroid \mathcal{M} is *entropic* if it can be got this way. The collection of entropic polymatroids on the ground set M is $\Gamma_M^* \subseteq \Gamma_M$. For $|M| \geq 3$ the set Γ_M^* is not closed (in the usual Euclidean topology). Polymatroids in the closure of Γ_M^* are called *almost entropic*, or just *aent*. Aent polymatroids form a full-dimensional convex cone, and every internal point of this cone is entropic [13]. For $|M| \geq 4$ there is a polymatroid in Γ_M with a positive distance from the aent cone [19]; and the aent cone is not polyhedral [14].

By an abuse of notation, we say that \mathcal{M} is an *entropic matroid* if \mathcal{M} is a matroid and for some positive real number λ the polymatroid $\lambda\mathcal{M}$ is entropic.

The singleton $e \in M$ in the polymatroid (f, M) is a *loop* if it has rank zero. In terms of entropic polymatroid being a loop means that the variable ξ_e is deterministic: takes a single value with probability 1. If not mentioned otherwise, polymatroids in this paper have no loops.

With an eye on entropic polymatroids, disjoint subsets A, B of the ground set M are called *independent* if $f(AB) = f(A) + f(B)$. If A and B are independent, $A' \subseteq A, B' \subseteq B$, then A' and B' are independent as well – this follows from the submodularity of the rank function. The single subset A is *independent* if any two disjoint subsets of A are independent. In other words, A is independent iff

$$f(A) = \sum \{f(i) : i \in A\}.$$

A *base* is a maximal independent subset which contains no loops; a *circuit* is a minimal dependent subset. In a loopless polymatroid every independent set can be extended to a base, and every dependent set contains a circuit. In the case when \mathcal{M} is a matroid every base has the same number of elements, and this number equals the rank of the ground set M . Moreover every subset $A \subseteq M$ contains an independent set of size $f(A)$, and every subsets $A \subseteq M$ with rank $f(A) < |A|$ contains a circuit, see [17].

The polymatroid (f, M) is *connected* if for every partition of M into two non-empty sets A and B we have $f(A) + f(B) > f(M)$, that is, A and B are *not* independent. Connected polymatroids have no loops. Indeed, if $i \in M$ is a loop then $f(M) = f(M-i)$, thus the partition $M = \{i\} \cup (M-i)$ contradicts the connectedness.

For an element $i \in M$, the *private info* of i is $f(M) - f(M-i)$, as this is the amount of information which only i and nobody else in M has. If i has no private information, then we say that the polymatroid is *tight at i* . *Tightening at i* means that i is stripped off its private info resulting in the function $f \downarrow i$ defined as

$$f \downarrow i : A \mapsto \begin{cases} f(A) & \text{if } i \notin A \\ f(A) - (f(M) - f(M-i)) & \text{if } i \in A. \end{cases}$$

Of course, $(f \downarrow i, M)$ is a polymatroid tight at i . If $\mathcal{M} = (f, M)$ is tight at every $i \in M$ then \mathcal{M} is *tight*. $\mathcal{M} \downarrow$ is the polymatroid got from \mathcal{M} after tightening at every element of its ground set (the result is independent of the order the elements are taken). Clearly, \mathcal{M} is tight if and only if $\mathcal{M} = \mathcal{M} \downarrow$. If \mathcal{M} is almost entropic then $\mathcal{M} \downarrow$ is almost entropic; this is a result of F. Matúš [16, Lemma 3]. In particular, the tight part of an entropic polymatroid is guaranteed to be almost entropic, but it is not necessarily entropic. A notable exception is the case of matroids: a matroid \mathcal{M} is entropic if and only if $\mathcal{M} \downarrow$ is entropic. It is so as if i is not tight, then $1 = f(M) - f(M-i) \leq f(i) \leq 1$ thus i is independent from all subsets of $M-i$, thus the random variable representing i can be discarded.

2.2 Secret sharing

In a perfect secret sharing scheme there is a *secret*, and each participant from the finite set P receives a *share* such that certain subsets of participants can recover the secret from their joint shares, while other subsets – based on the value of their shares – should have no information on the secret. Subsets who can recover

the secret are *qualified*, the qualified subsets form the *access structure* $\mathcal{A} \subseteq 2^P$. Sets not in \mathcal{A} are called *forbidden* or *unqualified*. An access structure is clearly upward closed. To avoid exceptional cases, \mathcal{A} is assumed to be non-empty (thus all participants together can recover the secret), and the empty set not to be in \mathcal{A} (there must be a secret at all).

The participant $i \in P$ is *important* if there is an unqualified subset such that when i joins this subset, it becomes qualified. If i is not important, then it can join or leave any subset without affecting its status. Consequently the share of an unimportant participant does not play any role, unimportant participants can be discarded. The access structure \mathcal{A} is *connected* if every participant is important. This terminology comes from the relationship between access structures and polymatroids realizing them, see Claims 4 and 5 below. In the rest of the paper, if not mentioned otherwise, access structures are assumed to be connected.

There are several definitions of what secret sharing schemes are. The following definition is considered to be the most general one encompassing all other natural notions [1]. P is the set of participants and $s \notin P$ denotes the secret. A *distribution scheme* is a collection of discrete random variables $\xi = \langle \xi_i : i \in sP \rangle$ with some joint distribution. The value of ξ_s is the *secret*, while the value of ξ_i is the *share* of participant $i \in P$. The secret must be non-trivial, namely it must take at least two different values with positive probability.

The distribution scheme ξ *realizes* an access structure if a) the collection of shares of a qualified subset determine the secret, and b) the collection of shares of an unqualified subset is independent of the secret. Let $\mathcal{M}_\xi = (f, sP)$ be the entropic polymatroid associated with ξ . Shares of the subset $A \subseteq P$ determine the secret iff $\mathbf{H}(\xi_s | \xi_A) = 0$, which translates to $f(sA) = f(A)$. The same collection is independent of the secret if $\mathbf{H}(\xi_s | \xi_A) = \mathbf{H}(\xi_s)$, which translates to $f(sA) = f(A) + f(s)$. This justifies the following definition.

Definition (realizing an access structure). The polymatroid $\mathcal{M} = (f, sP)$ *realizes the access structure* $\mathcal{A} \subseteq 2^P$ if a) $A \in \mathcal{A}$ if and only if $f(sA) = f(A)$, and b) $A \notin \mathcal{A}$ if and only if $f(sA) = f(A) + f(s)$. Polymatroids realizing an access structure are called *secret sharing polymatroids*.

The entropic polymatroid \mathcal{M}_ξ realizes the access structure \mathcal{A} if and only if ξ is a distribution scheme realizing \mathcal{A} . Indeed, if ξ is a distribution scheme then $f(s) = \mathbf{H}(\xi_s)$ is positive, thus one cannot have $f(sA) = f(A)$ and $f(sA) = f(A) + f(s)$ at the same time. Conversely, if \mathcal{M}_ξ realizes \mathcal{A} , then $f(s) > 0$ (otherwise both $f(As) = f(A)$ and $f(As) = f(A) + f(s)$ hold simultaneously), thus the secret is not trivial. Other conditions follow easily.

The proof of the following well-known fact illustrates the ease of reasoning when using polymatroids rather than using entropies directly.

Claim 1. *Suppose \mathcal{M} realizes \mathcal{A} . Then $f(i) \geq f(s)$ for every important participant $i \in P$.*

Proof. As $i \in P$ is important, there is an unqualified subset $A \subseteq P$ (A can be empty) such that iA is qualified. Then $f(sA) = f(A) + f(s)$, and $f(siA) =$

$f(iA)$. Using that $f(i) + f(sA) \geq f(siA)$ and $f(iA) \geq f(A)$ (submodularity and monotonicity) one gets

$$f(i) \geq f(siA) - f(sA) = f(iA) - (f(A) - f(s)) \geq f(s),$$

which proves the claim. \square

All participants together can always determine the secret, thus $f(sP) = f(P)$. This means that the secret has no private info. The private info of the participants does not help at all.

Claim 2. *The polymatroid \mathcal{M} realizes \mathcal{A} if and only if $\mathcal{M}\downarrow$ realizes \mathcal{A} .*

Proof. As observed above, the secret is tight, so let $i \in P$ and $\mathcal{M}\downarrow i = (f^*, sP)$ be the polymatroid after taking away the private info of i . For every $A \subseteq P$, either i is in both A and sA , or i is in none of them, thus

$$f^*(sA) - f^*(A) = f(sA) - f(A).$$

This means that if one of \mathcal{M} or $\mathcal{M}\downarrow i$ realizes \mathcal{A} , then the other does the same. the claim follows after tightening at each participant. \square

Given an access structure it would be tempting to consider tight polymatroids only among those which realize it. But, as was mentioned at the end of Section 2.1, there is no guarantee that the tight part of an entropic polymatroid is also entropic.

Corollary 3. *Suppose \mathcal{A} is connected, and \mathcal{M} realizes \mathcal{A} . If $f(i) = f(s)$ then $i \in P$ is tight.*

Proof. By Claim 2 $\mathcal{M}\downarrow i = (f^*, sP)$ also realizes \mathcal{A} . As $i \in P$ is important, Claim 1 gives $f^*(i) \geq f(s)$. Now $f^*(i) \leq f(i) = f(s)$, thus $f^*(i) = f(i)$ showing that i is tight. \square

According to Claim 4 below, a polymatroid realizing a connected access structure must be connected. The converse is not true in general. In the special case when the polymatroid is a matroid the converse follows from some standard properties of matroid circuits [17].

Claim 4. *Suppose the polymatroid \mathcal{M} realizes the access structure \mathcal{A} . If the access structure is connected, then \mathcal{M} is connected.*

Proof. Assume, by contradiction, that $\mathcal{M} = (h, sP)$ is not connected, which means $sA \cup B$, $B \neq \emptyset$ is a partition of the ground set sP and $h(sAB) = h(sA) + h(B)$. In other words, sA and B are independent, consequently subsets of sA and B are independent as well. Let $b \in B$ and assume $A'B'$ is not qualified while $A'bB'$ is qualified with $A' \subseteq A$ and $B' \subseteq B - b$. Then $h(sA'B') = h(A'B') + h(s)$, from where

$$h(sA') + h(B') = h(sA'B') = h(A'B') + h(s) = h(A') + h(B') + h(s).$$

On the other hand, $h(sA'bB') = h(A'bB')$, which gives

$$h(sA') + h(bB') = h(sA'bB') = h(A'bB') = h(A') + h(bB').$$

From the first line $h(sA') = h(A') + h(s)$, while from the second $h(sA') = h(A')$, a contradiction. \square

Claim 5. *Suppose \mathcal{M} is a matroid which realizes the access structure \mathcal{A} . If \mathcal{M} is connected then so is the access structure \mathcal{A} .*

Proof. Using matroid terminology, A is *dependent* if $h(A) < |A|$, and A is a *circuit* if it is a minimal dependent set. If C is a circuit, then $h(C) = |C| - 1$ and for each $i \in C$, $h(C-i) = |C| - 1$. A circuit *connects* two points if it contains both of them.

Let the ground set of the matroid be sP and pick some $a \in P$. To show that a is important it is enough to find a circuit C connecting a and s . Indeed, let $A = C-s$, then $a \in A$ and $h(sA) = h(C) = h(C-s) = h(A)$, thus A is qualified. C is minimal dependent, thus $sA-a = C-a$ is independent, and then $h(sA-a) = h(s) + h(A-a)$ which means $A-a$ is not qualified.

To finish the proof it suffices to quote the following result from matroid theory [17, Proposition 4.1.4]: *a matroid is connected if and only if any two points can be connected by a circuit.* For a quick proof see the Appendix. \square

2.3 Complexity

Distribution schemes realizing an access structure scale up: taking n independent copies of the scheme all entropies are multiplied by n and the composite scheme still realizes the same access structure. Similarly, whether a polymatroid realizes an access structure or not is invariant for multiplying the polymatroid by any positive constant. When defining the efficiency one has to take into account this scalability. The usual way is to measure everything in multiples of the secret size. For example, if $\mathcal{M} = (f, sP)$ is a secret sharing polymatroid, then the *relative share size* of participant $i \in P$ is $f(i)/f(s)$, and the (worst case) *complexity* of \mathcal{M} is

$$\sigma(\mathcal{M}) = \max \left\{ \frac{f(i)}{f(s)} : i \in P \right\},$$

Other complexity measures, not considered here, include average relative size, and the scaled total randomness. If \mathcal{M} realizes the connected access structure \mathcal{A} , then $\sigma(\mathcal{M}) \geq 1$ by Claim 1. Access structures where this lower bound is attained are called *ideal*.

Definition (ideal and almost ideal structures). The access structure \mathcal{A} is *ideal* if it can be realized by an entropic polymatroid with complexity 1. The access structure \mathcal{A} is *almost ideal* if it can be realized by an almost entropic polymatroid with complexity 1.

In general, the usual definition of the complexity of an access structure is the infimum of the complexity of all secret sharing schemes realizing it:

$$\sigma(\mathcal{A}) = \inf \{ \sigma(\mathcal{M}) : \mathcal{M} \text{ is entropic and realizes } \mathcal{A} \}.$$

Interestingly, there is a non-ideal (according to our definition) access structure with complexity 1, see [2, Section 6], thus the infimum here is not necessarily taken. The cone of almost entropic polymatroids is closed, thus an access structure with complexity 1 is almost ideal. It is an interesting open question whether the converse is true. When approximating an aent polymatroid by an entropic one, the only guarantee is that the rank functions differ by a small (negligible) amount. This means that qualified subsets can recover the secret with “overwhelming probability” only (as $\mathbf{H}(s|A)$ is not necessarily zero, only negligible), and unqualified subsets might get information on the secret (as $\mathbf{H}(s|A)$ can be strictly smaller than $\mathbf{H}(s)$), but this information is negligible. The question is can we patch these imperfections by adding a small amount of entropy to the secret? It seems that for secret recovery the answer is *yes*; for independence the author tends to believe that the answer is *no*.

Next to $\sigma(\mathcal{A})$ other complexity measures can be defined by considering other polymatroid classes. Realizing \mathcal{A} by an entropic polymatroid is the same as realizing it by a distribution scheme. Realizing by an almost entropic polymatroids instead means that one relaxes the strict requirements of recoverability and independence “up to a negligible amount”. Linearly representable polymatroids are important from both practical and theoretical point of view. Such polymatroids arise from linear error correcting codes [6], they are studied extensively and typically provide concise, efficient and low complexity schemes. We consider the following polymatroid classes, listed in decreasing order:

- a) all polymatroids,
- b) almost entropic polymatroids,
- c) entropic polymatroids,
- d) (conic hull of) linearly representable polymatroids.

Every access structure can be realized by a linearly representable polymatroid, thus every class gives a complexity notion on access structures. For classes a), c) and d) they are denoted by κ , σ , and λ [18]. For class b) we use $\bar{\sigma}$ to indicate that we are considering the closure of entropic polymatroids. The earlier definition of $\sigma(\mathcal{A})$ is the same as given here.

$$\begin{aligned} \kappa(\mathcal{A}) &= \inf \{ \sigma(\mathcal{M}) : \mathcal{M} \text{ realizes } \mathcal{A} \}, \\ \bar{\sigma}(\mathcal{A}) &= \inf \{ \sigma(\mathcal{M}) : \mathcal{M} \text{ is aent and realizes } \mathcal{A} \}, \\ \sigma(\mathcal{A}) &= \inf \{ \sigma(\mathcal{M}) : \mathcal{M} \text{ is entropic and realizes } \mathcal{A} \}, \\ \lambda(\mathcal{A}) &= \inf \{ \sigma(\mathcal{M}) : \mathcal{M} \text{ is linear and realizes } \mathcal{A} \}. \end{aligned}$$

For the same access structure these values increase (as less and less polymatroids are considered). Each pair of these measures is known to be separated except for σ and $\bar{\sigma}$, see [1, 18].

2.4 Duals

Let P be the set of participants, and $\mathcal{A} \subseteq 2^P$ be an access structure. The qualified subsets in the *dual access structure* \mathcal{A}^\perp are the complements of unqualified subsets of \mathcal{A} :

$$\mathcal{A}^\perp = \{A \subseteq P : P-A \notin \mathcal{A}\}.$$

Clearly, the dual of \mathcal{A}^\perp is \mathcal{A} ; $\emptyset \notin \mathcal{A}^\perp$ and $P \in \mathcal{A}^\perp$ as these are true for \mathcal{A} .

Claim 6. \mathcal{A} is connected if and only if \mathcal{A}^\perp is connected.

Proof. Suppose \mathcal{A} is connected, we show that \mathcal{A}^\perp is connected. The other direction follows from $(\mathcal{A}^\perp)^\perp = \mathcal{A}$. Let $a \in P$, and $A \subset P$ unqualified in \mathcal{A} such that $aA \in \mathcal{A}$. Such an A exists as \mathcal{A} is connected. Then $a \in P-A$ is qualified in \mathcal{A}^\perp and $(P-A)-a \notin \mathcal{A}^\perp$ is unqualified in \mathcal{A}^\perp , as required. \square

Let $\mathcal{M} = (f, M)$ be a polymatroid. Define the discrete measure μ on subsets of M by $\mu(i) = f(i)$. As the measure is additive, for every subset $A \subseteq M$ we have

$$\mu(A) = \sum \{f(i) : i \in A\}.$$

The *dual of the polymatroid* \mathcal{M} is $\mathcal{M}^\perp = (f^\perp, M)$ where the function f^\perp is defined for subsets of M as

$$f^\perp : A \mapsto f(M-A) + \mu(A) - f(M).$$

By submodularity, f^\perp is non-negative; submodularity holds by an easy inspection, thus \mathcal{M}^\perp is a polymatroid. If \mathcal{M} is integer-valued then so is \mathcal{M}^\perp ; moreover if \mathcal{M} is a matroid (the rank of a singleton is zero or one), then so is the dual.

Claim 7. a) \mathcal{M} is connected if and only if \mathcal{M}^\perp is connected. b) \mathcal{M} realizes the access structure \mathcal{A} if and only if \mathcal{M}^\perp realizes \mathcal{A}^\perp .

Proof. a) Suppose $A \cup B$ is a partition of M , then $\mu(A) + \mu(B) = \mu(M)$. By the definition of f^\perp we have

$$f^\perp(A) + f^\perp(B) - f^\perp(M) = f(B) + f(A) - f(M).$$

If one of them is positive, then the other is positive, as required.

b) Let $M = sP$, $A \subseteq P$, then $\mu(sA) - \mu(A) = \mu(s) = f(s)$, thus

$$(f^\perp(sA) - f^\perp(A)) + (f(sP-A) - f(P-A)) = f(s).$$

If \mathcal{M} realizes \mathcal{A} , then $f(sP-A) - f(P-A)$ is either zero or $f(s)$ depending on whether $P-A \in \mathcal{A}$ or not. Consequently $f^\perp(sA) - f^\perp(A)$ is either zero or $f(s)$ depending on whether $P-A \notin \mathcal{A}$ or not. Thus f^\perp realizes \mathcal{A}^\perp . The converse is similar. \square

The dual polymatroid \mathcal{M}^\perp is always tight as

$$f^\perp(M-i) = f(i) + \mu(M-i) - f(M) = \mu(M) - f(M) = f^\perp(M).$$

Consequently the dual of \mathcal{M}^\perp is also tight, and if \mathcal{M} was not tight, the dual of \mathcal{M}^\perp cannot be the same as \mathcal{M} . However, if \mathcal{M} is tight, then it equals $\mathcal{M}^{\perp\perp}$, in particular $\mathcal{M}^{\perp\perp\perp} = \mathcal{M}^\perp$ always true.

Claim 8. a) Suppose \mathcal{M} is tight. Then $\mathcal{M}^{\perp\perp} = \mathcal{M}$, moreover \mathcal{M} and \mathcal{M}^\perp has the same value on singletons. b) For every polymatroid \mathcal{M} , $\mathcal{M}^{\perp\perp} = \mathcal{M}\downarrow$.

Proof. a) We start with the second claim. By the assumption, $f(M) = f(M-i)$,

$$f^\perp(i) = f(M-i) + \mu(i) - f(M) = \mu(i) = f(i),$$

as claimed. It means that $\mu^\perp(A) = \mu(A)$, and then

$$\begin{aligned} f^\perp(M-A) &= f(A) + \mu(M-A) - f(M), \\ f^\perp(M) &= \mu(M) - f(M), \end{aligned}$$

thus

$$\begin{aligned} f^{\perp\perp}(A) &= f^\perp(M-A) + \mu^\perp(A) - f^\perp(M) = \\ &= f(A) + \mu(M-A) + \mu(A) - \mu(M) = f(A), \end{aligned}$$

proving $\mathcal{M}^{\perp\perp} = \mathcal{M}$.

b) It is enough to show that the dual of \mathcal{M} and the dual of $\mathcal{M}\downarrow$ are the same, from here the claim follows by a). In $\mathcal{M}\downarrow$ the rank of every set containing $i \in M$ decreases by the same amount. In the expression

$$f(M-A) + \mu(A) - f(M)$$

this amount is added once in the first two terms, and subtracted once in the last term, thus it cancels. \square

2.5 Factor and principal extension

Let $\mathcal{M} = (h, M)$ be a polymatroid. Partitions of the ground set M can be considered as equivalence classes of an equivalence relation on M . Let \cong be an equivalence relation on M , $N = M/\cong$ be the set of equivalence classes, and $\varphi : M \rightarrow N$ be the map which assigns to each element its equivalence class. The *factor of \mathcal{M} by \cong* , denoted as \mathcal{M}/\cong , is the pair (g, N) where g assigns the value $g : A \mapsto h(\varphi^{-1}(A))$ to subsets of N (that is, union of complete equivalence classes). It is clear that \mathcal{M}/\cong is a polymatroid.

Let $a \in M$, and $\alpha \geq 0$ be a real number. The *principal extension $\mathcal{M}_{a,\alpha}$* is a one-point extension of \mathcal{M} defined on the set $M \cup \{a'\}$ assigning the value

$$h : a' A \mapsto \min \{h(A) + \alpha, h(aA)\}$$

to new subsets. It is a routine to check that the principal extension is a polymatroid [9]. Principal extension of an almost entropic polymatroid is almost entropic. This is an immediate consequence of (and actually, is equivalent to) a result of F. Matúš [13, Theorem 2], see also [16, Lemma 3]. We state this result without proof.

Theorem 9 (F. Matúš). *If the polymatroid \mathcal{M} is almost entropic, then so is the principal extension $\mathcal{M}_{a,\alpha}$.* \square

Matúš' proof guarantees the extension to be only almost entropic even if \mathcal{M} is entropic. In fact, there is an entropic polymatroid where some principal extension is not entropic.

Principal extensions can be used to “split atoms” of a polymatroid, which, in turn, will be used to prove that integer polymatroids are factors of matroids. Let us see the details. In what follows $\mathcal{M} = (h, M)$ is a polymatroid.

Lemma 10. *Let $a \in M$, and α_1, α_2 be non-negative numbers whose sum is $h(a)$. There is a polymatroid $\mathcal{M}' = (h', a_1 a_2 \cup M - a)$ such that $h'(a_i) = \alpha_i$, and \mathcal{M} is a factor of \mathcal{M}' collapsing a_1 and a_2 to a . Moreover, \mathcal{M} is almost entropic if and only if so is \mathcal{M}' .*

Proof. Let \mathcal{M}' be the principal extension $\mathcal{M}_{a, \alpha_1}$ adding the new point a_1 , so that $M' = a_1 \cup M$; then let \mathcal{M}'' be the principal extension $\mathcal{M}'_{a_1, \alpha_2}$ adding the new point a_2 . Then for each $A \subseteq M - a$ we have

$$\begin{aligned} h''(A) &= h(A), \\ h''(a_1 A) &= \min \{h(A) + \alpha_1, h(aA)\}, \\ h''(a_2 A) &= \min \{h(A) + \alpha_2, h(aA)\}, \\ h''(a_1 a_2 A) &= \min \{h(A) + \alpha_1 + \alpha_2, h(aA)\}. \end{aligned}$$

As $h(A) + \alpha_1 + \alpha_2 = h(A) + h(a) \geq h(aA)$, we have $h''(a_1 a_2 A) = h(aA)$. This shows that \mathcal{M}'' restricted to the ground set $a_1 a_2 \cup M - a$ is the required splitting. If \mathcal{M} is aent, then both \mathcal{M}' and \mathcal{M}'' are aent by Theorem 9. A restriction and a factor of an aent polymatroid is trivially aent, proving the last claim. \square

Lemma 11. *Let $\mathcal{M} = (f, aN)$ be tight, and suppose $\mathcal{N} = (g, a_1 a_2 N)$ splits a in \mathcal{M} as $g(a_i) = \alpha_i$. Then \mathcal{N}^\perp splits a in \mathcal{M}^\perp at the same way.*

Proof. Let $A \subseteq N$, then $g(A) = f(A)$ and $g(a_1 a_2 A) = f(aA)$. Calculating $g^\perp(A)$ one gets

$$\begin{aligned} g^\perp(A) &= g(a_1 a_2 N - A) + \mu(A) - g(a_1 a_2 N) = \\ &= f(aN - A) + \mu(A) - f(aN) = f^\perp(A), \end{aligned}$$

and similarly $g^\perp(a_1 a_2 A) = f^\perp(aA)$. Finally,

$$\begin{aligned} g^\perp(a_1 A) &= g(a_1 a_2 N - a_1 A) + \mu(a_1 A) - g(a_1 a_2 N) \\ &= g(a_2 N - A) + \mu(a_1) + \mu(A) - f(aN) \\ &= \min \{f(N - A) + \alpha_2, f(aN - A)\} + \alpha_1 + \mu(A) - f(aN) \\ &= \min \{f(N - A) + \mu(aA) - f(aN), f(aN - A) + \mu(A) - f(aN) + \alpha_1\} \\ &= \min \{f^\perp(aA), f^\perp(A) + \alpha_1\}, \end{aligned}$$

thus \mathcal{N}^\perp splits a as claimed as $f^\perp(a) = f(a) = \alpha_1 + \alpha_2$ using that \mathcal{M} is tight. \square

Factors of a matroid are integer polymatroids. Helgason's theorem [5] says that the converse is true: every integer polymatroid is a factor of some matroid. We need the following strengthening of this result.

Theorem 12. *For each integer polymatroid \mathcal{M} there is a matroid $\varphi(\mathcal{M})$ such that a) \mathcal{M} is a factor of $\varphi(\mathcal{M})$, b) \mathcal{M} is aent if and only if $\varphi(\mathcal{M})$ is aent, c) if \mathcal{M} is tight, then $\varphi(\mathcal{M}^\perp)$ is the dual of $\varphi(\mathcal{M})$.*

Proof. Let \mathcal{M} be an integer polymatroid. The matroid $\varphi(\mathcal{M})$ is generated by a series of splitting. If all singletons have rank zero or one, then \mathcal{M} is a matroid, and we are done. Otherwise some $a \in M$ has rank $h(a) > 1$. Using Lemma 10 split a into two with ranks 1 and $h(a) - 1$. All ranks in the split polymatroid \mathcal{M}' remain integer, and by Lemma 10 \mathcal{M}' is aent if and only if \mathcal{M} is aent. Continue this way to get the matroid $\varphi(\mathcal{M})$. Clearly \mathcal{M} is a factor of $\varphi(\mathcal{M})$, and c) holds by Lemma 11. \square

2.6 The duality conjecture

Fix the connected access structure $\mathcal{A} \subset 2^P$ and consider all polymatroids on the ground set sP which realize \mathcal{A} . We are interested in $\kappa(\mathcal{A})$, the minimal complexity of these polymatroids. By Claim 2 the search can be restricted to tight polymatroids. Suppose the infimum is attained by the tight polymatroid \mathcal{M} . (It is attained as polymatroids form a closed set.) Claim 8 a) implies that \mathcal{M} and \mathcal{M}^\perp have the same complexity. According to Claim 7 b) \mathcal{M}^\perp realizes \mathcal{A}^\perp , thus $\kappa(\mathcal{A}^\perp) \leq \kappa(\mathcal{A})$. Applying the same reasoning to the dual structure we get $\kappa(\mathcal{A}^{\perp\perp}) \leq \kappa(\mathcal{A}^\perp)$. As $\mathcal{A}^{\perp\perp}$ and \mathcal{A} are the same, we have

Claim 13. *For every access structure we have $\kappa(\mathcal{A}) = \kappa(\mathcal{A}^\perp)$.* \square

Every access structure can be realized by some linearly representable polymatroid, the complexity measure $\lambda(\mathcal{A})$ defines the infimum of the complexity of such representations. It is well-known that the conic hull of linearly representable polymatroids is a closed subset of the entropic polymatroids, and it is closed for taking duals. Therefore it is also closed for tightening by Claim 8 b). The corresponding complexity measure is $\lambda(\mathcal{A})$, and the same reasoning as above gives

Claim 14. *For every access structure we have $\lambda(\mathcal{A}) = \lambda(\mathcal{A}^\perp)$.* \square

Every explicitly defined access structure \mathcal{A} with known exact complexity value $\sigma(\mathcal{A})$ satisfies $\lambda(\mathcal{A}) = \sigma(\mathcal{A}) = \kappa(\mathcal{A})$ – consequently the same is true for the dual structure, and then $\sigma(\mathcal{A}) = \sigma(\mathcal{A}^\perp)$. It is a long-standing open problem whether the statement similar to Claims 13 and 14 holds for the entropic complexity σ .

Conjecture 1 (complexity of dual structure). *For every access structure we have $\sigma(\mathcal{A}) = \sigma(\mathcal{A}^\perp)$.*

The conjecture is probably not true, but even the particular case when \mathcal{A} is an ideal access structure resisted all efforts. Recall, that \mathcal{A} is ideal if it can be realized by an ideal entropic polymatroid, or, equivalently, by an ideal distribution scheme.

Conjecture 2 (dual of ideal structure). *The dual of an ideal access structure is ideal.*

Refuting the second conjecture does not necessarily refutes Conjecture 1 as the dual might be non-ideal while having complexity 1. In Section 3 we prove that Conjecture 2 is equivalent to a question about matroid representability. Using results of that section, and a construction by Tarik Kaced [7] the duality question for almost ideal schemes is settled.

3 Ideal structures and matroids

First we give a self-contained proof of a somewhat extended result of Blakley and Kabatianski [3, 15] connecting ideal access structures and matroids. Using this connection we present a statement about matroid representability which is equivalent to Conjecture 2.

Fix the connected access structure $A \subset 2^P$ and suppose the polymatroid $\mathcal{M} = (f, sP)$ realizes it. Assume furthermore that \mathcal{M} has complexity 1, that is, the rank of all singletons equals $f(s)$. The following lemmas establish some structural properties of \mathcal{M} . In the lemmas A is a subset of P , $a \in P$, and s denotes the secret.

Lemma 15. *Suppose $A \in \mathcal{A}$ and $A-a \notin \mathcal{A}$. Then $f(A) - f(A-a) = f(s)$.*

Proof. By submodularity of the rank function f , we have

$$\begin{aligned} f(a) &\geq f(A) - f(A-a) = f(sA) - (f(sA-a) - f(s)) = \\ &= (f(sA) - (f(sA-a))) + f(s) \geq f(s). \end{aligned}$$

As $f(a) = f(s)$, the conclusion follows. \square

Lemma 16. *Let $a \in A' \subseteq A$, Suppose $A-a$ and A' are qualified and $A'-a$ is not. Then $f(A) = f(A-a)$.*

Proof. For qualified subsets $f(sA') = f(A')$, etc., for the unqualified subset $f(sA'-a) = f(A'-a) + f(s)$. Thus

$$\begin{aligned} f(A) - f(A-a) &= f(sA) - f(sA-a) \leq f(sA') - f(sA'-a) = \\ &= (f(A') - f(A'-a)) - f(s) = 0, \end{aligned}$$

where the inequality follows from submodularity and the last equality from Lemma 15. Thus $0 \leq f(A) - f(A-a) \leq 0$, proving the claim. \square

Lemma 17. *Suppose $f(A) - f(A-a) = f(s)$, and $a \in A' \subseteq A$. Then $f(A') - f(A'-a) = f(s)$.*

Proof. By submodularity, $f(A) - f(A-a) \leq f(A') - f(A'-a) \leq f(a)$. As both sides equal $f(s)$, the claim follows. \square

Theorem 18 (Blakley–Kabatianski). *Let $\mathcal{A} \subset 2^P$ be a connected access structure and $\mathcal{M} = (f, sP)$ be a polymatroid realizing \mathcal{A} such that $f(a) = f(s) = 1$ for all $a \in P$. Then \mathcal{M} is a matroid which is uniquely determined by the access structure.*

Proof. All singletons have rank 1, thus \mathcal{M} is a matroid if all ranks are integer. The basic idea is to show that for any subset A of the ground set sP one can find an element a of A such that $f(A) - f(A-a)$ is either zero or one. The additional claim that \mathcal{M} is uniquely determined by \mathcal{A} follows from the fact that for the chosen element $a \in A$ the value of $f(A) - f(A-a)$ depends only on the access structure, and not on the particular realization.

If the subset contains the secret s , then $f(sA) - f(A)$ is either zero or $f(s) = 1$ depending on whether $A \in \mathcal{A}$ or $A \notin \mathcal{A}$, which settles this case. So assume $A \subseteq P$.

When A is qualified, then there are two cases. If $A-a$ is not qualified for some $a \in A$, then Lemma 15 gives that this difference is $f(s) = 1$. If all $A-a$ is qualified, then pick a minimal qualified $A' \subseteq A$ and use Lemma 16 with any $a \in A'$.

Thus assume A is unqualified. As \mathcal{A} is connected, there is an unqualified subset B such that AB is qualified (pick any element of A and let B show that this element is important). Choose such an unqualified B such that the set $B-A$ has minimal cardinality, and within this constrain $A \cap B$ has maximal cardinality. Then $AB-k$ is unqualified for any $k \in B-A$ (as otherwise $B-A$ is not minimal), and aB is qualified for any $a \in A-B$ (as otherwise $A \cap B$ is not maximal). Fix $a \in A-B$. With any $k \in B-A$ we have that AB is qualified, $AB-k$ is not. Lemma 15 gives $f(AB) - f(AB-k) = 1$, and by Lemma 17, $f(A') - f(A'-k) = 1$ for all $k \in A' \subseteq AB$. By induction this gives both $f(AB) - f(A) = |A-B|$ and $f(AB-a) - f(A-a) = |A-B|$. Therefore $f(A) - f(A-a) = f(AB) - f(AB-a)$. Now AB is qualified. If $AB-a$ is unqualified, then by Lemma 15 this difference is $f(s) = 1$. If $AB-a$ is qualified, then $f(AB) = f(AB-a)$ using Lemma 16 with $A' = aB$. \square

The main result of this section is the equivalence of a statement about matroid representability and Conjecture 2. Recall that \mathcal{M} is an *entropic matroid* if for some positive λ the polymatroid $\lambda\mathcal{M}$ is entropic.

Theorem 19. *The following statements are equivalent.*

- a) *The dual of every ideal access structure is ideal.*
- b) *The dual of every entropic matroid is entropic.*

Proof. Let us first make some simplifying assumptions. In a) the access structure can be assumed to be connected: simply forget about the unimportant participants, they will be unimportant in the dual structure. In b) the matroid can be assumed to be tight and connected. This is so as the matroids \mathcal{M} and $\mathcal{M}\downarrow$ are entropic at the same time: if i has a non-zero private info, then it is

completely independent of the rest of the matroid. Furthermore, if \mathcal{M} is not connected, then it is an independent sum of the connected components, and then \mathcal{M}^\perp is the sum of the duals of the components.

a) \rightarrow b) As remarked above, we may assume that the entropic matroid \mathcal{M} is tight and connected. Pick any element of its ground set and name it s , the remaining elements are in P . Since \mathcal{M} is connected, it has no loops, thus $f(s) = 1$. Define the access structure $\mathcal{A} \subset 2^P$ by

$$\mathcal{A} = \{A \subseteq P : f(sA) = f(A)\}.$$

Clearly \mathcal{M} realizes this access structure, consequently \mathcal{A} is ideal, and by Claim 5 it is also connected. By Claim 7 b), the dual matroid \mathcal{M}^\perp realizes \mathcal{A}^\perp .

As \mathcal{A} is a connected ideal structure, assumption a) says that \mathcal{A}^\perp is ideal. Let \mathcal{M}' be the scaled entropic polymatroid which realizes \mathcal{A}^\perp with $f'(s) = f'(a) = 1$. As \mathcal{A}^\perp is connected by Claim 6, conditions of Theorem 18 hold. Consequently \mathcal{M}' is the unique matroid realizing \mathcal{A}^\perp . As \mathcal{M}^\perp also realizes the same access structure, \mathcal{M}' and \mathcal{M}^\perp are the same matroids. Now \mathcal{M}' is a scaled version of an entropic polymatroid, thus $\mathcal{M}' = \mathcal{M}^\perp$ is an entropic matroid, as was required.

b) \rightarrow a) Let \mathcal{A} be an ideal connected access structure realized by the entropic polymatroid \mathcal{M}^* . As \mathcal{A} is connected and ideal, we have $f^*(i) = f^*(s) > 0$ for all participants $i \in P$. Let $\lambda = 1/f^*(s)$ and $\mathcal{M} = \lambda\mathcal{M}^*$. Then \mathcal{M} also realizes \mathcal{A} and $f(i) = f(s) = 1$ for all $i \in P$. By Corollary 3 \mathcal{M} is tight, and by Theorem 18 \mathcal{M} is a matroid. As \mathcal{A} is connected, by Claim 4 \mathcal{M} is connected. Consequently \mathcal{M} is a tight, connected, entropic matroid which realizes the access structure \mathcal{A} . By assumption b) \mathcal{M}^\perp is an entropic matroid, realizes \mathcal{A}^\perp by Claim 7 b); finally by Claim 8 a) \mathcal{M}^\perp and \mathcal{M} have the same value on singletons. Thus $\lambda^\perp\mathcal{M}^\perp$ is an entropic polymatroid for some positive λ^\perp , realizes \mathcal{A}^\perp , and has complexity $\sigma(\mathcal{M}^\perp) = \sigma(\mathcal{M}) = 1$. Therefore \mathcal{A}^\perp is ideal. \square

Almost entropic polymatroids form a closed cone, which means that positive multiplies of an entropic polymatroid are entropic. Consequently the definition of almost entropic matroids does not require scaling as was the case for entropic matroids. The matroid \mathcal{M} is *almost entropic* if it is almost entropic as a polymatroid. Repeating the proof above word by word while replacing “ideal” by “almost ideal” and “entropic” by “almost entropic” everywhere one gets the following theorem.

Theorem 20. *The following statements are equivalent.*

- a) *The dual of every almost ideal access structure is almost ideal.*
- b) *The dual of every almost entropic matroid is almost entropic.* \square

4 Duals of almost entropic matroids

We have almost all the pieces together to prove the main result:

Theorem 21. *There is an almost ideal access structure whose dual is not almost ideal.*

By Theorem 20 we need to exhibit an almost entropic matroid whose dual is not almost entropic. The existence of such a matroid was proved by Tarik Kaced [7, Theorem 2], this section is a detailed account of that result. The proof starts with the construction of an entropic polymatroid whose dual is not entropic. Using a continuity argument and linear scaling, one gets an integer polymatroid with the same properties. Theorem 12 established a connection between integer polymatroids and matroids which preserves duality and almost entropicity. To complete the tour apply this theorem to get the required matroid. Now let us see the details.

Finding an entropic polymatroid whose dual is not entropic was a long-standing open problem. The example below is due to Kaced [7]. The polymatroid is specified by a distribution on five binary random variables. To show that its dual is not entropic, Kaced used a 5-variable non-Shannon type information inequality, see [10, 12]. Such an inequality is a hyperplane in the $(2^{|M|} - 1)$ -dimensional space which a) contains all entropic points on its non-negative side (consequently all aent points as well), and b) cuts into the polymatroid cone Γ_M . Entropy inequalities are typically written using abbreviations originating in information theory. For disjoint subsets A, B, C we write

$$\begin{aligned} h(A|B) &= h(AB) - h(B), \\ h(A, B) &= h(A) + h(B) - h(AB), \\ h(A, B | C) &= h(AC) + h(BC) - h(ABC) - h(C), \end{aligned}$$

corresponding to conditional entropy, mutual information, and conditional mutual information, respectively. In any polymatroid these expressions are always non-negative. The MMRV inequality written for the singletons of the five-element set $\{abcde\}$ is

$$\begin{aligned} &(h(a, b|c) + h(b, c|a) + h(c, a|b)) + \\ &+ (h(b, c|d) + h(b, c|e) + h(d, e) - h(b, c)) \geq 0. \end{aligned} \tag{1}$$

For a short proof that this inequality holds for aent polymatroids see the Appendix.

Claim 22. *There is a tight, integer and aent polymatroid whose dual is not aent.*

Proof. The distribution on five random variables ξ_a, \dots, ξ_e is specified in Table 1. Each of the variables takes either zero or one, there are only eight combinations with positive probability. The associated polymatroid \mathcal{M}_ξ is entropic, the left hand side of (1) evaluates to 0.106736. The dual \mathcal{M}_ξ^\perp is not aent as the left hand side of (1) is -0.070148.

The duality operation is continuous, thus duals of the polymatroids in a small neighborhood of \mathcal{M}_ξ are still violating the MMRV inequality. The entropic polymatroid \mathcal{M}_ξ is on the boundary of the aent cone (for example, d and e have no private info), but there is another polymatroid \mathcal{M}' arbitrarily close to \mathcal{M}_ξ

ξ_a	ξ_b	ξ_c	ξ_d	ξ_e	Prob
0	0	0	0	0	0.077
0	0	1	1	0	0.182
0	1	0	0	1	0.182
0	1	1	0	0	0.077
1	0	0	0	0	0.105
1	0	1	0	0	0.136
1	1	0	0	0	0.136
1	1	1	0	0	0.105

Table 1: Distribution on five variables

inside that cone. By [13] interior points of the aent cone are entropic. Take \mathcal{M}'' very close to \mathcal{M}' such that all coordinates rational. Let n be the smallest common denominator of the fractions in the coordinates. Coordinates in $n\mathcal{M}''$ are integer. The dual of $n\mathcal{M}''$ violates the MMRV as the dual of \mathcal{M}'' violates it, and the left hand side of (1) also multiplies by n . Finally, $n\mathcal{M}''$ is entropic: to realize it take n independent copies of the random variables realizing \mathcal{M}'' . The tight part of $n\mathcal{M}''$ is integer and almost entropic, its dual is the same as the dual of $n\mathcal{M}''$ by Claim 8 b), proving the claim.

Using the distribution ξ above, such an integer polymatroid can be constructed directly. For a subset $A \subseteq M$ define the the polymatroid \mathbf{r}_A as

$$\mathbf{r}_A : I \mapsto \begin{cases} 1 & \text{if } A \cap I \neq \emptyset, \\ 0 & \text{otherwise.} \end{cases}$$

Clearly, $\lambda \mathbf{r}_A$ is entropic for every positive λ . As A runs over all non-empty subsets of M these polymatroids are linearly independent and span a full-dimensional subcone of Γ_M^* consisting of entropic polymatroids only [13]. The idea is that take an integer multiple of \mathcal{M}_ξ (which is entropic), and use some linear combination of \mathbf{r}_A 's to round up the coordinates to integer values. This idea works. The polymatroid

$$\begin{aligned} \mathcal{M} = & 51\mathcal{M}_\xi + 0.3725518(\mathbf{r}_{abd} + \mathbf{r}_{acd} + \mathbf{r}_{abe} + \mathbf{r}_{ace}) + \\ & 0.1431172(\mathbf{r}_b + \mathbf{r}_c) + 0.0272887\mathbf{r}_{bc} + 0.4970901\mathbf{r}_{abc} + \\ & 0.6449149(\mathbf{r}_{bd} + \mathbf{r}_{cd} + \mathbf{r}_{be} + \mathbf{r}_{ce}) + 0.0129470(\mathbf{r}_{bcd} + \mathbf{r}_{bce}) + \\ & 0.1185108\mathbf{r}_a + 0.5029099(\mathbf{r}_{ab} + \mathbf{r}_{ac}) + 0.3230136(\mathbf{r}_{ad} + \mathbf{r}_{ae}) + \\ & 0.3408074(\mathbf{r}_{abcd} + \mathbf{r}_{abce}) + 0.4828705\mathbf{r}_{bcde} + 0.5732815\mathbf{r}_{abcde} \end{aligned}$$

is clearly entropic, and it is integer. This is so as the coefficients in this formula are the solutions of a system of linear equations yielding exact values. Table 2 shows the coordinates of $51\mathcal{M}_\xi$ (left column), the integer entropic polymatroid \mathcal{M} (middle column), and the tightening of \mathcal{M} (right column). The value of the MMRV inequality for the dual of \mathcal{M} is -1 , thus \mathcal{M}^\perp is not almost entropic. As $\mathcal{M}^\perp = (\mathcal{M}^\downarrow)^\perp$, the tight part of \mathcal{M} is a tight, integer, aent polymatroid whose dual is not aent. \square

<i>a</i>	49.987448	55	37	<i>abc</i>	146.585502	155	89
<i>b</i>	50.031000	55	31	<i>abd</i>	111.373534	119	77
<i>c</i>	50.031000	55	31	<i>acd</i>	111.373534	119	77
<i>d</i>	34.232147	38	38	<i>abe</i>	111.373534	119	77
<i>e</i>	34.232147	38	38	<i>ace</i>	111.373534	119	77
<i>ab</i>	100.018448	107	65	<i>ade</i>	90.899025	99	81
<i>ac</i>	100.018448	107	65	<i>bde</i>	97.350040	105	81
<i>ad</i>	74.201801	81	63	<i>cde</i>	97.350040	105	81
<i>ae</i>	74.201801	81	63	<i>bcd</i>	113.027026	121	73
<i>bc</i>	97.350040	105	57	<i>bce</i>	113.027026	121	73
<i>bd</i>	73.690520	80	56	<i>abcd</i>	146.585502	155	89
<i>be</i>	73.690520	80	56	<i>abce</i>	146.585502	155	89
<i>cd</i>	73.690520	80	56	<i>abde</i>	122.728619	131	89
<i>ce</i>	73.690520	80	56	<i>acde</i>	122.728619	131	89
<i>de</i>	65.520445	72	72	<i>bcde</i>	128.704012	137	89
				<i>abcde</i>	146.585502	155	89

Table 2: An integer entropic polymatroid

Let the tight integer polymatroid provided by Claim 22 be \mathcal{N} , and consider the matroid $\varphi(\mathcal{N})$ provided by Theorem 12. As \mathcal{N} is aent, $\varphi(\mathcal{N})$ is almost entropic; \mathcal{N}^\perp is not aent, thus $\varphi(\mathcal{N}^\perp)$ is not aent. Consequently the matroid $\varphi(\mathcal{N})$ is aent and its dual, $\varphi(\mathcal{N}^\perp)$, is not aent either – completing the proof of Theorem 21.

Using the tight almost-entropic polymatroid of Table 2 the construction in the proof of Theorem 19 gives an almost-ideal access structure on 174 participants (as the corresponding aent matroid has $f(a) + f(b) + f(c) + f(d) + f(e) = 175$ atoms, one of them is the secret, others are the participants) whose dual is not almost-ideal. It is left to the interested reader to describe the qualified subsets for different choices of the secret.

According to Theorem 20, to construct a counterexample to Conjecture 2 we need an entropic matroid whose dual is not entropic. Entropic matroids (and their multiplies) are always on the boundary of the aent cone; the boundary has an intricate and complicated structure. There seems to be no other way to show that a matroid is entropic than giving the probability distribution explicitly. But it is not clear how to guarantee $\mathbf{H}(\xi_A)/\mathbf{H}(\xi_s)$ to be an integer. No entropic matroid is known which is not a multiple of a linearly representable polymatroid. Finding such a matroid would be very interesting.

Acknowledgment

The author would like to thank the encouragement and fruitful discussion at the Prague Stochastics Conference, 2019. He is particularly indebted to Oriol

Farrás and Carles Padró. Special thanks to Gabor Tardos for figuring out the subtleties of Claim 5.

The research reported in this paper was supported by GACR project number 19-04579S, and partially by the Lendület program of the HAS.

Appendix

Theorem. *A matroid is connected if and only if any two points can be connected by a circuit.*

Proof. Let $\mathcal{M} = (h, M)$ be a matroid. Using matroid terminology, A is *dependent* if $h(A) < |A|$, and A is a *circuit* if it is a minimal dependent set. Every dependent set contains a circuit. Points x and y are *connected*, written as $a \approx b$, if there is a circuit containing both of them. First we prove that \approx is an equivalence relation: if $x \approx z$ and $z \approx y$ then $x \approx y$. This is done in three steps. In claims a), b), c), C_1 and C_2 are circuits.

a) *Suppose $z \in C_1 \cap C_2$. There is a circuit $E \subseteq C_1 \cup C_2$ which avoids z (exchange property of circuits):*

Proof. As C_1, C_2 are circuits, $h(C_i) = |C_i| - 1$. Using submodularity,

$$\begin{aligned} h(C_1 \cup C_2) &\leq h(C_1) + h(C_2) - h(C_1 \cap C_2) = \\ &= |C_1| + |C_2| - h(C_1 \cap C_2) - 2 \leq |C_1| + |C_2| - |C_1 \cap C_2| - 2 = \\ &= |C_1 \cup C_2| - 2. \end{aligned}$$

Consequently $h(C_1 \cup C_2 - z) \leq |C_1 \cup C_2 - z| - 1$, which means that $C_1 \cup C_2 - z$ is dependent, thus contains a circuit.

b) *Let $x \in C_1 - C_2$, and $z \in C_1 \cap C_2$. There is a circuit in $C_1 \cup C_2$ which contains x and avoids z .*

Proof. By induction on $|C_1 \cup C_2|$. By a) there is a circuit $E \subseteq C_1 \cup C_2$ which avoids z . Then $E \cap (C_2 - C_1)$ is not empty, as E is dependent while $E \cap C_1$, as a proper subset of C_1 , is independent. If $x \in E$ then we are done. If $x \notin E$, then pick $z' \in E \cap (C_2 - C_1)$. By a) there is circuit $F \subseteq E \cup C_2$ which avoids z' . Use induction on C_1 and F .

c) *Let $x \in C_1 - C_2$, $y \in C_2 - C_1$, and $C_1 \cap C_2 \neq \emptyset$. There is a circuit $E \subseteq C_1 \cup C_2$ which contains x and y .*

Proof. By induction on $|C_1 \cup C_2|$. Let $z \in C_1 \cap C_2$. By b) there is a circuit $E \subseteq C_1 \cup C_2$ which contains x and avoids z . If $y \in E$, then we are done. If $y \notin E$, then pick $z' \in E \cap (C_2 - C_1)$. By b) there is a circuit $F \subseteq E \cup C_2$ such that $y \in F$ and $z' \notin F$. Use induction on C_1 and F .

This proves that \approx is an equivalence relation. Any two points of the matroid are connected by a circuit if and only if there is only a single equivalence class for \approx . First assume that the matroid is connected, and by contradiction that A is a proper equivalence class of \approx . Consider the partition $A \cup B$ where B is the complement of A . Choose the independent sets $A' \subseteq A$ and $B' \subseteq B$ such that $h(A) = |A'|$ and $h(B) = |B'|$. As A and B are not independent,

$h(A') + h(B') = h(A) + h(B) > |A'| + |B'|$, thus $A' \cup B'$ contains a circuit E . But E must intersect both A' and B' (as A' and B' are independent), contradicting that elements from $A' \subseteq A$ and from $B' \subseteq B$ are not connected.

Conversely, if the matroid is not connected, say the elements of the partition $A \cup B$ are independent, then no circuit can intersect both A and B . Indeed, first the independence of A and B implies $h(E) = h(E \cap A) + h(E \cap B)$ for all subsets $E \subseteq M$. Second, assume the circuit E intersects both A and B . Then $E \cap A$ and $E \cap B$ are independent (as proper subsets of E), and then

$$h(E) = h(E \cap A) + h(E \cap B) = |E \cap A| + |E \cap B| = |E|,$$

a contradicting that E is dependent. \square

Theorem. *If $\xi = \langle \xi_a, \dots, \xi_e \rangle$ is a distribution on five elements, then the polymatroid \mathcal{M}_ξ satisfies the MMRV inequality*

$$\begin{aligned} & (h(a, b|c) + h(b, c|a) + h(c, a|b)) + \\ & + (h(b, c|d) + h(b, c|e) + h(d, e) - h(b, c)) \geq 0, \end{aligned} \quad (2)$$

written as $\text{MMRV}(\mathcal{M}_\xi) \geq 0$.

Proof. Observe first that in *any* polymatroid $\mathcal{M} = (h, M)$ the inequality

$$\text{MMRV}(\mathcal{M}) + 3h(a, de|bc) \geq 0$$

always holds. This is so as expanding $\text{MMRV}(\mathcal{M}) + 3h(a, de|bc)$ as a linear combination of rank values, and expanding the clearly non-negative sum below, the results are the same:

$$\begin{aligned} & h(a, d|b) + h(a, d|c) + h(a, e|b) + h(a, e|c) + \\ & + h(b, c|ad) + h(b, c|ae) + h(a, bc|de) + \\ & + h(d, e|a) + h(a, e|bcd) + h(a, d|bce). \end{aligned}$$

The MMRV inequality (2) has been grouped into two parts. The first part depends only on ranks of subsets of abc , and the second part depends only on subsets of bcd . In other words, the value of the first (and second) part depends only on the marginal distribution ξ_{abc} and ξ_{bcd} , respectively. Σ denotes the collection of all distributions $\eta = \langle \eta_a, \dots, \eta_e \rangle$ where each of these five variables takes the same values as the corresponding variable does in ξ but with arbitrary joint probability. Consider the optimization problem of maximizing the entropy of $\eta \in \Sigma$ under the constraints that certain marginal distributions are fixed:

$$\max_{\eta} \{ \mathbf{H}(\eta) : \eta \in \Sigma, \eta_{abc} = \xi_{abc}, \eta_{bcd} = \xi_{bcd} \}.$$

As $\mathbf{H}(\eta)$ is a strictly convex function of the probabilities, this is a convex optimization problem with linear constraints, consequently it has a single unique optimal solution $\eta^* \in \Sigma$. Considering the distribution with the maximal entropy is often referred to as the *maximum entropy principle*. As the marginals

on abc and bcd of ξ and η^* are the same, $\text{MMRV}(\mathcal{M}_\xi) = \text{MMRV}(\mathcal{M}_{\eta^*})$. The extremal distribution η^* has the additional property that η_a^* and η_{de}^* are independent given η_{bc}^* . This is so, as fixing the value of η_{bc}^* , one can redefine the distribution while keeping the probabilities on abc and on bcd fixed such that a and de becomes independent. This would increase the total entropy, thus a and de must be independent – giving the claimed conditional independence. Consequently the polymatroid \mathcal{M}_{η^*} satisfies additionally $h^*(a, de|bc) = 0$, and then $\text{MMRV}(\mathcal{M}_{\eta^*}) \geq 0$, proving the theorem. \square

References

- [1] A. Beimel (2011), Secret-sharing schemes: a survey, in: *IWCC 2011, volume 6639 of LNCS*, Springer, 2011, pp 11-46
- [2] A. Beimel, N. Livne (2006) On matroids and non-ideal secret sharing In: Halevi S., Rabin T. (eds) *Theory of Cryptography, volume 3876 of LNCS*, Springer, Berlin, Heidelberg pp 482-501
- [3] G. Blakley, G. Kabatianski (1995), On general perfect secret sharing schemes, in: *LNCS 963, Advances in Cryptology, Proceedings of Crypto'95*, Springer 1995, pp. 367–371
- [4] S. Fujishige (1978), Polymatroidal dependence structure of a set of random variables. *Information and Control* **39** 55–72.
- [5] T. Helgason (1974) Aspects of the theory of hypermatroids, In: Berge C., Ray-Chaudhuri D. (eds) *Hypergraph Seminar*, Lecture Notes in Mathematics, vol 411. Springer, Berlin, Heidelberg
- [6] W. C. Huffman and V. Pless (2003), *Fundamentals of error correcting codes*, Cambridge University Press, 2003
- [7] T. Kaced (2018), Information Inequalities are Not Closed Under Polymatroid Duality, *IEEE Transactions on Information Theory*, 64, pp 4379–4381
- [8] J. Katz, Y. Lindell (2007), *Introduction to modern cryptography*, Chapman & Hall/CRC
- [9] L. Lovász (1982), Submodular functions and convexity. *Mathematical Programming – The State of the Art* (A. Bachem, M. Grötschel and B. Korte, eds.), Springer-Verlag, Berlin, 234–257.
- [10] K. Makarichev, Y. Makarichev, A. Romashchenko, N. Vereshchagin (2002), A new class of non-Shannon type inequalities for entropies. *Communications in Information and Systems*, vol 2, pp 147–166
- [11] F. Matúš (1994), Probabilistic conditional independence structures and matroid theory: background. *Int. Journal of General Systems* **22** 185–196.

- [12] F. Matúš (2007), Adhesivity of polymatroids, *Discrete Mathematics* **307** pp. 2464–2477
- [13] F. Matúš (2007), Two constructions on limits of entropy functions. *IEEE Transactions on Information Theory* 53, pp 320-330.
- [14] F. Matúš (2007), Infinitely many information inequalities. *Proceedings IEEE ISIT 2007*, Nice, France, pp 41–44.
- [15] F. Matúš (2012), Polymatroids and polyquantoids. in: *Proceedings of WU-PES'2012* (eds. J. Vejnarová and T. Kroupa) Mariánské Lázně, Prague, Czech Republic, pp 126-136.
- [16] F. Matúš, L. Csirmaz (2016), Entropy region and convolution, *IEEE Trans. Inf. Theory* **62** 6007–6018
- [17] J.G. Oxley (1992) *Matroid Theory*, Oxford Science Publications. The Clarendon Press, Oxford University Press, New York
- [18] C. Padró (2012), Lecture notes in secret sharing, *Cryptology ePrint archive*, report 2012/674
- [19] R.W. Yeung (2002), *A First Course in Information Theory*, Kluwer Academic/Plenum Publishers, New York.