

# On Rejection Sampling in Lyubashevsky’s Signature Scheme

Julien Devevey<sup>1</sup>, Omar Fawzi<sup>1,2</sup>, Alain Passelègue<sup>1,2</sup>, and Damien Stehlé<sup>1,3</sup>

<sup>1</sup> ENS de Lyon, Lyon, France

<sup>2</sup> Inria Lyon, Lyon, France

<sup>3</sup> Institut Universitaire de France, Paris, France

**Abstract.** Lyubashevsky’s signatures are based on the Fiat-Shamir with aborts paradigm, whose central ingredient is the use of rejection sampling to transform secret-dependent signature samples into samples from (or close to) a secret-independent target distribution. Several choices for the underlying distributions and for the rejection sampling strategy can be considered. In this work, we study Lyubashevsky’s signatures through the lens of rejection sampling, and aim to minimize signature size given signing runtime requirements. Several of our results concern rejection sampling itself and could have other applications.

We prove lower bounds for compactness of signatures given signing runtime requirements, and for expected runtime of perfect rejection sampling strategies. We also propose a Rényi-divergence-based analysis of Lyubashevsky’s signatures which allows for larger deviations from the target distribution, and show hyperball uniforms to be a good choice of distributions: they asymptotically reach our compactness lower bounds and offer interesting features for practical deployment. Finally, we propose a different rejection sampling strategy which circumvents the expected runtime lower bound and provides a worst-case runtime guarantee.

## 1 Introduction

<sup>1</sup>Lyubashevsky’s signature scheme [Lyu09,Lyu12] may be viewed as a lattice variant of Schnorr’s group-based signature scheme [Sch91], with a core conceptual difference being the use of rejection sampling and the associated introduction of aborts and repeats in the Fiat-Shamir heuristic [FS86]. The use of rejection sampling in Lyubashevsky’s scheme is the focus of the present work. It is hard to overstate the importance of Lyubashevsky’s signature scheme in lattice-based cryptography. Thanks to its elementary and flexible design, numerous variants and optimizations have been proposed (see [AFLT16, GLP15, DDLL13, BG14], or [Lyu16], for instance). Notably, it led to the TESLA [ABB<sup>+</sup>17, AAB<sup>+</sup>19] and Dilithium [DKL<sup>+</sup>18, BDK<sup>+</sup>20] candidates to the NIST standardization project on post-quantum cryptography. It also led to lattice-based zero-knowledge proofs (see [LNP22] and the references therein).

---

<sup>1</sup> A shorter version of this article appeared in the proceedings of Asiacrypt’22. This is the full version, dated Dec. 5, 2022.

Lyubashevsky’s scheme involves a publicly shared matrix  $\mathbf{A} \in \mathbb{Z}_q^{n \times m}$  (note that other algebraic setups are possible, but this is not relevant to the present discussion). The signing key is a matrix  $\mathbf{S} \in \mathbb{Z}^{m \times k}$ . It is small in the sense that all its entries have absolute values significantly smaller than  $q$ . The verification key associated to  $\mathbf{S}$  is  $\mathbf{T} = \mathbf{AS}$ . Given a message  $\mu \in \{0, 1\}^*$ , the signer samples a small masking vector  $\mathbf{y} \in \mathbb{Z}^m$  and computes a random-looking commitment  $\text{com} = \mathbf{Ay}$ . By using a hash function  $H$  taking small values in  $\mathbb{Z}^k$ , it computes a challenge  $\mathbf{c} = H(\text{com}, \mu)$ . Finally, if some (possibly probabilistic) test passes, it outputs a signature  $\sigma = (\mathbf{z}, \mathbf{c})$  with  $\mathbf{z} = \mathbf{y} + \mathbf{Sc}$ , and else it restarts from scratch. Given a signature  $\sigma = (\mathbf{z}, \mathbf{c})$  for a message  $\mu$ , the verifier accepts if and only if  $\mathbf{z}$  is small and  $H(\mathbf{Az} - \mathbf{Tc}, \mu) = \mathbf{c}$ . We refer the reader to Figure 2 for a formal description. As suggested by the choice of terminology, Lyubashevsky’s signature can be viewed as an identification protocol made non-interactive by relying on the Fiat-Shamir heuristic, i.e., by replacing a truly random  $\mathbf{c}$  by the output of a hash function. The security proof relies on the Random Oracle Model (ROM) as it models  $H$  as a function such that each image is distributed as  $\mathbf{c}$  is supposed to be.

Compared to Schnorr’s signature scheme, the signing key and mask do not belong to a finite set, preventing the use of a uniform mask  $\mathbf{y}$  to hide the sensitive term  $\mathbf{Sc}$ .<sup>2</sup> One possibility (see, e.g., [DPSZ12]) is to sample  $\mathbf{y}$  exponentially larger than  $\mathbf{Sc}$  as a function of the security parameter, so that the distributions of  $\mathbf{y}$  and  $\mathbf{y} + \mathbf{Sc}$  have exponentially small statistical distance. As  $q$  must be larger than  $\mathbf{y}$  and the smallness of  $\mathbf{S}$  relative to  $q$  impacts security, this flooding approach leads to large parameters. Instead, Lyubashevsky [Lyu09, Lyu12] put forward the notion of Fiat-Shamir with aborts. This is the reason for the test concerning  $\mathbf{z}$  in the signing algorithm: it is so that the output signature  $(\mathbf{z}, \mathbf{c})$  follows a distribution that is independent of the sensitive term  $\mathbf{Sc}$ .

A classic application of rejection sampling (see, e.g., [Dev86, Chapter 2]) is to use a source distribution  $Q$  that is convenient to sample from, to create samples from a target distribution  $P$ . In Lyubashevsky’s scheme, the purpose differs: we start from a pre-source distribution  $Q$  for  $\mathbf{y}$ ; it is shifted by  $\mathbf{Sc}$ , leading to a distribution  $Q_{+\mathbf{Sc}}$  for  $\mathbf{y} + \mathbf{Sc}$ ; the latter is the source distribution; it is rejected to a target distribution  $P$  for  $\mathbf{z}$  that does not depend on the signing key  $\mathbf{S}$ . The purpose of rejection sampling here is to hide the sensitive data  $\mathbf{Sc}$ . Diverse choices of pairs of distributions have been put forward in the literature: uniform in hypercubes [Lyu09], Gaussian with the same standard deviation while allowing for some small statistical inaccuracy in the target distribution [Lyu12], a bimodal Gaussian source distribution with a Gaussian target distribution in association with an accommodating arithmetic modification of the scheme [DDLL13] (the modification consists in replacing  $q$  with  $2q$  and changing key generation to ensure that  $\mathbf{T} = -\mathbf{T} = \mathbf{qI} \pmod{2q}$ ). The pre-source distribution  $Q$  is shifted by  $(-1)^b \mathbf{Sc}$  for a uniform bit  $b$ , leading to a source distribution  $Q_{\pm \mathbf{Sc}}$ . We refer

<sup>2</sup> If we view  $\mathbf{y}$  and  $\mathbf{S}$  over  $\mathbb{Z}_q$  rather than  $\mathbb{Z}$ , then they do belong to a finite set; but for security, the masking should preserve smallness relative to  $q$ , which the uniform distribution modulo  $q$  does not achieve.

to this as the bimodal setting. By opposition, we now refer to the former two cases where the source distribution is  $Q_{+\mathbf{s}_c}$  as the unimodal setting. The first choice (uniform distributions in hypercubes) leads to a simple design, whereas the latter two allow for more compact signatures. One may also want to add constraints on the number of loop iterations, notably to guarantee a signing runtime upper bound. In the extreme case of removing rejection altogether, it was recently shown in [ASY22] that a limited flooding suffices, compared to the exponential flooding discussed earlier. This leads us to the question we address in this work:

*Given signing runtime requirements, which rejection sampling strategy leads to the most compact signatures?*

In a signature  $\sigma = (\mathbf{z}, \mathbf{c})$ , the second component contributes to a small fraction of the bitsize: the main requirement on  $\mathbf{c}$  is that it has sufficiently high min-entropy to make it hard to guess. On the other hand, the contribution of  $\mathbf{z}$  towards signature length is mostly driven by  $\|\mathbf{z}\|$ , as this directly impacts security: for a given security level, the smaller  $\|\mathbf{z}\|$ , the more compact the signatures. For this reason, we simplify the overall objective to minimizing  $\mathbb{E}_{\mathbf{x} \leftarrow P}(\|\mathbf{x}\|)$  under signing runtime requirements.

**Contributions.** Our main contributions concern the optimality of rejection sampling design choices towards optimizing signature sizes and signing runtime. We provide lower bounds, and study ways to reach and circumvent them.

Before describing the main results, we need to quantify the runtime of rejection sampling strategies. We note that for classic rejection sampling with target  $P$  and source  $Q$ , the expected number of samples needed is  $R_\infty(P\|Q)$  where  $R_\infty(D_1\|D_2) = \sup_x D_1(x)/D_2(x)$  refers to the Rényi divergence of infinite order. Indeed, for classic rejection sampling, one samples  $x$  from  $Q$  and accepts with probability  $P(x)/(M \cdot Q(x))$ , for  $M = R_\infty(P\|Q)$ . This justifies using  $R_\infty(P\|Q)$  to quantify the runtime for rejecting  $Q$  to  $P$ .

We start with our lower bounds.

- Considering Lyubashevsky’s scheme with perfect rejection sampling to the target distribution  $P$  (as in [Lyu09]), the relevant quantity measuring the signing runtime is then given by  $M = \max_{\mathbf{s}, \mathbf{c}} R_\infty(P\|Q_{+\mathbf{s}_c})$ . We show (under a mild assumption discussed below) that for all  $P$  and  $Q$  such that  $M$  is finite, the expected norm  $\mathbb{E}_{\mathbf{x} \leftarrow P}(\|\mathbf{x}\|)$  is  $\Omega((m/\log M) \cdot \max_{\mathbf{s}, \mathbf{c}} \|\mathbf{S}\mathbf{c}\|)$ . Interestingly, this bound is a factor  $\sqrt{m}$  lower than what is obtained for the typical choice of  $P$  and  $Q$  set as uniform distributions in hypercubes.
- In the case of perfect rejection with the accommodating arithmetic modification from [DDLL13], then the relevant quantity for measuring the signing runtime is  $M = \max_{\mathbf{s}, \mathbf{c}} R_\infty(P\|Q_{\pm\mathbf{s}_c})$ , where  $Q_{\pm\mathbf{s}_c}$  denotes the balanced mixture of  $Q_{+\mathbf{s}_c}$  and  $Q_{-\mathbf{s}_c}$ . In this case, we show (under the same mild assumption) that for all  $P$  and  $Q$  such that  $M$  is finite, the expected norm  $\mathbb{E}_{\mathbf{x} \leftarrow P}(\|\mathbf{x}\|)$  is  $\Omega(\sqrt{m/\log M} \cdot \max_{\mathbf{s}, \mathbf{c}} \|\mathbf{S}\mathbf{c}\|)$ . This lower bound is actually reached (up to a constant factor) for  $P$  and  $Q$  Gaussian as in [DDLL13].

- We show that for any algorithm (terminating with probability 1) that selects one out of many samples from  $Q$  to get a sample from  $P$ , the expected number of required samples from  $Q$  is  $\geq R_\infty(P\|Q)$ . This lower bound is reached by classic rejection sampling. In the case of Lyubashevsky’s signatures with exact rejection sampling, this general result implies that classic rejection sampling is the appropriate strategy when it comes to minimize the expected runtime.

The lower bounds above seem to give little margin of improvement in the design choices of Lyubashevsky’s signatures, except for the unimodal case, for which uniform distributions in hypercubes do not reach the lower bound. Our second set of main results considers ways to reach or circumvent these lower bounds.

- Concerning the unimodal case, one way to circumvent the results above is to consider imperfect rejection sampling, by allowing for an approximation to  $P$  whose accuracy is parameterized by some  $\varepsilon > 0$  (as introduced in [Lyu12]). Then the relevant quantity to bound the runtime becomes  $\max_{\mathbf{S}, \mathbf{c}} R_\infty^\varepsilon(P\|Q_{+\mathbf{S}\mathbf{c}})$ , where  $R_\infty^\varepsilon$  is a smoothed variant of  $R_\infty$  that we define. In this case, we improve the signature security analysis from [Lyu12] by using the Rényi divergence instead of the statistical distance to quantify the closeness to  $P$  of the output distribution. This allows choosing  $\varepsilon$  larger than previously, leading to a (limited) signature compactness improvement.
- Gaussian distributions provide better signature compactness in the bimodal and imperfect unimodal regimes, than uniforms in hypercubes in the perfect unimodal regime. However, uniforms in hypercubes are sometimes preferred (see, e.g., Dilithium), because they lead to a simpler implementation, which in turn makes protection against timing attacks easier. We consider uniforms in hyperballs as a new alternative for the choice of source and target distributions. We show that this choice reaches the two lower bounds for  $\mathbb{E}_{\mathbf{x} \leftarrow P}(\|\mathbf{x}\|)$  for perfect rejection sampling and is as good as Gaussians for imperfect rejection sampling in practice in the imperfect regime. Interestingly, the rejection test for uniforms in hyperballs is very simple in the unimodal case, similarly to uniforms in hypercubes. We not only study the choice of uniforms in hyperballs in the asymptotic regime, but also compare it to Dilithium.
- Finally, imperfect rejection from  $Q$  to  $P$  allows us to describe and analyze variants of rejection sampling where the maximum number of loop iterations is bounded. This provides trade-offs between maximum signing runtime and signature sizes. When instantiated to rejection-free sampling, we recover the scheme and analysis from [ASY22], whereas it quickly converges to Lyubashevsky’s signature scheme when the signing runtime bound grows.

The results concerning signature compactness for unbounded (perfect and imperfect) rejection sampling are summarized in Table 1.

**Technical overview.** In Section 2, we provide the background necessary to this work, including rejection sampling and Lyubashevsky’s signature scheme.

After identifying the notion of expected number of iterations during rejection sampling with the notion of smooth-Rényi divergence that we define, we start

	Unimodal ( $\varepsilon = 0$ )	Unimodal ( $\varepsilon \geq 2^{-o(m)}$ and $\varepsilon = o(1/m)$ )	Bimodal ( $\varepsilon = 0$ )
Hypercube	$\frac{tm^{3/2}}{\log M}$	$\frac{tm^{3/2}}{\log M}$	$\frac{tm^{3/2}}{\log M}$
Gaussian	$\infty$	$\frac{t\sqrt{m}\sqrt{\log \frac{1}{\varepsilon} + \log M}}{\log M}$	$\frac{t\sqrt{m}}{\sqrt{\log M}}$
Hyperball	$\frac{tm}{\log M}$ (Lemma 5.1)	$\frac{t\sqrt{m}\sqrt{\log \frac{1}{\varepsilon} + \log M}}{\log M}$ (Lemma 5.1)	$\frac{t\sqrt{m}}{\sqrt{\log M}}$ (Lemma 5.2)
Lower bound	$\frac{tm}{\log M}$ (Corollary 3.3)	?	$\frac{t\sqrt{m}}{\sqrt{\log M}}$ (Corollary 3.5)

**Table 1.** This table expresses the compactness of the signature modeled as  $\mathbb{E}_{\mathbf{x} \leftarrow P}(\|\mathbf{x}\|)$  given the signing runtime constraint for various choices of distributions  $P$  and  $Q$ . The column indicates the signing runtime constraint which is modeled in the unimodal case by  $\max_{\mathbf{v} \in \mathcal{B}_m(t)} R_\infty^\varepsilon(P\|Q_{+\mathbf{v}}) \leq M$  where  $\varepsilon$  quantifies the accuracy of rejection sampling and in the bimodal case by  $\max_{\mathbf{v} \in \mathcal{B}_m(t)} R_\infty(P\|Q_{\pm\mathbf{v}}) \leq M$ . In the first row,  $P$  and  $Q$  are chosen to be uniform in  $m$ -dimensional hypercubes of appropriate side-lengths, in the second row, they are chosen to be  $m$ -dimensional Gaussians of appropriate variance. In the third row, they are chosen to be uniform in the  $m$ -dimensional hyperballs of appropriate radii. The last row gives a lower bound on the compactness for any choice of  $P$  and  $Q$ . Multiplicative constants are omitted in this table, and we make the assumption that  $\log M \leq m$ .

addressing our main question of understanding to which extent the expected norm of a signature can be small for target expected signing runtime constraints. *Lower Bounds.* In Section 3, we prove lower bounds in the case of exact rejection sampling in both unimodal and bimodal settings. These lower bounds are obtained following a similar path. In what follows, we focus on the unimodal setting. To ease the analysis, we place ourselves in a slightly simplified setup where shifts belong to a hyperball  $\mathcal{B}_m(t)$  of radius  $t$  instead of being defined as  $\mathbf{S}\mathbf{c}$ . Given that  $\mathbf{S}$  is unknown, this simplification seems reasonable and allows avoiding significant complications in the proof. In this setting, we prove that for a given constraint  $\max_{\mathbf{v} \in \mathcal{B}_m(t)} R_\infty(P\|Q_{+\mathbf{v}}) \leq M$ , we have  $\mathbb{E}_{\mathbf{x} \leftarrow P}(\|\mathbf{x}\|) \geq (t/M^{1/(m-1)} - 1) - \sqrt{m}/2$ .

Our lower bounds are obtained in three steps: (1) considering the same setting with continuous distributions, we first prove that we can restrict ourselves to the case of isotropic distributions over  $\mathbb{R}^m$ , where isotropic means that their densities only depend on the norm. Specifically, we prove that for any two densities  $f, g$ , there exist isotropic distributions  $f^*, g^*$  satisfying  $\max_{\mathbf{v} \in \mathcal{B}_m(t)} R_\infty(f^*\|g^*_{+\mathbf{v}}) \leq M$  as well as  $\mathbb{E}_{\mathbf{x} \leftarrow f^*}(\|\mathbf{x}\|) = \mathbb{E}_{\mathbf{x} \leftarrow f}(\|\mathbf{x}\|)$ . The latter distributions are essentially obtained from  $f, g$  by averaging their respective densities on hyperspheres. (2) Starting with  $f$  and  $g$  isotropic, we show that  $\mathbb{E}_{\mathbf{x} \leftarrow f}(\|\mathbf{x}\|) = \mu_m/\mu_{m-1}$  where  $\mu_k = \int_0^\infty r^k f(r) dr$ . The main technicality consists in proving an intermediate lower bound  $\mu_{m-1}/\mu_{m-2} \geq (t/M^{1/(m-1)} - 1)$  which results from the constraint  $\max_{\mathbf{v} \in \mathcal{B}_m(t)} R_\infty(f\|g_{+\mathbf{v}}) \leq M$ . Our lower bound is then obtained by applying the Cauchy-Schwarz inequality  $|\mathbb{E}(XY)|^2 \leq \mathbb{E}(X^2)\mathbb{E}(Y^2)$  to random variables  $X = \|\mathbf{x}\|^{m/2}$  and  $Y = \|\mathbf{x}\|^{(m-2)/2}$ , where  $\mathbf{x} \leftarrow f$ . Indeed, it immediately leads

to inequality  $\mu_m \cdot \mu_{m-2} \geq (\mu_{m-1})^2$ , which results in  $\mu_m/\mu_{m-1} \geq \mu_{m-1}/\mu_{m-2} \geq (t/M^{1/(m-1)} - 1)$ . (3) A similar lower bound in the discrete setting is obtained by considering the continuous density  $p(\mathbf{x}) = P(\lceil \mathbf{x} \rceil)$  with  $P$  being a discrete probability. These lower bounds provide us with a target to reach, and we can compare them with the signature size obtained when instantiating the above scheme with various distributions.

*On Alternative Rejection Sampling Strategies.* In Section 3.3, we investigate the question of the relevance of rejection sampling strategies differing from the classic one. We consider the following setting. As above, the goal is to sample from a distribution  $P$  given access to a sampler from a distribution  $Q$ , and we consider a sequence of samples  $(X_i)_{i \geq 1}$  from distribution  $Q$ . Any strategy is allowed as long as we output one of the  $X_i$ 's. A strategy is given by a sequence of algorithms  $(A_i)_{i \geq 1}$  that take samples  $(X_j)_{j \leq i}$  as input and return either an index  $j \in [i]$ , which corresponds to halting with output  $X_j$ , or a special symbol  $r$  which corresponds to rejecting and moving to  $A_{i+1}$ . We restrict ourselves to the case of procedures that terminate with probability 1. Considering  $i^*$  the random variable denoting the number of samples observed in a strategy, our objective is then to measure how small  $\mathbb{E}(i^*)$  can be. We prove that for any  $P, Q$ , we have  $\mathbb{E}(i^*) \geq R_\infty(P||Q)$ . This result is obtained by proving that for any  $x$ , we have  $P(x) \leq \mathbb{E}(i^*) \cdot Q(x)$ , leading to the former inequality by definition of  $R_\infty$ .

*Rényi-Based Analysis for Imperfect Rejection Sampling.* All lower bounds are for perfect rejection sampling, in the sense that one obtains a sample from (exactly)  $P$ . In [Lyu12], Lyubashevsky showed that one can consider imperfect rejection sampling, and shows that it is particularly beneficial in the case of Gaussians. We propose an analysis that replaces the use of the statistical distance as done in [Lyu12] by that of the smooth Rényi divergence, and allows loosening the constraints on imperfectness. We first recall that in [Lyu12], the statistical distance is used to bound the statistical distance between a (single) execution of the imperfect rejection sampling algorithm and the target distribution. Using imperfect rejection sampling in a signature scheme and given bound  $\varepsilon$  for the above statistical distance, one can then bound the distinguishing advantage of an adversary between the real security game and the ideal game (where signatures are simulated by sampling them from the target distribution) by  $q_{\text{sig}} \cdot \varepsilon$ . Here  $q_{\text{sig}}$  is a bound on the number of signature queries an adversary can make. In Section 4, we prove that for  $P, Q$  such that  $R_\infty^\varepsilon(P||Q)$  is finite, the Rényi divergence of infinite order between a (single) execution of the imperfect rejection sampling algorithm and the target distribution is bounded by  $1/(1 - \varepsilon)$ . Combining this result with the multiplicativity of the Rényi divergence, we can then bound the Rényi divergence of infinite order between the adversary's view in the real game and its view in the ideal game by  $1/(1 - \varepsilon)^{q_{\text{sig}}}$  for the resulting signature. The probability preservation property of the Rényi divergence then allows completing the analysis. Our analysis leads to potential improvements as the former statistical bound  $q_{\text{sig}} \cdot \varepsilon$  imposes that  $\varepsilon = 2^{-\Omega(\lambda)}$ , while our bound can be used setting  $\varepsilon = 1/q_{\text{sig}}$ . Since  $q_{\text{sig}}$  is a (possibly large) polynomial of the

security parameter  $\lambda$ , this puts less constraint on the condition  $P$  and  $Q$  must satisfy, which results in compactness improvement.

*Hyperball Uniforms.* We show that (continuous) uniform distributions over hyperballs reach the signature compactness lower bound (up to constant factors) in both unimodal and bimodal settings, as shown in Section 5.1. We also show that they are experimentally as good as Gaussians for imperfect rejection sampling. These results reduce to Rényi divergence computations, which involve geometric properties of hyperballs. We emphasize that while Gaussian distributions also achieve similar signature size in both unimodal and bimodal settings (but only in the case of imperfect rejection sampling with polynomial loss for the unimodal case), using uniform distributions over hyperballs makes the rejection test as simple as computing  $\|\mathbf{z}\|$  in the unimodal case since it consists only in checking that  $\mathbf{z}$  is in the hyperball of the target distribution  $P$ . In the bimodal case, the rejection test involves computing two norms and flipping a coin. In order to use this distribution in a signature, we propose a generalization of Lyubashevsky’s signature that allows for continuous source and target distributions, by adding a rounding step after accepting a sample. Its security relies on the same mechanisms as the discrete case. This strategy could also benefit to Gaussian distributions, by allowing to replace discrete Gaussian sampling with possibly simpler continuous Gaussian sampling. To assess the practicality of this new choice of distributions, we propose parameters for a variant of Dilithium with uniform distributions in hyperballs. If considering the sum of bitsizes of a verification key and a signature, the gains range from  $\sim 25\%$  to  $\sim 30\%$ , depending on the security level, just like for the Gaussian variant, whose parameters we also update.

*Bounded Rejection Sampling.* We conclude this work by proposing an original strategy to use rejection sampling while guaranteeing a (moderate) worst-case runtime. This could be beneficial in the context of real-time systems. A simple strategy could consist in fixing a (very large) bound  $i$  on the number of iterations such that it fails to produce a sample with negligible probability. While this guarantees a worst-case runtime, the change is mainly cosmetic since it has to be large enough for the sampling to succeed. In Section 6, we propose an alternative solution that leaves the choice of  $i$  open without ever failing: for a fixed bound  $i$ , it performs (up to)  $i - 1$  iterations of the classic rejection sampling and outputs a sample if it ever succeeds, otherwise, the last ( $i$ -th) iteration uses one-shot flooding techniques (as done in [ASY22]) to guarantee an output. The analysis makes heavy use of the smooth Rényi divergence and its properties. Different choices for the bound  $i$  offer various trade-offs, ranging from one-shot signatures ( $i = 1$ ) as in [ASY22] to Lyubashevsky’s expected polynomial-time signatures ( $i$  going to  $\infty$ ).

**Open problems.** Our results suggest that instantiating the Fiat-Shamir with aborts using uniform distributions in hyperballs is a relevant choice, both in the unimodal and bimodal settings, as it provides more compact signatures than uniform distributions in hypercubes but also much simpler rejection test than Gaussians. We believe it is an interesting open question to investigate a constant-time

implementation with this choice. Regarding further improvements of signatures, our results show that there is not much room for improvement if the goal is to minimize signature size or  $\mathbb{E}(i^*)$ . However, other quantities could be considered, such as the shape of the tail of the distribution of  $i^*$ .

## 2 Preliminaries

Due to space limitations, we postpone notations and some standard background to Appendix A. This includes definitions pertaining to digital signatures and to specific distributions.

We introduce a relaxed version of the Rényi divergence, termed the smooth Rényi divergence, where one is able to remove a few problematic points from the support, including those that may lie in  $\text{Supp}(p) \setminus \text{Supp}(q)$ . Doing so, we can compare a wider set of probability distributions. For instance, while the Rényi divergence of infinite order between  $D_{\mathbb{Z}^m, \sigma}$  and  $D_{\mathbb{Z}^m, \sigma, \mathbf{v}}$  is infinite when  $\mathbf{v} \neq \mathbf{0}$ , their smooth divergence is finite, as we show in Lemma C.2 and is implicit in [Lyu12]. We could give this definition for any order  $a \in [1, +\infty]$ . However, only the case  $a = +\infty$  is relevant for this work.

This definition is useful to link previous works on rejection sampling and the Rényi divergence. A similar quantity has been previously defined in the quantum information literature [Ren05, Dat09], though the specific notion of smoothing we consider here is slightly different.

**Definition 2.1 (Smooth Rényi Divergence).** *Let  $\varepsilon \geq 0$ . Let  $p, q$  be two probability densities such that  $\int_{\text{Supp}(q)} p(x) d\mu(x) \geq 1 - \varepsilon$ . Their  $\varepsilon$ -smooth Rényi divergence of infinite order is*

$$R_{\infty}^{\varepsilon}(p||q) := \inf_{\substack{S \subseteq \text{Supp}(q) \\ \int_S p(x) d\mu(x) \geq 1 - \varepsilon}} \text{ess sup}_{x \in S} \frac{p(x)}{q(x)}.$$

*This definition is equivalent to*

$$R_{\infty}^{\varepsilon}(p||q) := \inf\{M > 0 \mid \Pr_{x \leftarrow p}(p(x) \leq Mq(x)) \geq 1 - \varepsilon\}.$$

*By convention, if  $\int_{\text{Supp}(q)} p(x) d\mu(x) < 1 - \varepsilon$ , we define  $R_{\infty}^{\varepsilon}(p||q) = +\infty$ .*

In Appendix A.3, we prove that the two definitions are indeed equivalent and give useful properties of the smooth Rényi divergence.

### 2.1 Rejection Sampling

Given two close enough densities  $p_t$  and  $p_s$ , either both continuous or both discrete, rejection sampling is a way to generate samples from  $p_t$  given access to samples from  $p_s$ , as explained for instance in [Dev86]. It was used mainly to generate samples from complex distributions that were “close” to easier-to-sample distributions. However, in cryptography and particularly in the line of

works started with [Lyu09], it found a peculiar use that diverged from its primary use. Given a family of densities  $(p_s^{(v)})$ , rejection sampling can be used to hide the parameter  $v$  given a density  $p_t$  that is close to every density in this family. It was later observed in [Lyu12] that an “imperfect” rejection procedure is sufficient for this use and leads to smaller parameters, notably standard deviation of  $p_s$ .

In the case of Lyubashevsky’s signature scheme [Lyu09,Lyu12], a signature is a pair of vectors  $(\mathbf{y} + \mathbf{S}\mathbf{c}, \mathbf{c})$  where  $\mathbf{y} \leftarrow P_{\mathbf{y}}$  and  $\mathbf{c}$  would ideally be sampled from  $P_{\mathbf{c}}$ . Here  $P_{\mathbf{c}} : \mathcal{C} \rightarrow \mathbb{R}$  and  $P_{\mathbf{y}} : \mathbb{Z}^m \rightarrow \mathbb{R}$  are two discrete probability distributions, where  $\mathcal{C} \subset \mathbb{Z}^k, m, k \geq 1$ , and  $\mathbf{S} \in \mathbb{Z}^{m \times k}$  is fixed (it is the signing key). The joint distribution of this pair corresponds to the source distribution  $P_s^{(\mathbf{S}\mathbf{c})}$  above, which depends on the sensitive data  $\mathbf{S}\mathbf{c}$ . Rejection sampling is used to ensure that the output of the signing algorithm is of the form  $(\mathbf{z}, \mathbf{c})$  where  $\mathbf{z} \leftarrow P_{\mathbf{z}}$  and  $\mathbf{c} \leftarrow P_{\mathbf{c}}$  are statistically independent and  $P_{\mathbf{z}}$  is well-chosen. Their joint distribution corresponds to the target distribution  $P_t$  above. The case of BLISS [DDLL13] is identical, except that signatures are of the form  $(\mathbf{y} + (-1)^b \mathbf{S}\mathbf{c}, \mathbf{c})$ , where  $b \leftarrow U(\{0, 1\})$ .

We consider the following algorithms from Figure 1, which take some  $M \geq 1$  as a parameter. Algorithm  $\mathcal{A}^{\text{ideal}}$  corresponds to what we would like to have, whereas  $\mathcal{A}^{\text{real}}$  is the algorithm corresponding to the real distribution. We are typically interested in calling these algorithms until they output something, which is what  $\mathcal{B}_{\infty}^{\text{real}}$  and  $\mathcal{B}_{\infty}^{\text{ideal}}$  do. It remains to understand when the outputs of these algorithms are statistically close. For completeness, we prove the following lemma in Appendix B.1.

<p>Algorithm <math>\mathcal{A}^{\text{real}}</math>:</p> <ol style="list-style-type: none"> <li>1: <math>x \leftarrow p_s</math></li> <li>2: with probability <math>\min\left(\frac{p_t(x)}{M \cdot p_s(x)}, 1\right)</math>, return <math>x</math></li> <li>3: return <math>\perp</math></li> </ol>	<p>Algorithm <math>\mathcal{A}^{\text{ideal}}</math>:</p> <ol style="list-style-type: none"> <li>1: <math>x \leftarrow p_t</math></li> <li>2: with probability <math>\frac{1}{M}</math>, return <math>x</math></li> <li>3: return <math>\perp</math></li> </ol>
<p>Algorithm <math>\mathcal{B}_{\infty}^{\text{real}}</math>:</p> <ol style="list-style-type: none"> <li>1: <math>z \leftarrow \perp</math></li> <li>2: <b>while</b> <math>z = \perp</math> <b>do</b></li> <li>3:     <math>z \leftarrow \mathcal{A}^{\text{real}}</math></li> <li>4: <b>end while</b></li> <li>5: return <math>z</math></li> </ol>	<p>Algorithm <math>\mathcal{B}_{\infty}^{\text{ideal}}</math>:</p> <ol style="list-style-type: none"> <li>1: <math>z \leftarrow \perp</math></li> <li>2: <b>while</b> <math>z = \perp</math> <b>do</b></li> <li>3:     <math>z \leftarrow \mathcal{A}^{\text{ideal}}</math></li> <li>4: <b>end while</b></li> <li>5: return <math>z</math></li> </ol>

**Fig. 1.** Rejection sampling algorithms.

**Lemma 2.2 (Adapted from [Lyu12, Lemma 4.7]).** *Assume that  $M \geq 1$  and  $\varepsilon \in [0, 1/2]$  are such that*

$$\Pr_{z \leftarrow p_t} (p_t(z) \leq M \cdot p_s(z)) \geq 1 - \varepsilon,$$

which can be rewritten in terms of smooth Rényi divergence as  $R_\infty^\varepsilon(p_t \| p_s) \leq M$ . Then the probability  $\mathcal{A}^{\text{real}}(\perp)$  that  $\mathcal{A}^{\text{real}}$  aborts is such that

$$\frac{M-1}{M} \leq \mathcal{A}^{\text{real}}(\perp) \leq \frac{M-1+\varepsilon}{M}.$$

Moreover, we have

$$\Delta(\mathcal{A}^{\text{real}}, \mathcal{A}^{\text{ideal}}) \leq \varepsilon/M \quad \text{and} \quad \Delta(\mathcal{B}_\infty^{\text{real}}, \mathcal{B}_\infty^{\text{ideal}}) \leq \varepsilon.$$

## 2.2 Lyubashevsky's signature scheme

All the following parameters are functions of a security parameter  $\lambda$ . We let  $k, m, n \geq 1$  and  $q \geq 2$  specify matrix spaces over  $\mathbb{Z}_q$ , with  $m > n$ . The distribution  $P_{\mathbf{S}}$  over  $\mathbb{Z}^{m \times k}$  is for signing keys and has support  $\mathcal{S} = \text{Supp}(P_{\mathbf{S}})$ . Let  $\mathcal{M}$  be the message space. Let  $\mathcal{C} \subset \mathbb{Z}^k$  finite and  $H : \mathbb{Z}_q^n \times \mathcal{M} \rightarrow \mathcal{C}$  a hash function, which is modeled as a random oracle in the signature scheme analysis. The parameter  $\gamma > 0$  is used in the verification algorithm to quantify the smallness of vectors corresponding to valid signatures. To obtain a  $2^\lambda$  security against known attacks, one typically sets  $m, n, k = \Omega(\lambda)$  and  $\gamma, q = \text{poly}(\lambda)$ .

Let  $\varepsilon \geq 0$  and  $M \geq 1$  be parameters related to rejection sampling, for a source distribution  $Q$  and a target distribution  $P$  over  $\mathbb{Z}^m$ . Most works directly instantiate these distributions. For example, uniform distributions in well-chosen hypercubes are used in [Lyu09] and  $P = Q$  Gaussian are used in [Lyu12]. We assume that the support of  $Q$  is contained in  $(-q/2, q/2]^m$ .

We consider the scheme presented in Figure 2, borrowed from [Lyu12] with the aforementioned rejection sampling generalization. For simplicity, we assume that the verification key  $\mathbf{A} \in \mathbb{Z}_q^{n \times m}$  is in Hermite normal form, i.e., we have  $\mathbf{A} = (\mathbf{I}_n | \mathbf{B})$  for some matrix  $\mathbf{B}$  and with  $\mathbf{I}_n \in \mathbb{Z}_q^{n \times n}$  denoting the identity matrix. Up to mild conditions on  $k, n, m, q$ , this is without loss of generality. Runtime and correctness follow from the two lemmas below.

**Lemma 2.3 (Sign Runtime).** *Let  $\varepsilon \geq 0$ ,  $M \geq 1$  and  $B = \lceil \lambda / \log \frac{M}{M-1+\varepsilon} \rceil$ . Assume that  $P$  and  $Q$  satisfy  $\max_{(\mathbf{s}, \mathbf{c}) \in \mathcal{S} \times \mathcal{C}} R_\infty^\varepsilon(P \| Q + \mathbf{s}\mathbf{c}) \leq M$ . Let  $(\mathbf{y}_0^\top | \mathbf{y}_1^\top)^\top \leftarrow Q$ , where  $\mathbf{y}_0$  takes values in  $\mathbb{Z}^n$ . In the ROM, the number of loop iterations  $i^*$  of a Sign execution satisfies*

$$\forall i : \Pr(i^* \geq i) \leq \left(1 - \frac{1-\varepsilon}{M}\right)^i + B^2 \cdot 2^{-H_\infty(\mathbf{y}_0 | \mathbf{y}_1)_Q} + 2^{-\lambda}.$$

*Note that when  $M \leq \text{poly}(\lambda)$ ,  $\varepsilon \leq 1 - 1/\text{poly}(\lambda)$  and  $2^{-H_\infty(\mathbf{y}_0 | \mathbf{y}_1)_Q} \leq \text{negl}(\lambda)$ , we have that  $B^2 \cdot 2^{-H_\infty(\mathbf{y}_0 | \mathbf{y}_1)_Q} + 2^{-\lambda} \leq \text{negl}(\lambda)$ .*

**Lemma 2.4 (Correctness).** *Let  $\varepsilon \geq 0$  and  $M \geq 1$ . Let  $P$  and  $Q$  satisfy  $\max_{(\mathbf{s}, \mathbf{c}) \in \mathcal{S} \times \mathcal{C}} R_\infty^\varepsilon(P \| Q + \mathbf{s}\mathbf{c}) \leq M$ . Let  $(\mathbf{y}_0^\top | \mathbf{y}_1^\top)^\top \leftarrow Q$ , where  $\mathbf{y}_0$  takes value in  $\mathbb{Z}^n$ . Further assume that  $2^{-H_\infty(\mathbf{y}_0 | \mathbf{y}_1)_Q} \leq \text{negl}(\lambda)$ ,  $\varepsilon \leq \text{negl}(\lambda)$  and the probability that Sign terminates is  $\geq 1 - \text{negl}(\lambda)$ . Then, in the ROM, the scheme is correct if  $\gamma \geq \gamma_P$  with  $\gamma_P$  such that  $\Pr_{\mathbf{z} \leftarrow P}(\|\mathbf{z}\| \geq \gamma_P) \leq \text{negl}(\lambda)$ .*

```

KeyGen( $1^\lambda$ ) :
1:  $\mathbf{B} \leftarrow \mathbb{Z}_q^{n \times (m-n)}$  and  $\mathbf{S} \leftarrow P_{\mathbf{S}}$ 
2:  $\mathbf{A} \leftarrow (\mathbf{I}_n | \mathbf{B})$ 
3:  $\mathbf{T} \leftarrow \mathbf{AS}$ 
4: return  $\text{vk} = (\mathbf{A}, \mathbf{T})$  and  $\text{sk} = (\mathbf{A}, \mathbf{S})$ 

Sign( $\mu, \mathbf{A}, \mathbf{S}$ ) :
1:  $\mathbf{y} \leftarrow Q$ 
2:  $\mathbf{c} \leftarrow H(\mathbf{Ay}, \mu)$ 
3:  $\mathbf{z} \leftarrow \mathbf{y} + \mathbf{Sc}$ 
4:  $u \leftarrow U([0, 1])$ 
5: if  $u \leq \min\left(\frac{P(\mathbf{z})}{M \cdot Q(\mathbf{y})}, 1\right)$  then
6:   return  $(\mathbf{z}, \mathbf{c})$ 
7: else
8:   go to Step 1
9: end if

Verify( $\mu, \mathbf{z}, \mathbf{c}, \mathbf{A}, \mathbf{T} = \mathbf{AS}$ ) :
1: if  $\|\mathbf{z}\| \leq \gamma$  and  $\mathbf{c} = H(\mathbf{Az} - \mathbf{Tc}, \mu)$ 
   then
2:   return 1
3: else
4:   return 0
5: end if

```

**Fig. 2.** Lyubashevsky’s signature scheme.

We only highlight components of typical security proofs that are relevant to our work, and refer to prior works for more details [Lyu09,Lyu12,AFLT16]. The security proofs of Lyubashevsky’s signature scheme all proceed by sequences of games and argue that the adversary’s advantages in successive games differ by small amounts and that no efficient adversary can solve the last game with a significant advantage.

An early step in the sequence of games is to replace the calls to  $H$  at Step 2 of the Sign algorithm by truly uniform and independent samples  $\mathbf{c} \leftarrow U(\mathcal{C})$ . To ensure that the adversary cannot notice the difference in the ROM, this requires that a given input  $(\mathbf{Ay}, \mu)$  to  $H$  cannot occur twice. This is obtained by having the conditional min-entropy  $H_\infty(\mathbf{y}_0 | \mathbf{y}_1)_Q$  satisfy:

$$H_\infty(\mathbf{y}_0 | \mathbf{y}_1)_Q = \Omega(\lambda).$$

An important other game hop consists in making Steps 1 to 6 of the Sign algorithm signing-key independent. Concretely, this means arguing that the distributions of the pair  $(\mathbf{z}, \mathbf{c})$  in the experiments from Figure 3 are statistically close, by using Lemma 2.2. (Note that this also requires programming  $H$  consistently with all appearing  $\mathbf{c}$ ’s.)

To complete the security proof, Lyubashevsky [Lyu12] reduces the SIS problem to the sEU-CMA security of a signing-key independent simulation of the Sign algorithm, by relying on the forking lemma. At this stage of the security proof, rejection sampling does not play a role anymore. We only note that the SIS instance has parameters  $q, m, n$  and  $\beta = 2(\gamma + \gamma')$ , with  $\gamma$  as in the Verify algorithm and  $\gamma' = \max_{(\mathbf{s}, \mathbf{c}) \in \mathcal{S} \times \mathcal{C}} \|\mathbf{Sc}\|$ . Note that  $\gamma$  is always significantly larger than  $\gamma'$ . We stress that there is a tension in setting  $\gamma$ : it should be sufficiently high to provide correctness (see Lemma 2.4 above) and as small as possible to provide higher security and hence allow more compact instantiations.

1: $\mathbf{y} \leftarrow Q$	1: $\mathbf{c} \leftarrow U(\mathcal{C})$
2: $\mathbf{c} \leftarrow U(\mathcal{C})$	2: $\mathbf{z} \leftarrow P$
3: $\mathbf{z} \leftarrow \mathbf{y} + \mathbf{S}\mathbf{c}$	3: $u \leftarrow U([0, 1])$
4: $u \leftarrow U([0, 1])$	4: <b>if</b> $u \leq \frac{1}{M}$ <b>then</b>
5: <b>if</b> $u \leq \min\left(\frac{P(\mathbf{z})}{M \cdot Q(\mathbf{y})}, 1\right)$ <b>then</b>	5:     return $(\mathbf{z}, \mathbf{c})$
6:     return $(\mathbf{z}, \mathbf{c})$	6: <b>else</b>
7: <b>else</b>	7:     return $(\perp, \perp)$
8:     return $(\perp, \perp)$	8: <b>end if</b>
9: <b>end if</b>	

**Fig. 3.** Simulating signatures.

### 3 Lower Bounds in the Case of Perfect Rejection Sampling

We start by studying *the case of perfect rejection sampling*, which corresponds to the setting of [Lyu09,DDLL13]. That is, we set  $\varepsilon = 0$  in the formalism of Section 2.2. We prove two lower bounds: (1) regarding signature size in both unimodal and bimodal settings (Sections 3.1 and 3.2), and (2) regarding the expected number of iterations of the rejection step (Section 3.3).

First, we analyze to which extent the expected norm of a distribution  $P$  can be decreased, under the constraint that we can reject to it using shifted samples from  $Q$ , where the Euclidean norm of the shift is bounded from above. This gives lower bounds on the norm of the signature vector  $\mathbf{z}$  in Lyubashevsky’s signature scheme, as recalled in Section 2.2. We start by studying the easier case of continuous distributions, and then provide a way to discretize the results.

Second, we prove that the classical rejection sampling strategy described above is optimal if one aims to minimize the expected number of iterations of the rejection step in the case of perfect rejection sampling from  $P$  to  $Q$ . Specifically, the expected number of iterations of any strategy is at least  $R_\infty(P\|Q)$ , which is reached by classical rejection sampling.

#### 3.1 Optimal Compactness in the Unimodal Setting

The main result of this subsection is the following.

**Theorem 3.1.** *Let  $m > 1, t > 0, V = \mathcal{B}_m(t)$  and  $M > 1$ . Let  $f, g : \mathbb{R}^m \rightarrow [0, 1]$  be two probability densities over  $\mathbb{R}^m$  such that  $\sup_{\mathbf{v} \in V} R_\infty(f\|g_{+\mathbf{v}}) \leq M$ . Then we have:*

$$\mathbb{E}_{\mathbf{x} \leftarrow f}(\|\mathbf{x}\|) \geq \frac{t}{M^{1/(m-1)} - 1}.$$

Note that we place ourselves in a setup where shifts belong to a hyperball. In the context of Lyubashevsky’s signature scheme, the shift is  $\mathbf{S}\mathbf{c}$ , where  $\mathbf{S}$  is the signing key and  $\mathbf{c}$  is the challenge (which is part of the signature). Given that  $\mathbf{S}$  is unknown, replacing the set of  $\mathbf{S}\mathbf{c}$ ’s by a hyperball seems to be a reasonable approach. Refining this approximation would lead to significant difficulties in the proof, with unlikely gains.

We now discuss the parameters  $M$  and  $m$ . As exhibited in Lemma 2.3, the variable  $M$  is related to the rejection probability. The smaller  $M$ , the faster we expect signing to be. To obtain a signing algorithm that terminates in polynomial time with overwhelming probability, we are interested in  $M \leq \text{poly}(\lambda)$ . Recall that we have  $m = \Omega(\lambda)$ . In this parameter regime, we have  $t/(M^{1/(m-1)} - 1) \approx t(m-1)/\log M$ .

The role of distribution  $g$  in Theorem 3.1 may seem puzzling, as it does not appear in the result. It acts as a control of the discrepancy of  $f$ : distribution  $f$  must be sufficiently wide to hide (in the Rényi divergence sense) a version of  $V$  that is blurred by  $g$ . This forces  $\mathbb{E}_{\mathbf{x} \leftarrow f}(\|\mathbf{x}\|)$  to be rather large. The proof proceeds in two steps. The first one consists in showing that there is no point favoring any direction and that we can restrict the study to isotropic distributions, i.e., distributions whose density is a function of the norm of the vector. The proof, which may be found in Appendix B, proceeds by averaging on shells. Theorem 3.1 is then obtained by integrating the local constraint  $\sup_{\mathbf{v} \in V} R_\infty(f\|g_{+\mathbf{v}}) \leq M$  over the whole support, with appropriate scaling.

**Lemma 3.2.** *Let  $m \geq 1, t > 0$  and  $V = \mathcal{B}_m(t)$ . Let  $f, g : \mathbb{R}^m \rightarrow [0, 1]$  be two probability densities over  $\mathbb{R}^m$  and define  $M = \sup_{\mathbf{v} \in V} R_\infty(f\|g_{+\mathbf{v}})$ . Then there exist two probability densities  $f^*, g^*$  that satisfy*

- $\sup_{\mathbf{v} \in V} R_\infty(f^*\|g_{+\mathbf{v}}^*) \leq M$ ,
- $\|\mathbf{x}\| = \|\mathbf{y}\| \implies g^*(\mathbf{x}) = g^*(\mathbf{y})$  and  $f^*(\mathbf{x}) = f^*(\mathbf{y})$ ,
- $\mathbb{E}_{\mathbf{z} \leftarrow f}(\|\mathbf{z}\|) = \mathbb{E}_{\mathbf{z} \leftarrow f^*}(\|\mathbf{z}\|)$ .

*Proof (Theorem 3.1).* Thanks to Lemma 3.2, we can, without loss of generality, assume that both  $f$  and  $g$  are isotropic. For  $k \geq 0$ , we define  $\mu_k = \int_0^\infty r^k f(r) dr$ , which is the  $k$ -th order moment of  $f$ . In particular, we have  $\mu_{m-1} = 1/S_m$  and  $\mu_m = \mathbb{E}_{\mathbf{x} \leftarrow f}(\|\mathbf{x}\|)/S_m$ . Indeed, using a hyperspherical variable change, we see that, for any  $\beta \in \{0, 1\}$ :

$$\begin{aligned} \mathbb{E}_{\mathbf{x} \leftarrow f}(\|\mathbf{x}\|^\beta) &= \int_{\mathbb{R}^m} \|\mathbf{x}\|^\beta f(\mathbf{x}) d\mathbf{x} \\ &= \int_0^\infty \rho^{m-1+\beta} f(\rho) \int_{[0, \pi]^{m-2} \times [0, 2\pi]} D(\vec{\theta}) d\vec{\theta} d\rho \\ &= S_m \cdot \mu_{m-1+\beta}. \end{aligned}$$

The above implies that  $\mathbb{E}_{\mathbf{x} \leftarrow f}(\|\mathbf{x}\|) = \mu_m/\mu_{m-1}$ .

For any  $x \geq 0$  and  $u \in [-t, t]$ , it holds that  $f(x) \leq M \cdot g(|x-u|)$ . In particular, for  $x \geq t$ , we have  $f(x-t) \leq M \cdot g(x)$ . Let us multiply both sides by  $x^{m-1}$  and integrate over  $[t, +\infty)$ . With a change of variable on the left-hand side, this gives

$$\begin{aligned} \int_0^\infty (x+t)^{m-1} f(x) dx &\leq M \cdot \int_t^\infty x^{m-1} g(x) dx \\ &\leq M \cdot \int_0^\infty x^{m-1} g(x) dx \\ &= M \cdot \int_0^\infty x^{m-1} f(x) dx, \end{aligned}$$

by recognizing that the right-hand side is  $M \cdot \mu_{m-1}$  (which is the same for  $f$  and  $g$ ). Grouping everything on the same side, we have

$$0 \leq \int_0^\infty (Mx^{m-1} - (x+t)^{m-1}) f(x) dx. \quad (1)$$

Let  $C = t/(M^{1/(m-1)} - 1)$ . For  $m > 2$ , we rewrite the integrand as

$$\begin{aligned} Mx^{m-1} - (x+t)^{m-1} &= \left(M^{\frac{1}{m-1}}x - (x+t)\right) \cdot \sum_{k=0}^{m-2} \left(xM^{\frac{1}{m-1}}\right)^k (x+t)^{m-2-k} \\ &= \left(M^{\frac{1}{m-1}} - 1\right) (x-C) \cdot \sum_{k=0}^{m-2} \left(xM^{\frac{1}{m-1}}\right)^k (x+t)^{m-2-k}. \end{aligned}$$

For  $m = 2$ , the above holds by replacing the sum by 1. Now, note that the inequality  $xM^{1/(m-1)} \geq x+t$  holds if and only if  $x \geq C$ . Hence the following upper bound holds for any  $x \geq 0$ , if  $m > 2$ :

$$(x-C) \cdot \sum_{k=0}^{m-2} (xM^{\frac{1}{m-1}})^k (x+t)^{m-2-k} \leq (x-C)(m-1)M^{\frac{m-2}{m-1}}x^{m-2}.$$

When  $m > 2$ , we can divide by  $(M^{1/(m-1)} - 1)M^{(m-2)/(m-1)}(m-1) > 0$  in Equation (1), and obtain:

$$C \cdot \int_0^\infty x^{m-2} f(x) dx \leq \int_0^\infty x^{m-1} f(x) dx.$$

Note that it also holds for  $m = 2$ . This can be rewritten as  $\mu_{m-1}/\mu_{m-2} \geq C$ .

Now, observe that  $\mu_m \cdot \mu_{m-2} \geq (\mu_{m-1})^2$ . Indeed, the Cauchy-Schwarz inequality states that for any real random variables  $X, Y$ , it holds that  $|\mathbb{E}(XY)|^2 \leq \mathbb{E}(X^2)\mathbb{E}(Y^2)$ . We instantiate it with the (non-independent) random variables  $X = \|\mathbf{x}\|^{m/2}$  and  $Y = \|\mathbf{x}\|^{(m-2)/2}$ , where  $\mathbf{x} \leftarrow f$ . Then  $XY = \|\mathbf{x}\|^{\frac{m}{2} + \frac{m-2}{2}} = \|\mathbf{x}\|^{m-1}$ . To conclude, note  $\mu_m \cdot \mu_{m-2} \geq (\mu_{m-1})^2$  implies that  $\mu_m/\mu_{m-1} \geq \mu_{m-1}/\mu_{m-2} \geq C$ . This completes the proof.  $\square$

For the discrete case, we observe that given a discrete distribution  $P$ , letting  $f : \mathbf{x} \mapsto P(\lceil \mathbf{x} \rceil)$  be a probability density over  $\mathbb{R}^m$ , we have, by the triangle inequality

$$\mathbb{E}_{\mathbf{x} \leftarrow f}(\|\mathbf{x}\|) \leq \mathbb{E}_{\mathbf{x} \leftarrow P}(\|\mathbf{x}\|) + \frac{\sqrt{m}}{2}.$$

Theorem 3.1 can then be adapted to the discrete case, up to subtracting  $\sqrt{m}/2$  from the lower bound. In all setups considered in this work, this term is significantly smaller than  $t/(M^{1/(m-1)} - 1)$ .

**Corollary 3.3.** *Let  $m > 1$ ,  $t > 0$ ,  $V = \mathcal{B}_m(t) \cap \mathbb{Z}^m$  and  $M > 1$ . Let  $P$  and  $Q$  be two discrete probability distributions over  $\mathbb{Z}^m$  such that  $\sup_{\mathbf{v} \in V} R_\infty(P \| Q_{+\mathbf{v}}) \leq M$ . Then we have:*

$$\mathbb{E}_{\mathbf{x} \leftarrow P}(\|\mathbf{x}\|) \geq \frac{t}{M^{1/(m-1)} - 1} - \frac{\sqrt{m}}{2}.$$

### 3.2 Optimal Compactness in the Bimodal Setting

We obtain the following result in the bimodal setting. As the proof is similar to the one of Theorem 3.1, it is postponed to Appendix B.3.

**Theorem 3.4.** *Let  $m \geq 3, t > 0, V = \mathcal{B}_m(t)$  and  $M > 1$ . Let  $f, g : \mathbb{R}^m \rightarrow [0, 1]$  be two probability densities over  $\mathbb{R}^m$  such that  $\sup_{\mathbf{v} \in V} R_\infty(f \| g_{\pm \mathbf{v}}) \leq M$ , where  $g_{\pm \mathbf{v}}$  is the density  $\mathbf{x} \mapsto \frac{1}{2}(g(\mathbf{x} - \mathbf{v}) + g(\mathbf{x} + \mathbf{v}))$ . Then the following holds:*

$$\mathbb{E}_{\mathbf{x} \leftarrow f}(\|\mathbf{x}\|) \geq \frac{t}{\sqrt{M^{\frac{2}{m-2}} - 1}}.$$

For  $M \leq \text{poly}(\lambda)$  and  $m = \Omega(\lambda)$  as in the discussion following Theorem 3.1, we have  $t/(M^{2/(m-2)} - 1)^{1/2} \approx t\sqrt{(m-2)/(2 \log M)}$ . Similarly to the unimodal case, the lower bound can be adapted to integer distributions with limited loss (for all setups considered in this work).

**Corollary 3.5.** *Let  $m \geq 3, t > 0, V = \mathcal{B}_m(t) \cap \mathbb{Z}^m$  and  $M > 1$ . Let  $P$  and  $Q$  be two discrete probability distributions over  $\mathbb{Z}^m$  such that  $\sup_{\mathbf{v} \in V} R_\infty(P \| Q_{\pm \mathbf{v}}) \leq M$ , where  $Q_{\pm \mathbf{v}}$  is as in Theorem 3.4. Then the following holds:*

$$\mathbb{E}_{\mathbf{x} \leftarrow P}(\|\mathbf{x}\|) \geq \frac{t}{\sqrt{M^{\frac{2}{m-2}} - 1}} - \frac{\sqrt{m}}{2}.$$

### 3.3 Optimality of the Expected Number of Iterations

We now analyze to which extent the expected number of iterations of the rejection step could be reduced in the case of exact rejection sampling from  $P$  to  $Q$ , and prove the classical strategy to be optimal. This question arises from the variety of rejection sampling techniques that have been studied in other fields.

There exist multiple variants of rejection sampling. For instance, a procedure described in [HJMR07] and recalled in Appendix D takes a greedy approach to rejection sampling and differs from the one we presented up until now. We are in the setting where we have access to a sampler from distribution  $Q$ . These samples are denoted by  $(X_i)_{i \geq 1}$  with  $X_i \in \mathcal{X}$  for some set  $\mathcal{X}$  and we are required to output a sample from the distribution  $P$  over  $\mathcal{X}$ . Any design of procedure is allowed, as long as the output is one of the observed samples  $X_i$ . Let  $i^*$  be the random variable denoting the number of samples observed by an algorithm and we wish to determine how small  $\mathbb{E}(i^*)$  can be. We note that the work of [HJMR07], establishes that there exists a rejection sampling algorithm achieving  $\mathbb{E}(\log i^*) = \log R_1(P \| Q)$  up to lower order terms in  $R_1(P \| Q)$ , and that this is optimal. Here, we show that the minimum value for  $\mathbb{E}(i^*)$  is  $R_\infty(P \| Q)$ .

We model a rejection sampling algorithm by a family of randomized functions  $A_i : \mathcal{X}^i \rightarrow \{1, \dots, i\} \cup \{r\}$ . At step  $i$ , it sees the new sample  $X_i$  and based on  $X_1, \dots, X_i$  it computes  $A_i(X_1, \dots, X_i)$ . If it is equal to  $r$ , the algorithm asks for one more sample and otherwise if  $A_i(X_1, \dots, X_i) \in \{1, \dots, i\}$ , the algorithm terminates and outputs the sample  $X_{A_i(X_1, \dots, X_i)}$ . Note that the running time

of the algorithm is defined by  $i^* = \inf\{i \geq 1 : A_i(X_1, \dots, X_i) \neq r\}$ . We only consider algorithms for which  $i^* < \infty$  almost surely. We define the random variable  $J = A_{i^*}(X_1, \dots, X_{i^*}) \in \mathbb{N}_+$ , note that  $J \leq i^*$  and the output of the algorithm is  $X_J$  (i.e., the output sample may not be the last one that was generated).

**Theorem 3.6.** *Let  $P, Q$  be two discrete probability distributions. Any rejection sampling algorithm  $(A_i)_{i \geq 1}$  sampling from  $P$  satisfies  $\mathbb{E}(i^*) \geq R_\infty(P\|Q)$ .*

*Proof.* We have by assumption for any  $x \in \mathcal{X}$ ,

$$P(x) = \Pr[X_J = x] = \sum_{j=1}^{\infty} \Pr[J = j, X_j = x] \leq \sum_{j=1}^{\infty} \Pr[i^* \geq j, X_j = x],$$

where we used the fact that the event  $[J = j]$  is contained in  $[i^* \geq j]$ . Now, observe that the event  $[i^* < j]$  only depends on  $X_1, \dots, X_{j-1}$  and as such it is independent of the event  $[X_j = x]$ . This implies that  $[i^* \geq j]$  is independent of  $[X_j = x]$ . As a result, we have

$$P(x) \leq \sum_{j=1}^{\infty} \Pr[i^* \geq j] \Pr[X_j = x] = \mathbb{E}(i^*)Q(x),$$

which proves the desired result.

In the context of Lyubashevsky's signature schemes with source distribution  $Q'$ , target distribution  $P'$ , challenge set  $\mathcal{C}$  and signing key  $\mathbf{S}$ , we would have  $P = P' \otimes U(\mathcal{C})$  and  $Q$  would be the distribution of the pair  $(\mathbf{z}, \mathbf{c})$  obtained by sampling  $\mathbf{y}$  from  $Q'$ ,  $\mathbf{c}$  from  $U(\mathcal{C})$  and defining  $\mathbf{z} = \mathbf{y} + \mathbf{S}\mathbf{c}$ .

The above proof can be adapted in the setting where  $P$  and  $Q$  are continuous distributions by considering a sequence of balls converging to  $\{x\}$  instead of  $x$ .

## 4 Improved Analysis via the Rényi Divergence

For the rest of the paper, we flip our focus and prove positive results (upper bounds). In this section, we propose an improved analysis of Lyubashevsky's signatures that relies on the Rényi divergence rather than the statistical distance, allowing larger sampling errors in the case of imperfect rejection sampling. Then, in Section 5 propose a new choice of distributions that (asymptotically) reaches our lower bounds. Finally, in Section 6, we propose a way to circumvent the lower bound for the expected number of iterations by providing an alternate strategy which allows to fix a-priori a maximal number of loop iterations.

Our lower bounds apply to perfect rejection sampling, but rejecting to an inaccurate approximation to the target distribution also allows to instantiate Lyubashevsky's signature, as done in [Lyu12] and already mentioned in Section 2.2 (when  $\varepsilon > 0$  in Lemma 2.2). In particular, imperfect rejection sampling is used when instantiating the signature scheme with Gaussian distributions.

In this section, we study the case of imperfect rejection sampling and describe a way to improve the analysis of the digital signature from Section 2.2, by replacing the statistical distance (in Lemma 2.2) with the Rényi divergence to quantify the closeness between ideal and real rejection sampling algorithms. As already observed in prior works (see in particular the discussion in [BLR<sup>+</sup>18]), the Rényi divergence is well-suited for improving the analyses of digital signatures, as the security game is of a search type. While the analysis based on the statistical distance imposes  $\varepsilon = 2^{\Omega(\lambda)}$ , as it requires the statistical distance to be negligible, our analysis allows larger sampling errors as it only imposes  $\varepsilon \approx 1/q_{\text{sig}}$  where  $q_{\text{sig}}$  is the number of signing queries (which is  $\text{poly}(\lambda) \ll 2^{\Omega(\lambda)}$ ).

#### 4.1 Rényi Divergence Bounds for Imperfect Rejection Sampling

Let  $p_t$  and  $p_s$  be two probability densities, both continuous or both discrete. We consider algorithms  $\mathcal{A}^{\text{real}}$ ,  $\mathcal{A}^{\text{ideal}}$ ,  $\mathcal{B}^{\text{real}}$  and  $\mathcal{B}^{\text{ideal}}$  from Figure 1.

**Lemma 4.1.** *Assume that  $M > 1$  and  $\varepsilon < 1$  are such that  $R_\infty^\varepsilon(p_t \| p_s) \leq M$ . Then for any  $a \in (1, +\infty)$  we have:*

$$R_a(\mathcal{A}^{\text{real}} \| \mathcal{A}^{\text{ideal}}) \leq \left( \frac{1}{M} + \frac{M-1+\varepsilon}{M} \cdot \left( 1 + \frac{\varepsilon}{M-1} \right)^{a-1} \right)^{\frac{1}{a-1}},$$

$$R_a(\mathcal{B}_\infty^{\text{real}} \| \mathcal{B}_\infty^{\text{ideal}}) \leq \frac{1}{(1-\varepsilon)^{a/(a-1)}}.$$

Moreover, for  $a = \infty$ , we have:

$$R_\infty(\mathcal{A}^{\text{real}} \| \mathcal{A}^{\text{ideal}}) \leq 1 + \frac{\varepsilon}{M-1} \quad \text{and} \quad R_\infty(\mathcal{B}_\infty^{\text{real}} \| \mathcal{B}_\infty^{\text{ideal}}) \leq \frac{1}{1-\varepsilon}.$$

Note that for  $\varepsilon = 0$ , we recover the distributional equalities  $\mathcal{A}^{\text{real}} = \mathcal{A}^{\text{ideal}}$  and  $\mathcal{B}_\infty^{\text{real}} = \mathcal{B}_\infty^{\text{ideal}}$  of Lemma 2.2. We are interested in the case  $\varepsilon > 0$ .

*Proof.* Let  $\mathcal{A}^{\text{real}}(\perp)$  and  $\mathcal{A}^{\text{ideal}}(\perp)$  denote the probabilities that  $\mathcal{A}^{\text{real}}$  or  $\mathcal{A}^{\text{ideal}}$  output nothing. We have, using results from Lemma 2.2:

$$\begin{aligned} R_a(\mathcal{A}^{\text{real}} \| \mathcal{A}^{\text{ideal}})^{a-1} &= \left[ \int_{\text{Supp}(p_s)} \frac{\left( p_s(x) \min\left( \frac{p_t(x)}{M \cdot p_s(x)}, 1 \right) \right)^a}{(p_t(x)/M)^{a-1}} dx \right] + \frac{(\mathcal{A}^{\text{real}}(\perp))^a}{(\mathcal{A}^{\text{ideal}}(\perp))^{a-1}} \\ &\leq \int_{\text{Supp}(p_s)} \frac{\left( p_s(x) \frac{p_t(x)}{M \cdot p_s(x)} \right)^a}{(p_t(x)/M)^{a-1}} dx + \frac{(1 - (1-\varepsilon)/M)^a}{(1 - 1/M)^{a-1}} \\ &= \int_{\text{Supp}(p_s)} \frac{p_t(x)}{M} dx + \frac{M-1+\varepsilon}{M} \cdot \left( \frac{M-1+\varepsilon}{M-1} \right)^{a-1} \\ &\leq \frac{1}{M} + \frac{M-1+\varepsilon}{M} \cdot \left( 1 + \frac{\varepsilon}{M-1} \right)^{a-1}. \end{aligned}$$

We move on to bounding the second divergence. For any  $x \in \text{Supp}(p_s)$ :

$$\mathcal{B}_\infty^{\text{real}}(x) = \frac{\mathcal{A}^{\text{real}}(x)}{1 - \mathcal{A}^{\text{real}}(\perp)}.$$

This also holds for  $\mathcal{B}_\infty^{\text{ideal}}$  with  $\mathcal{A}^{\text{ideal}}$  instead of  $\mathcal{A}^{\text{real}}$ . We obtain:

$$\begin{aligned} R_a(\mathcal{B}_\infty^{\text{real}} \parallel \mathcal{B}_\infty^{\text{ideal}})^{a-1} &= \int_{\text{Supp}(p_s)} \frac{1}{M^{a-1}} \cdot \frac{\left(p_s(x) \min\left(\frac{p_t(x)}{M \cdot p_s(x)}, 1\right)\right)^a}{(\mathcal{A}^{\text{real}}(\perp))^a (p_t(x)/M)^{a-1}} \\ &\leq \frac{M}{(1-\varepsilon)^a} \int_{\text{Supp}(p_s)} \frac{\left(p_s(x) \min\left(\frac{p_t(x)}{M \cdot p_s(x)}, 1\right)\right)^a}{(p_t(x)/M)^{a-1}}. \end{aligned}$$

This sum was already computed just above and is at most  $1/M$ .

The continuity of  $a \mapsto R_a(P_t \parallel P_s)$  at  $a = +\infty$  gives the last bounds.  $\square$

## 4.2 Improved Analysis of Lyubashevsky's Scheme

We now go back to the scheme described in Section 2.2 with imperfect rejection sampling, and show that the analysis above allows setting  $\varepsilon \approx 1/q_{\text{sig}}$  instead of  $\varepsilon = 2^{-\Omega(\lambda)}$ . Here  $q_{\text{sig}}$  refers to the number of signing queries that an adversary can make. As a signing query requires an interaction with the signer, it is typically considered to be a large polynomial in  $\lambda$ , which is much smaller than  $2^{\Omega(\lambda)}$ . As a result, this refined analysis puts less stress on the condition that  $P_s$  and  $P_t$  must satisfy and hence to reach smaller values for  $\mathbb{E}_{\mathbf{z} \leftarrow P_t}(\|\mathbf{x}\|)$ : this is beneficial to security and then allows for smaller parameter sets.

To achieve this improvement, we replace the statistical distance with the Rényi divergence in the scheme analysis, when simulating signature queries (see Figure 3). By Lemma 4.1 and the data processing inequality of Lemma A.4, replacing  $\mathcal{A}^{\text{real}}$  by  $\mathcal{A}^{\text{ideal}}$  once in the security proof (i.e., in one loop iteration of one signature query) leads to a multiplicative loss of a factor  $\leq 1 + \varepsilon/(M-1)$  in the adversary's advantage. Now, note that the probability that at least one among the  $q_{\text{sig}}$  sign queries requires more than  $B = (\lambda + \log q_{\text{sig}})/\log(M/(M-1+\varepsilon))$  loop iterations is exponentially small. Assuming this is not the case, we can bound the number of times  $\mathcal{A}^{\text{real}}$  is replaced by  $\mathcal{A}^{\text{ideal}}$  in the security proof by  $B \cdot q_{\text{sig}}$ . By the Rényi divergence multiplicativity property (see Lemma A.4), this induces a multiplicative loss of a factor  $\leq (1 + \varepsilon/(M-1))^{B \cdot q_{\text{sig}}}$  in the adversary's advantage.

## 5 Reaching the Lower Bounds with Hyperballs

In this section, we show that continuous uniform distributions in hyperballs reach the lower bounds in both the unimodal and bimodal perfect rejection sampling settings. We also consider the imperfect unimodal setting and find parameters that are asymptotically at least as good as the ones obtained for the Gaussian distribution (using our analysis described in Section 4). As continuous hyperball

uniform distributions are easier both to study and implement than their discrete counterpart, we consider the case of continuous distributions. Further, we show that a slight modification of Lyubashevsky's signature allows for the target and source distributions to be continuous.

We also compare this choice of distributions with the uniform distributions in hypercubes and with Gaussians, both asymptotically and with concrete parameters.

### 5.1 Uniform Distributions in Hyperballs

The first step is to compute the divergence in the three settings: unimodal, either perfect or imperfect rejection sampling and bimodal perfect rejection sampling. The first case can actually be seen as a particular case of the second one, and we summarize both in the following lemma. The function  $I$  appearing in the statement is defined in Appendix A.6 using the beta function, and comes into play when dealing with hyperspherical caps.

**Lemma 5.1 (Smooth Divergence).** *Let  $m \geq 1$  and  $\mathbf{v} \in \mathbb{R}^m$ . Let  $\varepsilon \in [0, 1/2]$  and  $\eta \geq 1$  be such that  $2\varepsilon = I_{1-1/\eta^2}(\frac{m+1}{2}, \frac{1}{2})$ . Let  $r, r' > 0$  such that  $r'^2 \geq r^2 + \|\mathbf{v}\|^2 + 2r\|\mathbf{v}\|/\eta$ . Then it holds that:*

$$R_\infty^\varepsilon\left(U(\mathcal{B}_m(r))\|U(\mathcal{B}_m(r', \mathbf{v}))\right) = \left(\frac{r'}{r}\right)^m.$$

Let  $M > 1$ . The above is  $\leq M$  if  $r \geq \|\mathbf{v}\| \cdot \frac{\frac{1}{\eta} + \sqrt{\frac{1}{\eta^2} + M^{2/m} - 1}}{M^{2/m} - 1}$  and  $r' = M^{1/m}r$ .

Note that when  $\varepsilon = 0$ , we have  $\eta = 1$ . In that case, we can set  $r = \|\mathbf{v}\|/(M^{1/m} - 1)$ , which almost matches the lower bound from Theorem 3.1. As seen in Appendix A.6, for  $\varepsilon = 2^{-c \cdot m}$  with a constant  $c > 0$ , we have that  $1/\eta^2$  tends to  $1 - 2^{-c}$  when  $m$  goes to infinity. For  $\varepsilon$  satisfying  $\varepsilon \geq 2^{-o(m)}$  and  $\varepsilon = o(1/m)$  with  $m$  going to infinity, we have that  $1/\eta^2 \sim -\log(\varepsilon)/m$ .

*Proof.* Assume that there exists some cut  $\mathcal{C}$  with  $\text{vol}(\mathcal{C})/V_m(r) \leq \varepsilon$  such that the divergence is defined, i.e., with  $\mathcal{B}_m(r) \setminus \mathcal{C} \subseteq \mathcal{B}_m(r', \mathbf{v})$ . Then the divergence is  $(r'/r)^m$ , as the ratio of densities is constant and equal to  $(r'/r)^m$  over  $\mathcal{B}_m(r) \setminus \mathcal{C}$ . To prove the first claim, it hence suffices to show that such a cut  $\mathcal{C}$  exists.

We introduce the cut  $\mathcal{C}_\eta := \{\mathbf{x} \in \mathcal{B}_m(r) \mid \langle \mathbf{x}, \mathbf{v} \rangle \geq -\|\mathbf{v}\|r/\eta\}$ . This is the intersection of a ball with an affine half-space, i.e., an  $m$ -dimensional hyperspherical cap. By Lemma A.13, its volume is  $\frac{V_m(r)}{2} \cdot I_{1-1/\eta^2}(\frac{m+1}{2}, \frac{1}{2})$ . The definition of  $\eta$  ensures that  $\text{vol}(\mathcal{C}_\eta)/V_m(r) = \varepsilon$ . We now check that  $\mathcal{B}_m(r) \setminus \mathcal{C}_\eta \subseteq \mathcal{B}_m(r', \mathbf{v})$ . Let  $\mathbf{x} \in \mathcal{B}_m(r) \setminus \mathcal{C}_\eta$ . We have

$$\|\mathbf{x} - \mathbf{v}\| \leq \sqrt{r^2 + \|\mathbf{v}\|^2 + 2r\|\mathbf{v}\|/\eta}.$$

By assumption, the latter is no larger than  $r'$ , implying that  $\mathbf{x} \in \mathcal{B}_m(r', \mathbf{v})$ . This completes the proof of the first claim.

If we combine the condition on  $r$  and  $r'$  and the equality  $r' = M^{1/m}r$ , we get

$$r^2 + \|\mathbf{v}\|^2 + 2\frac{r\|\mathbf{v}\|}{\eta} \leq M^{2/m}r^2,$$

which is a degree-2 inequality on  $r$ . Solving it completes the proof.  $\square$

**Lemma 5.2 (Divergence in the Bimodal Setting).** *Let  $m \geq 1$  and  $\mathbf{v} \in \mathbb{R}^m$ . Let  $r, r' > 0$  such that  $r'^2 \geq r^2 + \|\mathbf{v}\|^2$ . Let  $U(\mathcal{B}_m(r'), \pm\mathbf{v})$  denote the continuous probability distribution which samples  $b \leftarrow U(\{0, 1\})$  and returns  $\mathbf{z} \leftarrow U(\mathcal{B}_m(r'), (-1)^b\mathbf{v})$ . Then it holds that:*

$$R_\infty\left(U(\mathcal{B}_m(r))\|U(\mathcal{B}_m(r'), \pm\mathbf{v})\right) = (1 + \chi_{<r+\|\mathbf{v}\|}(r')) \cdot \left(\frac{r'}{r}\right)^m,$$

where  $\chi_{<r+\|\mathbf{v}\|}$  denotes the indicator function of reals smaller than  $r + \|\mathbf{v}\|$ . Let  $M > 1$ . The above is  $\leq M$  if  $r \geq \|\mathbf{v}\|/\sqrt{(M/2)^{2/m} - 1}$  and  $r' = (M/2)^{1/m}r$ .

Note that the choice of  $r$  almost matches the lower bound from Theorem 3.4.

*Proof.* The support of  $U(\mathcal{B}_m(r'), \pm\mathbf{v})$  is exactly  $\mathcal{B}_m(r', \mathbf{v}) \cup \mathcal{B}_m(r', -\mathbf{v})$  and its density is  $\mathbf{z} \mapsto (\chi_{\mathcal{B}_m(r', \mathbf{v})}(\mathbf{z}) + \chi_{\mathcal{B}_m(r', -\mathbf{v})}(\mathbf{z})) / (2V_m(r'))$ . The divergence is finite when  $\mathcal{B}_m(r) \subseteq \mathcal{B}_m(r', \mathbf{v}) \cup \mathcal{B}_m(r', -\mathbf{v})$ . This is the case if any  $\mathbf{x}$  with  $\|\mathbf{x}\| \leq r$  satisfies  $\|\mathbf{x} - \mathbf{v}\| \leq r'$  or  $\|\mathbf{x} + \mathbf{v}\| \leq r'$ . Let us assume, w.l.o.g., that  $\|\mathbf{x} - \mathbf{v}\| \leq \|\mathbf{x} + \mathbf{v}\|$ . Then we write

$$\|\mathbf{x} - \mathbf{v}\| = \sqrt{\|\mathbf{x}\|^2 + \|\mathbf{v}\|^2 - 2\langle \mathbf{x}, \mathbf{v} \rangle} \leq \sqrt{\|\mathbf{x}\|^2 + \|\mathbf{v}\|^2}.$$

Thanks to the assumption on  $r$  and  $r'$ , we conclude that the divergence is finite.

Now, the ratio of the densities only takes three values. If  $\mathbf{x} \notin \mathcal{B}_m(r)$  then the ratio is 0. If  $\mathbf{x} \in \mathcal{B}_m(r) \cap \mathcal{B}_m(r', \mathbf{v}) \cap \mathcal{B}_m(r', -\mathbf{v})$  then the ratio is  $(r'/r)^m$ . Finally, if  $\mathbf{x}$  belongs to  $\mathcal{B}_m(r) \cap \mathcal{B}_m(r', \mathbf{v})$  but not to  $\mathcal{B}_m(r', -\mathbf{v})$ , then the ratio is  $2(r'/r)^m$ . This last case only occurs if  $\mathcal{B}_m(r) \not\subseteq \mathcal{B}_m(r', -\mathbf{v})$ . This is the case only if  $r' < r + \|\mathbf{v}\|$ . This completes the proof of the first claim.

For the second claim, note that the assumption on  $r$  and  $r'$  is satisfied, and that the divergence bound is indeed  $\leq M$ .  $\square$

In this bimodal case, the rejection test is as follows. One computes the norms of both  $\mathbf{z}$  and  $\mathbf{z} - 2(-1)^b\mathbf{v}$ , where  $\mathbf{z} = \mathbf{y} + (-1)^b\mathbf{v}$ . If only the first one is  $\leq r$ , then the sample is accepted. If both are  $\leq r$ , then it is accepted and rejected with probability 1/2.

Finally, in order to use the uniform distribution in a hyperball, we verify that there is sufficient min-entropy in the first  $n$  coordinates given the remaining  $m-n$  coordinates. The proof of the following lemma can be found in Appendix B.4.

**Lemma 5.3.** *Let  $m \geq 6, n \geq 1$  and  $r \geq 2\sqrt{m}$ . Let  $\mathbf{x} = (\mathbf{x}_0^\top | \mathbf{x}_1^\top)^\top$  be a random variable over  $\mathbb{R}^m$  whose distribution is  $U(\mathcal{B}_m(r))$ , where  $\mathbf{x}_0$  has dimension  $n$ . It holds that*

$$H_\infty(\lceil \mathbf{x}_0 \rceil | \lceil \mathbf{x}_1 \rceil)_{U(\mathcal{B}_m(r))} \geq \left(\log_2 \frac{1}{0.85}\right) \cdot n.$$

## 5.2 Lyubashevsky’s Signature with Continuous Distributions

We consider continuous distributions over hyperballs, which are not directly compatible with Lyubashevsky’s signature scheme, as recalled in Section 2. Switching to uniform distributions over the integer points inside hyperballs leads to several difficulties: sampling from such a distribution seems delicate, in particular if the radius of the ball is moderate. Similarly, adapting Lemmas 5.1 and 5.2 seems difficult. Rather, we argue that it is possible to extend Lyubashevsky’s signature scheme to the case of continuous distributions, and that this comes with very limited complications (in the case of Gaussians, it could be simpler to use continuous Gaussians with this modified scheme, than using discrete Gaussians with the original scheme).

In order to adapt Lyubashevsky’s signature scheme to continuous distributions, a rounding step is added after acceptance of a sample, as well as during hashing. Concretely, the changes compared to the construction described in Figure 2 are as follows: (i)  $\mathbf{y}$  is now sampled from a continuous distribution with density  $g$ , (ii)  $\mathbf{c}$  is now computed as  $H(\mathbf{A}\lceil\mathbf{y}\rceil, \mu)$ , (iii) with  $\mathbf{z}$  still being defined as  $\mathbf{y} + \mathbf{S}\mathbf{c}$ , if the test passes, and the returned signature is now  $(\lceil\mathbf{z}\rceil, \mathbf{c})$ . This adaptation is discussed in more details in Appendix C.3. We note that this leads to the requirement that the min-entropy of  $\lceil\mathbf{x}_0\rceil\lceil\mathbf{x}_1\rceil$  is large, where  $\mathbf{x} = (\mathbf{x}_0^\top | \mathbf{x}_1^\top)^\top$  is a random variable over  $\mathbb{R}^m$  whose distribution is  $g$  and  $\mathbf{x}_0$  has dimension  $n$ . In the case of the uniform distribution in a hyperball, this is provided by Lemma 5.3.

We further remark that this applies to the analysis relying on the statistical distance as well as our improved analysis which relies on the Rényi divergence. Also, we note that the modified scheme involves computations over real numbers. These can be securely replaced by finite precision computations, using standard techniques such as described in [Pre17].

## 5.3 Comparison with other Distributions

Let  $t = \max_{\mathbf{S}, \mathbf{c}} \|\mathbf{S}\mathbf{c}\|$ . In Table 2, we summarize the expected norm of signatures (up to a constant factor) for diverse distributions  $P$  and  $Q$ , and for a target expected number of iterations  $M$ . We consider three specific pairs of distributions, two of them being previously considered distributions (Gaussians and uniforms in hypercubes), and the last one being uniform distributions in hyperballs, introduced above. We also consider three different scenarios:

- unimodal distributions and perfect rejection sampling, corresponding to the column  $\varepsilon = 0$ ;
- unimodal distributions and imperfect rejection sampling – we use approximations specific to the choice of  $\varepsilon \geq 2^{-o(m)}$  and  $\varepsilon = o(1/m)$ ;
- bimodal source distribution and perfect rejection sampling, corresponding to column “Bimodal”.

Note that the second scenario relies on our improved analysis relying on the Rényi divergence for the imperfect case (see Section 4). This parameter range for  $\varepsilon$  is not appropriate when using the analysis relying on the statistical distance.

In the last column, we also emphasize if the test that decides to keep or reject a sample is simple or not. For hyperballs, it simply consists in comparing the norm of the sample with the radius of the target hyperball in the unimodal case. For the bimodal case, one needs to compute two norms, and if necessary, to flip a coin (see discussion after Lemma 5.2).

The entries in the table are approximations for  $m \rightarrow \infty$ ,  $t = \omega(1)$  and  $M = 2^{o(m)}$ , and for a given choice of  $P$ , we optimize the parametrization of  $Q$  (e.g., the radius in case of a hyperball) to minimize the signature norm.

Choices for $P$ and $Q$	$\varepsilon = 0$	$\varepsilon \geq 2^{-o(m)}$ and $\varepsilon = o(1/m)$	Bimodal	Rejection Test
Hypercubes	$\frac{tm^{3/2}}{\log M}$	$\frac{tm^{3/2}}{\log M}$	$\frac{tm^{3/2}}{\log M}$	Simple
Gaussians	$\infty$	$\frac{t\sqrt{m}\sqrt{\log \frac{1}{\varepsilon} + \log M}}{\log M}$	$\frac{t\sqrt{m}}{\sqrt{\log M}}$	Complex
Hyperballs	$\frac{tm}{\log M}$	$\frac{t\sqrt{m}\sqrt{\log \frac{1}{\varepsilon} + \log M}}{\log M}$	$\frac{t\sqrt{m}}{\sqrt{\log M}}$	Simple

**Table 2.** Expected norm of signatures depending on the choice of distributions and (im)perfectness of rejection sampling.

The values of the table are obtained by computing the parameters for the underlying distributions (radii  $r, r'$  of the hypercubes or hyperballs and standard deviation  $\sigma$  of Gaussians) for our constraints  $M$  and  $t$ . This is done by computing their (smooth) Rényi Divergence, as done in Lemmas 5.1 and 5.2 for hyperballs. Proofs for hypercubes and Gaussians can be found in Appendix C. Given these parameters, the expected norm immediately follows ( $r\sqrt{m}$  for a hypercube of radius  $r$ ,  $\sigma r$  for a Gaussian of standard deviation  $\sigma$ , and  $r$  for a hyperball of radius  $r$ ). To conclude this section, we emphasize the following points:

- Gaussians and Hyperballs are asymptotically equivalent and reach the lower bounds in the bimodal setting; Hyperballs further reach our lower bound in the exact unimodal setting as well;
- Hyperballs benefit from a significantly simpler rejection test compared to Gaussians;
- The bimodal setting (in both Gaussian and Hyperballs cases) leads to the most compact signatures.

#### 5.4 Concrete Parameters

<sup>3</sup> To study the concrete impact of the choice of distributions on signature size, we consider Dilithium. The left side of Table 3 shows the parameters for three

<sup>3</sup> This subsection has been significantly updated compared to the previous version of this article. First, parameters were set for SIS with  $B > q$ . As estimating the hardness of SIS for such large norm bounds is difficult, we decided to change the parameters so that the SIS bound  $B$  is always lower than the modulus  $q$ . Second, the impact of the cut for hyperballs was underestimated in the Python script. We now

security levels of the round-3 documentation of the CRYSTALS-Dilithium submission to the NIST post-quantum project [BDK<sup>+</sup>20]. The right side of Table 3 gives updated parameters for Dilithium-G, a modification of Dilithium using Gaussian distributions whose description is available in the first version of the eprint version of [DKL<sup>+</sup>18]. For this updated version, we set the value of  $M$  to 4 and aim for security levels consistent with those of Dilithium. We also update the quantum costs by plugging in the improvements from [CL21].

	Hypercube-Uniform			Previous Gaussian		
	Medium	Recommended	Very High	Medium	Recommended	Very High
Ring dimension $\ell$	256	256	256	256	256	256
$q$	8380417	8380417	8380417	918529	918529	918529
$(n, m - n)$	(4, 4)	(6, 5)	(8, 7)	(3, 4)	(4, 5)	(6, 7)
$\eta$	2	4	2	1	1	1
$S$	N/A	N/A	N/A	50	55	65
$\tau$	39	49	60	39	49	60
$t = S \cdot \sqrt{\tau}$	N/A	N/A	N/A	312	385	503
$B$	N/A	N/A	N/A	386K	313K	457K
$\gamma_2$	$\frac{q-1}{88}$	$\frac{q-1}{32}$	$\frac{q-1}{32}$	$\frac{q-1}{256}$	$\frac{q-1}{256}$	$\frac{q-1}{128}$
$d$	13	13	13	12	11	11
$M$	4.25	5.1	3.85	4	4	4
BKZ block-size $b$ to break SIS	423 (417)	638 (603)	909 (868)	408 (350)	639 (552)	1018 (887)
Best known classical bit-cost	123 (121)	186 (176)	265 (253)	119 (102)	186 (161)	297 (259)
Best known quantum bit-cost	108 (107)	163 (157)	233 (223)	104 (89)	164 (141)	261 (227)
BKZ block-size $b$ to break LWE	422	622	860	471	619	934
Best known classical bit-cost	123	181	251	137	181	273
Best known quantum bit-cost	108	159	221	121	159	240
Expected signature size	2420	3293	4595	2009	2571	3706
Expected public key size	1312	1952	2592	800	1184	1760

**Table 3.** Parameters for Dilithium and updated Dilithium-G.

In these schemes, the verification key is a module-LWE sample  $\mathbf{Bs}_1 + \mathbf{s}_2$  where  $\mathbf{s}_1$  and  $\mathbf{s}_2$  have  $\ell_\infty$ -norms  $\leq \eta$ . For each coordinate, the lowest  $d$  bits are dropped. A parameter  $\tau$  is used to control the  $\ell_1$ -norm of any hashed value  $\mathbf{c}$ , so that  $\mathbf{c}$  has sufficient min-entropy. In Dilithium-G, the bound  $t$  is  $S\sqrt{\tau}$ , where  $S$  is the median over the key generation randomness of the largest singular value of  $(\text{rot}(\mathbf{s}_1)^\top, \text{rot}(\mathbf{s}_2)^\top)^\top$ . A rejection step is added in `KeyGen` to check that the key satisfies this bound. The value of the SIS bound corresponding to unforgeability is computed using [BDK<sup>+</sup>20, Equation (6)]. The strong unforgeability bound is obtained by multiplying this bound by 2. The security is estimated using block-size optimized BKZ to break the module-SIS or module-LWE instances.<sup>4</sup>

For Dilithium, i.e., the hypercube version, we take  $t_\infty = \tau\eta$  as a bound on the  $\ell_\infty$ -norm of the secret key, which drives the radius of the hypercube and subsequently the unforgeability SIS bound (in  $\ell_\infty$ -norm).

---

provide a Maple worksheet in our security estimate repository showing that this is not the case anymore, and the obtained values are now hard-coded in the Python script. This flaw was leading to unnecessarily large parameters for the hyperball uniforms.

<sup>4</sup> We use the scripts from <https://github.com/pq-crystals/security-estimates>.

It was argued in [DKL<sup>+</sup>18] that it seems difficult for BKZ to solve SIS with  $\ell_\infty$ -norm bound close to  $q$ , i.e.,  $\ell_2$ -norm above  $q$ . To analyze the runtime of BKZ in the case of an  $\ell_2$ -norm bound  $B \geq q$ , one can remove the trivial vectors of the input basis (i.e., the vectors with coordinates in  $q\mathbb{Z}$ ) by some randomizing step. This approach was however not considered for Dilithium-G and  $q$  was chosen such that  $B < q$ , leading to bigger parameters overall. Our updated parameters keep this constraint, as it is difficult to analyze the effectiveness of the attack presented in [DKL<sup>+</sup>18] when  $B > q$  for Euclidean norm.

Finally, the computation of the verification key and signature sizes (in bytes) is performed as in [BDK<sup>+</sup>20] and [DKL<sup>+</sup>18], respectively, with a different encoding. Namely, to compute signature sizes for the Gaussian version, we rely on a strategy explained in [ETWY22, Section 5], called range Asymmetric Numeral System, which allows one to encode the signature with an average bit-length reaching its entropy plus some constant overhead that we ignore. This technique is used to obtain the sizes in the right hand side of Table 3.

	Hyperball-Uniform			Improved Gaussian		
	Medium	Recommended	Very High	Medium	Recommended	Very High
Ring dimension $\ell$	256	256	256	256	256	256
$q$	520193	520193	520193	758273	758273	758273
$(n, m - n)$	(3, 4)	(4, 5)	(6, 7)	(3, 4)	(4, 5)	(6, 7)
$\eta$	1	1	1	1	1	1
$S$	50	55	65	50	55	65
$\tau$	39	49	60	39	49	60
$t = S \cdot \sqrt{\tau}$	312	385	503	312	385	503
$B$	197K	259K	246K	367K	265K	375K
$\gamma_2$	$\frac{q-1}{16}$	$\frac{q-1}{8}$	$\frac{q-1}{8}$	$\frac{q-1}{256}$	$\frac{q-1}{256}$	$\frac{q-1}{256}$
$d$	11	11	10	12	11	11
$M$	4	4	4	4	4	4
BKZ block-size $b$ to break SIS	447 (381)	628 (541)	1091 (946)	404 (347)	650 (560)	1041 (906)
Best Known Classical bit-cost	130 (111)	183 (158)	319 (276)	118 (101)	190 (163)	304 (264)
Best Known Quantum bit-cost	114 (97)	161 (139)	280 (243)	103 (89)	167 (143)	267 (232)
BKZ block-size $b$ to break LWE	494	650	977	478	629	948
Best Known Classical bit-cost	144	190	285	140	183	277
Best Known Quantum bit-cost	127	167	251	123	161	243
Expected signature size	1903	2473	3461	1921	2462	3553
Expected public key size	800	1056	1760	800	1184	1760

**Table 4.** Parameters for hyperball-uniform and improved Dilithium-G.

Next, we apply to Dilithium-G two modifications introduced in this work and introduce the Hyperball variant. In Table 4 (right side), we show the improvements we obtain when the standard deviation  $\sigma$  is computed using our refined bound from Lemma C.2 on the smooth Rényi divergence between two Gaussians and instantiated with  $\varepsilon = 2^{-64}$  instead of  $\varepsilon = 2^{-128}$ , as allowed by the use of Rényi divergence (as discussed in Section 4). Keeping  $M = 4$ , the standard deviation  $\sigma$  drops from  $11t$  to  $6.85t$  and leads to an additional saving on the signature size. When compared to Dilithium, if we consider the sum of signature and verification key sizes, we obtain up to  $\approx 30\%$  savings for the ‘Recommended’ parameter set, and  $\approx 25\%$  for the others.

Finally, we explore the use of the continuous uniform distributions in hyperballs. We take the algorithms from Dilithium-G, which are adapted to radial distributions and replace the Gaussians with the continuous uniform distributions in hyperballs, adding coefficient-wise rounding to integers when computing commitments. To set parameters, the bound  $B$  is computed using the radius of the hyperball instead of the probabilistic upper bound on the norm of a Gaussian vector. In Table 4 (left side), we provide the instantiations that we obtained. We note that the signature sizes are very similar to the ones obtained with Gaussians.

All figures of Tables 3 and 4 can be reproduced using scripts available at <https://github.com/jdevevey/security-estimates>.

## 6 Circumventing the Second Lower Bound via Bounded Rejection Sampling

We conclude this work by investigating an alternative way to perform rejection sampling which circumvents our lower bound on the expected number of loop iterations from Section 3.3. Notably, this approach makes the resulting signature scheme run within a given amount of time, which may be required in some practical applications (e.g., in real-time systems).

A first solution could be to set a bound on the maximal number of iterations, based on the run-time analysis from Lemma 2.3. However, this leads to a very large bound, of the order of  $\omega(\log \lambda + \log q_{\text{sig}}) / \log(M / (M - 1 + \varepsilon))$ , to ensure that with probability  $1 - \lambda^{-\omega(1)}$ , no signature among  $q_{\text{sig}}$  requires more iterations.

In the following, we propose a rejection sampling strategy that lets us fix an arbitrary bound  $i \geq 1$  on the number of iterations while still guaranteeing an output is produced at the end of the process. This strategy consists in first running  $i - 1$  iterations of the rejection sampling procedure. If something was output, then we are done, but if all iterations failed, we have to sample something that is related to the target distribution, in one-shot. For this last step, we use some sort of flooding. Note that, setting  $i = 1$ , one obtains one-shot signatures based on flooding, as in [ASY22]. Hence, this strategy can be seen as a generalization of both rejection sampling and flooding techniques.

### 6.1 Bounded Rejection Sampling Lemma

Let  $i \geq 1$  be an arbitrary bound for the number of loop iterations. Instead of simply having one distribution  $P_s$  to sample from, we now use two distributions  $P_f$  and  $P_s$ , where  $P_s$  is used for the rejection sampling part (the first  $i - 1$  iterations) and  $P_f$  is used in case of  $i - 1$  successive failures. If the divergences  $R_\infty(P_f \| P_s)$  and  $R_\infty(P_s \| P_t)$  are small, this strategy works. Moreover, the resulting distribution has a divergence with  $P_s$  and is a weighted mean of the classical rejection sampling-resulting distribution and the flooding distribution. This is what we prove in the following lemma.

**Lemma 6.1 (Bounded Rejection Sampling).** *Let  $p_f, p_t, p_s$  be probability densities, either all continuous or all discrete, and  $\varepsilon_0, \varepsilon_1 \geq 0, M_0, M_1 \geq 1$  with*

$$R_\infty^{\varepsilon_0}(p_f \| p_t) \leq M_0 \quad \text{and} \quad R_\infty^{\varepsilon_1}(p_t \| p_s) \leq M_1.$$

Then

$$R_\infty^{\frac{M}{M_0} \varepsilon_0}(\mathcal{B}_i^{\text{real}} \| \mathcal{B}_i^{\text{ideal}}) \leq M,$$

where

$$M = \left(1 - \left(1 - \frac{1}{M_1}\right)^{i-1}\right) \frac{1}{1 - \varepsilon_1} + \left(1 - \frac{1 + \varepsilon_1}{M_1}\right)^{i-1} \cdot M_0,$$

and  $\mathcal{B}_i^{\text{real}}$  and  $\mathcal{B}_i^{\text{ideal}}$  are defined in Figure 4.

Note that in the case where  $i = 1$ , distribution  $p_s$  is useless, as  $\mathcal{B}_1^{\text{real}}$  samples  $z \leftarrow p_f$  and returns it: this is flooding. Our lemma captures this situation, as  $M = M_0$  in that case. It is then not only a generalization of rejection sampling but also of flooding techniques.

Algorithms  $\mathcal{B}_i^{\text{ideal}}$  and  $\mathcal{B}_i^{\text{ideal}'}$  produce the same distribution for variable  $z$ , and hence Lemma 6.1 also holds when replacing  $\mathcal{B}_i^{\text{ideal}}$  by  $\mathcal{B}_i^{\text{ideal}'}$ . Algorithm  $\mathcal{B}_i^{\text{ideal}'}$  is more convenient when analyzing the adapted Lyubashevsky signature scheme.

<p>Algorithm <math>\mathcal{B}_i^{\text{real}}</math>:</p> <ol style="list-style-type: none"> <li>1: <math>\ell \leftarrow 1</math></li> <li>2: <b>while</b> <math>\ell \leq i - 1</math> <b>do</b></li> <li>3:   <math>z \leftarrow p_s</math></li> <li>4:   with probability <math>\min\left(\frac{p_t(z)}{M_1 \cdot p_s(z)}, 1\right)</math>,       return <math>z</math></li> <li>5:   <math>\ell \leftarrow \ell + 1</math></li> <li>6: <b>end while</b></li> <li>7: return <math>z \leftarrow p_f</math></li> </ol>	<p>Algorithm <math>\mathcal{B}_i^{\text{ideal}}</math>:</p> <ol style="list-style-type: none"> <li>1: return <math>z \leftarrow p_t</math></li> </ol>	<p>Algorithm <math>\mathcal{B}_i^{\text{ideal}'}</math>:</p> <ol style="list-style-type: none"> <li>1: <math>\ell \leftarrow 1</math></li> <li>2: <b>while</b> <math>\ell \leq i - 1</math> <b>do</b></li> <li>3:   <math>z \leftarrow p_t</math></li> <li>4:   with probability <math>\frac{1}{M_1}</math>,       return <math>z</math></li> <li>5:   <math>\ell \leftarrow \ell + 1</math></li> <li>6: <b>end while</b></li> <li>7: return <math>z \leftarrow p_t</math></li> </ol>
---	---	---

**Fig. 4.** Bounded rejection sampling algorithms.

*Proof.* With  $p_t$  and  $p_s$ , for  $\mathfrak{t} \in \{\text{real}, \text{ideal}\}$ , we can view  $\mathcal{B}_i^{\mathfrak{t}}$  as calling  $i - 1$  times  $\mathcal{A}^{\mathfrak{t}}$  from Figure 1, returning the value of the first call that does not abort, and if all calls failed, returning some independent sample  $z \leftarrow p_f$  (or  $p_t$ ). Using probability bounds from Lemma 2.2 and letting  $\mathcal{A}^{\text{real}}(\perp)$  denote the probability that  $\mathcal{A}^{\text{real}}$  aborts, we know that

$$\begin{aligned} \mathcal{B}_i^{\text{real}}(x) &= \left[ \sum_{0 \leq j \leq i-2} (\mathcal{A}^{\text{real}}(\perp))^j \cdot \min\left(\frac{p_t(x)}{M_1}, p_s(x)\right) \right] + (\mathcal{A}^{\text{real}}(\perp))^{i-1} \cdot p_f(x) \\ &= \frac{1 - (\mathcal{A}^{\text{real}}(\perp))^{i-1}}{1 - \mathcal{A}^{\text{real}}(\perp)} \cdot \min\left(\frac{p_t(x)}{M_1}, p_s(x)\right) + (\mathcal{A}^{\text{real}}(\perp))^{i-1} \cdot p_f(x) \\ &\leq \frac{1 - \left(1 - \frac{1}{M_1}\right)^{i-1}}{\frac{1 - \varepsilon_1}{M_1}} \cdot \frac{p_t(x)}{M_1} + \left(\frac{M_1 - 1 + \varepsilon_1}{M_1}\right)^{i-1} \cdot p_f(x). \end{aligned}$$

Let us define

$$M = \left(1 - \left(1 - \frac{1}{M_1}\right)^{i-1}\right) \cdot \frac{1}{1 - \varepsilon_1} + \left(\frac{M_1 - 1 + \varepsilon_1}{M_1}\right)^{i-1} \cdot M_0.$$

For this to be an upper bound on  $R_\infty^{\frac{M}{M_0}\varepsilon_0}(\mathcal{B}_i^{\text{real}}\|\mathcal{B}_i^{\text{ideal}})$ , it suffices that

$$\Pr_{x \leftarrow \mathcal{B}_i^{\text{real}}}[\mathcal{B}_i^{\text{real}}(x) > M \cdot p_t(x)] \leq \frac{M}{M_0}\varepsilon_0.$$

For any output  $x$  such that  $\mathcal{B}_i^{\text{real}}(x) > Mp_t(x)$ , it holds  $p_f(x) > M_0p_t(x)$  according to the above upper bound on  $\mathcal{B}_i^{\text{real}}(x)$ . This yields, by definition of  $M_0$ :

$$\Pr_{x \leftarrow p_f}[\mathcal{B}_i^{\text{real}}(x) > M \cdot p_t(x)] \leq \varepsilon_0.$$

The probability is however not taken over the desired distribution for  $x$ . Note that if we combine  $p_f(x) > M_0 \cdot p_t(x)$  with the above bound on the distribution of the output of  $\mathcal{B}_i^{\text{real}}$ , we get

$$\mathcal{B}_i^{\text{real}}(x) < \frac{M}{M_0} \cdot p_f(x).$$

Then  $\Pr_{x \leftarrow \mathcal{B}_i^{\text{real}}}[\mathcal{B}_i^{\text{real}}(x) > Mp_t(x)] < \frac{M}{M_0}\varepsilon_0$ .  $\square$

## 6.2 Lyubashevsky's Signature with Bounded Rejection

In this section, we present a way to modify Lyubashevsky's signature scheme by relying on bounded rejection sampling, as described above. This can be seen as a hybrid version between one-shot signatures which use flooding, as in [ASY22], and Lyubashevsky's unbounded signature.

Let  $k, n, m, q \geq 1$  specify matrix spaces with  $m > n$ . Let  $\mathcal{M}$  be the message space. Let  $H$  be a hash function modeled as a random oracle with domain  $\mathbb{Z}_q^n \times \mathcal{M}$  and range some finite set  $\mathcal{C} \subseteq \mathbb{Z}^k$ . Let  $\gamma > 0$ . Let  $\varepsilon_0, \varepsilon_1 \geq 0, M_0, M_1 \geq 1, i \geq 1$  be parameters related to bounded rejection sampling. Let  $\mathcal{S} \subseteq \mathbb{Z}^{m \times k}$ . Let  $P_0, P_1$  and  $P_2$  be three probability distributions over  $\mathbb{Z}^m$  satisfying

$$\max_{(\mathbf{s}, \mathbf{c}) \in \mathcal{S} \times \mathcal{C}} R_\infty^{\varepsilon_0}((P_0)_{+\mathbf{s}\mathbf{c}}\|P_1) \leq M_0 \quad \text{and} \quad \max_{(\mathbf{s}, \mathbf{c}) \in \mathcal{S} \times \mathcal{C}} R_\infty^{\varepsilon_1}(P_1\|(P_2)_{+\mathbf{s}\mathbf{c}}) \leq M_1.$$

Let  $(\mathbf{x}_0^\top | \mathbf{x}_1^\top)^\top \leftarrow P_0$  and  $(\mathbf{y}_0^\top | \mathbf{y}_1^\top)^\top \leftarrow P_2$ , where  $\mathbf{y}_0$  and  $\mathbf{x}_0$  take values in  $\mathbb{Z}^n$ . We present the modified scheme in Figure 5. The key generation algorithm is unchanged from Figure 2.

Before moving to the scheme analysis, let us define

$$M = \left(1 - \left(1 - \frac{1}{M_1}\right)^{i-1}\right) \frac{1}{1 - \varepsilon_1} + \left(1 - \frac{1 + \varepsilon_1}{M_1}\right)^{i-1} \cdot M_0.$$

The runtime of `Sign` is deterministically bounded, by at most  $i$  loop iterations. The correctness statement from Lemma 2.4 can be adapted as follows.

<p><b>Sign'</b>(<math>\mu, \mathbf{A}, \mathbf{S}</math>) :</p> <ol style="list-style-type: none"> <li>1: <math>\ell \leftarrow 1</math></li> <li>2: <b>if</b> <math>\ell \leq i - 1</math> <b>then</b></li> <li>3:     <math>\mathbf{y} \leftarrow P_2</math></li> <li>4: <b>else</b></li> <li>5:     <math>\mathbf{y} \leftarrow P_0</math></li> <li>6: <b>end if</b></li> <li>7: <math>\mathbf{c} \leftarrow H(\mathbf{A}\mathbf{y}, \mu)</math></li> <li>8: <math>\mathbf{z} \leftarrow \mathbf{y} + \mathbf{S}\mathbf{c}</math></li> <li>9: <math>u \leftarrow U([0, 1])</math></li> <li>10: <b>if</b> <math>u \leq \frac{P_1(\mathbf{z})}{M_1 P_2(\mathbf{y})}</math> or <math>\ell = i</math> <b>then</b></li> <li>11:     return (<math>\mathbf{z}, \mathbf{c}</math>)</li> <li>12: <b>else</b></li> <li>13:     <math>\ell \leftarrow \ell + 1</math></li> <li>14:     go to Step 2</li> <li>15: <b>end if</b></li> </ol>	<p><b>Verify</b>(<math>\mu, \mathbf{z}, \mathbf{c}, \mathbf{A}, \mathbf{T} = \mathbf{A}\mathbf{S}</math>) :</p> <ol style="list-style-type: none"> <li>1: <b>if</b> <math>\ \mathbf{z}\  \leq \gamma</math> and <math>\mathbf{c} = H(\mathbf{A}\mathbf{z} - \mathbf{T}\mathbf{c}, \mu)</math> <b>then</b></li> <li>2:     return 1</li> <li>3: <b>else</b></li> <li>4:     return 0</li> <li>5: <b>end if</b></li> </ol>
--	--

**Fig. 5.** Lyubashevsky's signature scheme with bounded rejection.

**Lemma 6.2 (Correctness).** *Let  $\varepsilon_0, \varepsilon_1 \geq 0$  and  $M_0, M_1 \geq 1$ . Let  $P_0, P_1, P_2$  satisfy  $\max_{(\mathbf{s}, \mathbf{c}) \in \mathcal{S} \times \mathcal{C}} R_{\infty}^{\varepsilon_b}(P_b \| P_{b+1}, +\mathbf{s}\mathbf{c}) \leq M_b$  for  $b \in \{0, 1\}$ . Let  $(\mathbf{x}_0^\top | \mathbf{x}_1^\top)^\top \leftarrow P_0$  and  $(\mathbf{y}_0^\top | \mathbf{y}_1^\top)^\top \leftarrow P_2$ , where  $\mathbf{x}_0$  and  $\mathbf{y}_0$  take values in  $\mathbb{Z}^n$ . Assume that  $\varepsilon_0 \leq \text{negl}(\lambda)$ ,  $M \leq \text{poly}(\lambda)$  and  $2^{-H_{\infty}(\mathbf{x}_0 | \mathbf{x}_1)_{P_0}}, 2^{-H_{\infty}(\mathbf{y}_0 | \mathbf{y}_1)_{P_2}} \leq \text{negl}(\lambda)$ . Then, in the ROM, the scheme is correct if  $\gamma \geq \gamma_{P_1}$  with  $\gamma_{P_1}$  such that  $\Pr_{\mathbf{z} \leftarrow P_1}(\|\mathbf{z}\| \geq \gamma_{P_1}) \leq \text{negl}(\lambda)$ .*

The main modification towards analyzing the security of the scheme from Figure 5, compared to the one from Figure 2, resides in the observation that the distributions of the pair  $(\mathbf{z}, \mathbf{c})$  obtained by the two processes from Figure 6 have  $\frac{M}{M_0} \varepsilon_0$ -smooth Rényi divergence of infinite order bounded by  $M$ . This is obtained by applying Lemma 6.1. Note that the hash function  $H$  needs to be consistently programmed for every  $\mathbf{c}$  that is produced, which is why we use the formalism of Algorithm  $\mathcal{B}_i^{\text{ideal}'}$  rather than Algorithm  $\mathcal{B}_i^{\text{ideal}}$ .

By the multiplicativity of the smooth Rényi divergence (Lemma A.6), we obtain that the  $(q_{\text{sig}} \cdot M\varepsilon_0/M_0)$ -smooth Rényi divergence between the adversary's views in games where the changes from Figure 6 have been applied to all signature queries, is bounded by  $M^{q_{\text{sig}}}$ . Probability preservation (Lemma A.5) can then be used meaningfully if  $q_{\text{sig}} \cdot M\varepsilon_0/M_0 = 2^{-\Omega(\lambda)}$  and  $M^{q_{\text{sig}}} \leq \text{poly}(\lambda)$ .

Once the signature queries are simulated without the signing key, the security proof can be completed as in prior works (see [Lyu09, Lyu12, AFLT16]).

**Asymptotic trade-off.** We now discuss the choices of the distributions  $P_0, P_1$  and  $P_2$ . We require that  $M^{q_{\text{sig}}} = \text{poly}(\lambda)$  and  $q_{\text{sig}} \cdot M\varepsilon_0/M_0 = 2^{-\Omega(\lambda)}$ , with  $\varepsilon_0, \varepsilon_1, M_0, M_1$  and  $M$  as in Lemma 6.1. We are aiming at not too large divergence bounds  $M_0, M_1, M$  as signatures typically become less efficient when they increase. For this reason, we set  $\varepsilon_0 = 2^{-\Omega(\lambda)}$ . As the condition  $M^{q_{\text{sig}}} = \text{poly}(\lambda)$  forces  $M$  to be close to 1, the condition  $q_{\text{sig}} \cdot M\varepsilon_0/M_0 = 2^{-\Omega(\lambda)}$  is already satisfied. We now focus on  $\varepsilon_1, M_0$  and  $M_1$ .

<pre> 1: <math>\ell \leftarrow 1</math> 2: <b>if</b> <math>\ell \leq i - 1</math> <b>then</b> 3:   <math>\mathbf{y} \leftarrow P_2</math> 4: <b>else</b> 5:   <math>\mathbf{y} \leftarrow P_0</math> 6: <b>end if</b> 7: <math>\mathbf{c} \leftarrow U(\mathcal{C})</math> 8: <math>\mathbf{z} \leftarrow \mathbf{y} + \mathbf{S}\mathbf{c}</math> 9: <math>u \leftarrow U([0, 1])</math> 10: <b>if</b> <math>u \leq \frac{P_1(\mathbf{z})}{M_1 P_2(\mathbf{y})}</math> or <math>\ell = i</math> <b>then</b> 11:   <b>return</b> <math>(\mathbf{z}, \mathbf{c})</math> 12: <b>else</b> 13:   <math>\ell \leftarrow \ell + 1</math> 14:   <b>go to</b> Step 2 15: <b>end if</b> </pre>	<pre> 1: <math>\ell \leftarrow 1</math> 2: <math>\mathbf{y} \leftarrow P_1</math> 3: <math>\mathbf{c} \leftarrow U(\mathcal{C})</math> 4: <math>\mathbf{z} \leftarrow \mathbf{y} + \mathbf{S}\mathbf{c}</math> 5: <math>u \leftarrow U([0, 1])</math> 6: <b>if</b> <math>u \leq \frac{1}{M_1}</math> or <math>\ell = i</math> <b>then</b> 7:   <b>return</b> <math>(\mathbf{z}, \mathbf{c})</math> 8: <b>else</b> 9:   <math>\ell \leftarrow \ell + 1</math> 10:   <b>go to</b> Step 2 11: <b>end if</b> </pre>
---	---

**Fig. 6.** Simulating signatures.

When  $i$  tends to infinity, we have  $M \approx 1/(1 - \varepsilon_1)$ , so that we can set  $\varepsilon_1 \approx 1/q_{\text{sig}}$  as in Section 4. For  $i = 1$ , we have  $M = M_0$ , and we fall in the regime of [ASY22, Section 4]. Let us now consider the small  $i$  case, which is probably the most interesting one for applications requiring a bounded signature time. As  $M \geq 1/(1 - \varepsilon_1)$  and we must ensure that  $M^{q_{\text{sig}}} = \text{poly}(\lambda)$ , we set  $\varepsilon_1$  at most of the order of  $1/q_{\text{sig}}$ . This implies that  $M \approx 1 + M_0 \cdot (1 - 1/M_1)^{i-1}$ , and hence we set  $(M_0 - 1) \cdot (1 - 1/M_1)^{i-1} = O(1/q_{\text{sig}})$ . For Gaussian and hyperball-uniform instantiations, this leads to a standard deviation (resp. radius) growing polynomially in  $q_{\text{sig}}/(1 - 1/M_1)^{i-1}$ .

We argue now that the trade-off above (for small  $i$ ) seems essentially optimal. For  $i = 1$ , it was showed in [ASY22, Appendix C.2] that the folklore statistical attack against the Gaussian and rejection-free version of Lyubashevsky’s signature scheme runs in subexponential time when  $M_0 = q_{\text{sig}}^{o(1)}$ . Now, for larger  $i$  and sufficiently distinct target and flooding distributions, an adversary could consider the signatures for which all loop iterations failed (i.e., the output sample corresponds to the flooding distribution), and run the statistical attack described in [ASY22] for those samples. As the probability of rejecting all samples is essentially  $(1 - 1/M_1)^{i-1}$ , this attack matches with the trade-off above.

**Acknowledgments.** The authors thank Wonhee Cho, Hyeongmin Choe and Orel Cosserson for pointing out errors in prior versions of this work, and Léo Ducas, Thomas Espitau and Eamonn Postlethwaite for insightful discussions on the hardness of the SIS problem with large Euclidean norm bound. The authors were supported by the AMIRAL ANR grant (ANR-21-ASTR-0016), the European Union Horizon 2020 Research and Innovation Program Grant 780701, the PEPR quantique France 2030 programme (ANR-22-PETQ-0008) and the PEPR Cyber France 2030 programme (ANR-22-PECY-0003).

## References

- AAB<sup>+</sup>19. S. Akleylek, E. Alkim, P. S. L. M. Barreto, N. Bindel, J. Buchmann, E. Eaton, G. Gutoski, J. Krämer, P. Longa, H. Polat, J. E. Ricardini, and G. Zanon. qTESLA round-3 candidate to the NIST post-quantum cryptography standardisation project, 2019. Available at <https://qtesla.org/>.
- ABB<sup>+</sup>17. E. Alkim, N. Bindel, J. Buchmann, Ö. Dagdelen, E. Eaton, G. Gutoski, J. Krämer, and F. Pawlega. Revisiting TESLA in the quantum random oracle model. In *PQCrypto*, 2017.
- AFLT16. M. Abdalla, P.-A. Fouque, V. Lyubashevsky, and M. Tibouchi. Tightly secure signatures from lossy identification schemes. *J. Cryptol.*, 2016.
- ASY22. S. Agrawal, D. Stehlé, and A. Yadav. Round-optimal lattice-based threshold signatures, revisited. In *ICALP*, 2022.
- BDK<sup>+</sup>20. S. Bai, L. Ducas, E. Kiltz, T. Lepoint, V. Lyubashevsky, P. Schwabe, G. Seiler, and D. Stehlé. CRYSTALS-DILITHIUM round-3 candidate to the NIST post-quantum cryptography standardisation project, 2020. Available at <https://pq-crystals.org/dilithium/>.
- BG14. S. Bai and S. D. Galbraith. An improved compression technique for signatures based on learning with errors. In *CT-RSA*, 2014.
- BGMN05. F. Barthe, O. Guédon, S. Mendelson, and A. Naor. A probabilistic approach to the geometry of the  $\ell_p^n$ -ball. *Ann Probab*, 2005.
- BLR<sup>+</sup>18. S. Bai, T. Lepoint, A. Roux-Langlois, A. Sakzad, D. Stehlé, and R. Steinfeld. Improved security proofs in lattice-based cryptography: Using the Rényi divergence rather than the statistical distance. *J. Cryptol.*, 2018.
- BP98. K. Ball and I. Perissinaki. The subindependence of coordinate slabs in  $\ell_p^n$  balls. *Israel J Math*, 1998.
- CL21. A. Chailloux and J. Loyer. Lattice sieving via quantum random walks. In *ASIACRYPT*. Springer International Publishing, 2021.
- Dat09. N. Datta. Min-and max-relative entropies and a new entanglement monotone. *T. Inform. Theory*, 2009.
- DDLL13. L. Ducas, A. Durmus, T. Lepoint, and V. Lyubashevsky. Lattice signatures and bimodal gaussians. In *CRYPTO*, 2013.
- Dev86. L. Devroye. *Non-Uniform random variate generation*. 1986.
- DKL<sup>+</sup>18. L. Ducas, E. Kiltz, T. Lepoint, V. Lyubashevsky, P. Schwabe, G. Seiler, and D. Stehlé. CRYSTALS-DILITHIUM: A lattice-based digital signature scheme. *TCHES*, 2018.
- DPSZ12. I. Damgård, V. Pastro, N. P. Smart, and S. Zakarias. Multiparty computation from somewhat homomorphic encryption. In *CRYPTO*, 2012.
- ETWY22. T. Espitau, M. Tibouchi, A. Wallet, and Y. Yu. Shorter hash-and-sign lattice-based signatures. In *CRYPTO*, 2022.
- FS86. A. Fiat and A. Shamir. How to prove yourself: Practical solutions to identification and signature problems. In *CRYPTO*, 1986.
- GLP15. T. Güneysu, V. Lyubashevsky, and T. Pöppelmann. Lattice-based signatures: Optimization and implementation on reconfigurable hardware. *T. Comput.*, 2015.
- HJMR07. P. Harsha, R. Jain, D. McAllester, and J. Radhakrishnan. The communication complexity of correlation. In *CCC*, 2007.
- LNP22. V. Lyubashevsky, N. K. Nguyen, and M. Plançon. Lattice-based zero-knowledge proofs and applications: Shorter, simpler, and more general. In *CRYPTO*, 2022.

- LS15. A. Langlois and D. Stehlé. Worst-case to average-case reductions for module lattices. *Des. Codes Cryptogr.*, 2015.
- LSS14. A. Langlois, D. Stehlé, and R. Steinfeld. GGHLite: More efficient multilinear maps from ideal lattices. In *EUROCRYPT*, 2014.
- Lyu09. V. Lyubashevsky. Fiat-Shamir with aborts: Applications to lattice and factoring-based signatures. In *ASIACRYPT*, 2009.
- Lyu12. V. Lyubashevsky. Lattice signatures without trapdoors. In *EUROCRYPT*, 2012.
- Lyu16. V. Lyubashevsky. Digital signatures based on the hardness of ideal lattice problems in all rings. In *ASIACRYPT*, 2016.
- MR07. D. Micciancio and O. Regev. Worst-case to average-case reductions based on Gaussian measures. *J. Comput.*, 2007.
- MV10. D. Micciancio and P. Voulgaris. Faster exponential time algorithms for the shortest vector problem. In *SODA*, 2010.
- Pre17. T. Prest. Sharper bounds in lattice-based cryptography using the Rényi divergence. In *ASIACRYPT*, 2017.
- Ren05. R. Renner. *Security of quantum key distribution*. PhD thesis, ETH Zurich, 2005.
- RW04. R. Renner and S. Wolf. Smooth Rényi entropy and applications. In *ISIT*, 2004.
- Sch91. C.-P. Schnorr. Efficient signature generation by smart cards. *J. Cryptol.*, 1991.
- vEH14. T. van Erven and P. Harremoës. Rényi divergence and Kullback-Leibler divergence. *T. Inform. Theory*, 2014.

## A Additional Background Material

### A.1 Notations

When we consider a probability density, this is with respect to the canonical (i.e., Lebesgue or counting) measure  $\mu$  over their support. We may identify the notion of probability distribution and probability density in the discrete case. Given a distribution  $P$  with density  $p$ , we let  $x \leftarrow P$  denote the sampling of  $x$  according to  $P$ . We let  $\text{Supp}(P)$  denote the smallest (for inclusion) set such that for any set  $X$ ,  $P(X \cap \text{Supp}(P)) = P(X)$ . Namely, if  $P$  is discrete, then  $\text{Supp}(P) = \{x | P(x) \neq 0\}$ . Given a set  $S \subseteq \text{Supp}(P)$ , we let  $P^S$  denote the distribution  $P$  cut to  $S$ , i.e., the measure  $P/P(S)$  restricted to  $S$ . Given two probability distributions  $F$  and  $G$  with densities  $f$  and  $g$  we let  $F \otimes G$  denote the distribution of  $(x, y)$  where  $x \leftarrow F$  and  $y \leftarrow G$  are independent and  $f \otimes g$  one of its density. Given a distribution  $F$  with density  $f$  and an element  $\mathbf{x}$ , we let  $F_{+\mathbf{x}}$  denote the distribution with density  $f_{+\mathbf{x}} : \mathbf{y} \mapsto f(\mathbf{y} - \mathbf{x})$ . Given a finite set  $S$ , we let  $U(S)$  denote the uniform distribution over  $S$ . By notation abuse, we use algorithm names to denote the random variable associated to their output.

Given a dimension  $m \geq 1$ , a center  $\mathbf{c} \in \mathbb{R}^m$  and a radius  $r > 0$ , we let  $\mathcal{B}_m^p(r, \mathbf{c})$  (resp.  $\mathcal{S}_m^p(r, \mathbf{c})$ ) denote the  $p$ -norm ball (resp. sphere) of radius  $r$  and center  $\mathbf{c}$  for  $p \in [1, +\infty]$ . When  $p = 2$  (resp.  $\mathbf{c} = 0$ ), we omit it. We also let  $V_m(r) := \frac{\pi^{m/2}}{\Gamma(m/2+1)} r^m$  denote its volume as well as  $S_m = m \cdot V_m(1)$  denote the surface of the unit sphere.

Given a set and a subset  $S \subseteq Y$  we let  $\chi_S : x \mapsto \{1 \text{ if } x \in S, 0 \text{ if } x \in Y \setminus S\}$  denote the indicator function of  $S$ . Let  $\lceil \cdot \rceil : \mathbb{R} \rightarrow \mathbb{Z}$  be the rounding operator that maps  $x$  to the nearest integer (in case of a tie, it is rounded downwards). It is naturally extended to  $\mathbb{R}^n$  by coordinate-wise application. The notation  $\log$  refers to the natural logarithm.

For a parameter  $\lambda$  going to infinity, we use the notations  $\text{negl}(\lambda) = \lambda^{-\omega(1)}$  and  $\text{poly}(\lambda) = \lambda^{O(1)}$ . Let  $H : \mathcal{D} \rightarrow \mathcal{R}$  be a function with a finite range  $\mathcal{R}$ . It is said to be modeled as a random oracle if it is replaced by a uniformly sampled function among those from  $\mathcal{D}$  to  $\mathcal{R}$ .

## A.2 Information-Theoretic Tools

We will use the following instantiation of conditional min-entropy.

**Definition A.1 (Min-Entropy).** *Let  $X = (Y, Z)$  be a random variable. Let  $p_X, p_Z$  be the densities of  $X$  and  $Z$ , and  $p_{Y|Z=z}$  the density of  $Y$  conditioned on  $Z = z$ . The conditional min-entropy of  $Y$  on  $Z$  is:*

$$H_\infty(Y|Z)_{p_X} = -\log \left( \int_{\text{Supp}(p_Z)} p_Z(z) \max_{y \in \text{Supp}(p_{Y|Z=z})} p_{Y|Z=z}(y) \, d\mu(z) \right).$$

To quantify similarities between distributions, we consider the statistical distance or a Rényi divergence.

**Definition A.2 (Statistical Distance).** *Let  $P, Q$  be two probability distributions with respective densities  $p, q$ . Their statistical distance is*

$$\Delta(P, Q) := \frac{1}{2} \int_{\text{Supp}(P) \cup \text{Supp}(Q)} |p(x) - q(x)| \, d\mu(x).$$

**Definition A.3 (Rényi divergence).** *Let  $a \in (1, +\infty)$ . Let  $P, Q$  be two probability distributions with  $P$  absolutely continuous with respect to  $Q$ . Their Rényi divergence of order  $a$ , assuming that it exists, is*

$$R_a(P||Q) := \left( \int_{\text{Supp}(P)} \left( \frac{dP}{dQ}(x) \right)^{a-1} dP(x) \right)^{\frac{1}{a-1}}.$$

Their Rényi divergence of infinite order is

$$R_\infty(P||Q) := \text{ess sup}_{x \in \text{Supp}(P)} \frac{dP}{dQ}(x).$$

By notation abuse, we may use random variables instead of probability densities as arguments, for the notions defined above.

The following lemma lists standard properties of the Rényi divergence.

**Lemma A.4 ([vEH14]).** *Let  $X$  and  $Y$  be two random variables with probability distributions  $P_X$  and  $P_Y$  such that  $\text{Supp}(P_X) \subseteq \text{Supp}(P_Y)$ . The following holds for any order  $a \in (1, +\infty]$ .*

- **Log. Positivity:**  $R_a(P_X \| P_Y) \geq R_a(P_X \| P_X) = 1$ .
- **Data Processing Inequality:**  $R_a(P_{f(X)} \| P_{f(Y)}) \leq R_a(P \| Q)$  for any function  $f$ , where  $P_{f(Z)}$  denotes the distribution of  $f(Z)$  for  $Z = X$  or  $Y$ .
- **Multiplicativity:** Let  $X = (X_1, X_2)$  and  $Y = (Y_1, Y_2)$ . Let  $P_{X_1}$  and  $P_{Y_1}$  denote the probability distribution of  $X_1$  and  $Y_1$ . Let  $P_{X_2|X_1=x}$  and  $P_{Y_2|Y_1=x}$  denote the distribution of  $X_2$  conditioned on  $X_1 = x$  as well as  $Y_2$  conditioned on  $Y_1 = x$ . If  $X_1$  and  $X_2$  are independent and  $Y_1$  and  $Y_2$  are independent, then

$$R_a(P_X \| P_Y) = R_a(P_{X_1} \| P_{Y_1}) R_a(P_{X_2|X_1} \| P_{Y_2|Y_1}).$$

Otherwise

$$R_a(P_X \| P_Y) \leq R_\infty(P_{X_1} \| P_{Y_1}) \cdot \max_{x \in \text{Supp}(P_{X_1})} R_a(P_{X_2|X_1=x} \| P_{Y_2|Y_1=x}).$$

- **Probability Preservation:** For any event  $E \subseteq \text{Supp}(P_Y)$ ,

$$P_Y(E) \geq \frac{P_X(E)^{\frac{a}{a-1}}}{R_a(P_X \| P_Y)}.$$

Moreover the Rényi divergence is non-decreasing and continuous as a function of  $a \in [1, +\infty]$ , as long as it is finite.

### A.3 Properties of the Smooth Rényi Divergence

We first prove that the two definitions of Definition 2.1 are indeed equivalent.

*Proof (Definition 2.1).* We prove that both definitions are indeed equivalent. Let  $R_\infty^\varepsilon(p \| q)$  be the first quantity and  $\mathcal{R}_\infty^\varepsilon(p \| q)$  be the second one.

Let  $S \subseteq \text{Supp}(q)$  such that  $\int_S p(x) d\mu(x) \geq 1 - \varepsilon$ . Let  $M = \sup_{x \in S} \frac{p(x)}{q(x)}$ . This means that

$$\Pr_{x \leftarrow p} [p(x) \leq Mq(x)] \geq \int_S p(x) d\mu(x) \geq 1 - \varepsilon.$$

Then  $\mathcal{R}_\infty^\varepsilon(p \| q) \leq M$ . By definition of  $R_\infty^\varepsilon(p \| q)$ , this implies the first inequality

$$\mathcal{R}_\infty^\varepsilon(p \| q) \leq R_\infty^\varepsilon(p \| q).$$

Now let  $M > 0$  such that  $\Pr_{x \leftarrow p}(p(x) \leq Mq(x)) \geq 1 - \varepsilon$ . Define

$$S := \{x \in \text{Supp}(p) \cup \text{Supp}(q) \mid p(x) \leq Mq(x)\}.$$

Then  $\int_S p(x) d\mu(x) \geq 1 - \varepsilon$  by definition. Note that if we choose  $S' = S \cap \text{Supp}(p)$  we have  $\int_{S'} p(x) d\mu(x) = \int_S p(x) d\mu(x)$  as we only removed elements that were not in the support of  $p$ . Moreover, assume that there exists  $x \in S'$  such that  $x \notin \text{Supp}(q)$ . We would have  $p(x) \leq M \cdot 0 = 0$ , contradicting the fact that  $x \in \text{Supp}(p)$ . Then it holds that  $S' = \{x \in \text{Supp}(q) \mid p(x) \leq Mq(x)\}$ . We have

$$M \geq \sup_{x \in S'} \frac{p(x)}{q(x)}, \int_{S'} p(x) d\mu(x) \geq 1 - \varepsilon \quad \text{and} \quad S' \subseteq \text{Supp}(q).$$

This implies, by definition of  $R_\infty^\varepsilon(p\|q)$  that  $M \geq R_\infty^\varepsilon(p\|q)$ . By definition of the second quantity, we have the inequality

$$R_\infty^\varepsilon(p\|q) \leq \mathcal{R}_\infty^\varepsilon(p\|q),$$

thus completing the proof of the equality.  $\square$

We now give a few properties of the smooth Rényi divergence. The probability preservation and multiplicativity properties are used in the security proof of our signature variant with a bounded number of rejection steps. The comparison to the Rényi divergence is used to bound the smooth Rényi divergence between Gaussian distributions.

We start by proving a probability preservation property.

**Lemma A.5 (Probability Preservation).** *Let  $P, Q$  be two distributions. For any  $\varepsilon \geq 0$  such that  $R_\infty^\varepsilon(P\|Q)$  is finite, the following holds.*

$$\forall E \subseteq \text{Supp}(P), P(E) \leq R_\infty^\varepsilon(P\|Q) \cdot Q(E) + \varepsilon.$$

*Proof.* Let  $S := \{x \in \text{Supp}(P) \mid P(x) \leq R_\infty^\varepsilon(P\|Q) \cdot Q(x)\}$ . We decompose the event  $E$  into the disjoint union  $(E \cap S) \cup (E \setminus S)$ . The following holds:

- $P(E \cap S) \leq R_\infty^\varepsilon(P\|Q) \cdot Q(E \cap S) \leq R_\infty^\varepsilon(P\|Q) \cdot Q(E)$ , by definition of  $S$ ,
- $P(E \setminus S) \leq \varepsilon$ , by definition of  $R_\infty^\varepsilon(P\|Q)$ .

Combining both inequalities yields the result.  $\square$

In the case of signatures, the security loss may depend on the number of signing queries. The following result may then prove useful.

**Lemma A.6 (Multiplicativity).** *Let  $(X_1, X_2)$  (resp.  $(Y_1, Y_2)$ ) be a random variable with probability density  $p_{X_1 X_2}$  (resp.  $p_{Y_1 Y_2}$ ). Let  $p_{X_1}$  (resp.  $p_{Y_1}$ ) be the density of  $X_1$  (resp.  $Y_1$ ). For any  $x$ , let us denote by  $p_{X_2|X_1=x}$  (resp.  $p_{Y_2|Y_1=x}$ ) the probability density of  $X_2$  (resp.  $Y_2$ ) conditioned on  $X_1 = x$  (resp.  $Y_1 = x$ ). Then for any  $\varepsilon_1, \varepsilon_2 \geq 0$  it holds that*

$$R_\infty^{\varepsilon_1 + \varepsilon_2}(p_{X_1 X_2} \| p_{Y_1 Y_2}) \leq R_\infty^{\varepsilon_1}(p_{X_1} \| p_{Y_1}) \cdot \sup_{\substack{x \in \text{Supp}(p_{X_1}) \\ \cap \text{Supp}(p_{Y_1})}} R_\infty^{\varepsilon_2}(p_{X_2|X_1=x} \| p_{Y_2|Y_1=x}).$$

*Proof.* Let  $R_1 = R_\infty^{\varepsilon_1}(p_{X_1} \| p_{Y_1})$  and  $R_2 = \sup_{x \in S_1} R_\infty^{\varepsilon_2}(p_{X_2|X_1=x} \| p_{Y_2|Y_1=x})$ , with  $S_1 = \text{Supp}(p_{X_1}) \cap \text{Supp}(p_{Y_1})$ . If  $R_1 = +\infty$ , the statement is vacuously true. Let us now assume that this is not the case.

We now define  $R = R_1 \cdot R_2$  and

$$S = \{(x, y) \in \text{Supp}(p_{X_1 X_2}) \mid p_{X_1 X_2}(x, y) > R \cdot p_{Y_1 Y_2}(x, y)\}.$$

Any pair  $(x, y) \in S$  either satisfies  $p_{X_1}(x)p_{X_2|X_1=x}(y) > R p_{Y_1}(x)p_{Y_2|Y_1=x}(y)$  or  $x \notin \text{Supp}(p_{Y_1})$ . This implies that we have either  $p_{X_1}(x) > R_1 p_{Y_1}(x)$  or

$p_{X_2|X_1=x}(y) > R_2 p_{Y_2|Y_1=x}(y)$ . We then have, using the union bound,

$$\begin{aligned} \int_S p_{X_1 X_2}(\mathbf{x}) \, d\mathbf{x} &\leq \Pr_{x \leftarrow p_{X_1}} [p_{X_1}(x) > R_1 \cdot p_{Y_1}(x)] \\ &\quad + \sum_{x \in S_1} p_{X_1}(x) \cdot \Pr_{y \leftarrow p_{X_2|X_1=x}} [p_{X_2|X_1=x}(y) > R_2 \cdot p_{Y_2|Y_1=x}(y)] \\ &\leq \varepsilon_1 + \sum_{x \in S_1} p_{X_1}(x) \varepsilon_2 \\ &\leq \varepsilon_1 + \varepsilon_2. \end{aligned}$$

Define the set

$$\bar{S} := \text{Supp}(p_{Y_1 Y_2}) \setminus S = \{(x, y) \in \text{Supp}(p_{Y_1 Y_2}) \mid p_{X_1 X_2}(x, y) \leq R \cdot p_{Y_1 Y_2}(x, y)\}.$$

We have  $\bar{S} = (\text{Supp}(p_{X_1 X_2}) \cup \text{Supp}(p_{Y_1 Y_2})) \setminus S$ , as  $\text{Supp}(p_{X_1 X_2}) \setminus \text{Supp}(p_{Y_1 Y_2}) \subseteq S$ . Then it satisfies  $\int_{\bar{S}} p_{X_1 X_2}(\mathbf{x}) \, d\mathbf{x} \geq 1 - \varepsilon_1 - \varepsilon_2$ . The first definition of the smooth divergence provides the result.  $\square$

Noticing that  $R_a(P\|Q)^{a-1} = \mathbb{E}_{x \leftarrow P}((p(x)/q(x))^{a-1})$ , we can apply concentration inequalities to compare the smooth divergence and the Rényi divergence, as was done in [RW04] for entropies. We however recall that the smooth Rényi divergence may be finite for pairs of random variables for which the Rényi divergence is infinite, in which case our bound is trivial.

**Lemma A.7.** *Let  $X, Y$  be two discrete random variable with probability distributions  $P_X$  and  $P_Y$ . For any  $\varepsilon \geq 0$  and order  $a \in (1, +\infty)$  it holds*

$$R_\infty^\varepsilon(P_X\|P_Y) \leq \frac{R_a(P_X\|P_Y)}{\varepsilon^{1/(a-1)}} \quad \text{and} \quad R_\infty^\varepsilon(P_X\|P_Y) \leq R_\infty(P_X\|P_Y).$$

*Proof.* Markov's inequality gives that for any  $t > 0$ ,

$$\Pr_{x \leftarrow P_X} \left( \left( \frac{P_X(x)}{P_Y(x)} \right)^{a-1} \geq t \right) \leq \frac{R_a(P_X\|P_Y)^{a-1}}{t}.$$

Setting  $t_0$  such that  $R_a(P_X\|P_Y)^{a-1}/t_0 = \varepsilon$ , we have:

$$\Pr_{x \leftarrow P_X} [P_X(x) \geq t_0^{1/(a-1)} \cdot P_Y(x)] \leq \varepsilon.$$

By the second definition of  $R_\infty^\varepsilon(P_X\|P_Y)$ , this shows

$$R_\infty^\varepsilon(P_X\|P_Y) \leq t_0^{1/(a-1)} = \frac{R_a(P_X\|P_Y)}{\varepsilon^{\frac{1}{a-1}}}.$$

To conclude the proof, recall that the Rényi divergence is continuous as a function of  $a$ . Taking the limit of this upper bound when  $a$  tends to  $+\infty$  gives the second result.  $\square$

#### A.4 Digital Signatures

Here we briefly recall the formalism of digital signatures. Our definition slightly differs from more standard ones as we consider the case where correctness and signature runtime may only hold in the Random Oracle Model (ROM) rather than unconditionally.

**Definition A.8.** *A signature scheme is a tuple  $(\text{KeyGen}, \text{Sign}, \text{Verify})$  of algorithms with the following specifications:*

- $\text{KeyGen} : 1^\lambda \rightarrow (\text{vk}, \text{sk})$  takes as input a security parameter  $\lambda$  and outputs a verification key  $\text{vk}$  and a signing key  $\text{sk}$ .
- $\text{Sign} : (\text{sk}, \mu) \rightarrow \sigma$  takes as inputs a signing key  $\text{sk}$  and a message  $\mu$  and outputs a signature  $\sigma$ .
- $\text{Verify} : (\text{vk}, \mu, \sigma) \rightarrow b \in \{0, 1\}$  takes as inputs a verification key  $\text{vk}$ , a message  $\mu$  and a signature  $\sigma$  and accepts (1) or rejects (0).

We say that it is correct if for any pair  $(\text{vk}, \text{sk})$  in the range of  $\text{KeyGen}$  and  $\mu$ ,

$$\Pr(\text{Verify}(\text{vk}, \mu, \text{Sign}(\text{sk}, \mu)) = 1) \geq 1 - \text{negl}(\lambda),$$

where the probability is taken over the random coins of the two algorithms. We say that it is correct in the ROM if the above holds when the probability is also taken over the randomness of the random oracle modeling some hash function used in the scheme.

Second, we define the Strong Existential Unforgeability Chosen Message Attack (sEU-CMA) security game for digital signatures.

**Definition A.9.** *Let  $T, \delta > 0$ . A signature scheme  $(\text{KeyGen}, \text{Sign}, \text{Verify})$  is said to be  $(T, \delta)$ -sEU-CMA secure if no adversary  $\mathcal{A}$  with runtime  $\leq T$  given  $\text{vk}$  and access to a signing oracle has probability  $\geq \delta$  over the choice of the signing and verification keys  $(\text{vk}, \text{sk}) \leftarrow \text{KeyGen}(1^\lambda)$  and its random coins of outputting  $(\mu^*, \sigma^*)$  such that*

- If  $\mu^*$  was queried to the signing oracle, it did not return  $\sigma^*$ : the forged signature must be fresh,
- $\text{Verify}(\text{vk}, \mu^*, \sigma^*) = 1$ : the forged signature must be accepted.

The scheme is said  $(T, \delta)$ -sEU-CMA secure in the ROM if the above holds when the adversary can also make queries to a random oracle that models some hash function used in the scheme. The probability of forging a signature is also called the advantage of  $\mathcal{A}$ .

#### A.5 The LWE and SIS Problems

The Learning With Errors (LWE) and Short Integer Solution (SIS) problems serve as security foundation of Lyubashevsky's signature schemes. In the parameter instantiation section, we will use their module counterparts (see [LS15]).

**Definition A.10 (SIS).** Let  $m \geq n \geq 1$ ,  $q \geq 2$  and  $\beta > 0$ . The SIS problem with parameters  $m, n, q, \beta$  is as follows: given as input  $\mathbf{A} \leftarrow U(\mathbb{Z}_q^{n \times m})$ , the goal is to find  $\mathbf{x} \in \mathbb{Z}^m$  such that  $\mathbf{A}\mathbf{x} = \mathbf{0} \pmod{q}$  and  $0 < \|\mathbf{x}\| \leq \beta$ .

**Definition A.11 (LWE).** Let  $m \geq n \geq 1$ ,  $q \geq 2$  and  $\chi$  a distribution over  $\mathbb{Z}_q$ . The LWE problem with parameters  $m, n, q, \chi$  consists in distinguishing between the distributions  $(\mathbf{A}, \mathbf{A}\mathbf{s} + \mathbf{e})$  and  $(\mathbf{A}, \mathbf{u})$ , where  $\mathbf{A} \leftarrow U(\mathbb{Z}_q^{m \times n})$ ,  $\mathbf{u} \leftarrow U(\mathbb{Z}_q^n)$ ,  $\mathbf{s} \leftarrow \chi^n$  and  $\mathbf{e} \leftarrow \chi^m$ .

## A.6 Beta Function and Hyperspherical Caps

The beta function is a special function that is related to the gamma function and binomial coefficients.

**Definition A.12 (Regularized Incomplete Beta Function).** The incomplete beta function is defined over  $[0, 1] \times \mathbb{R}^+ \times \mathbb{R}^+$  as

$$B : (x; a, b) \mapsto \int_0^x t^{a-1}(1-t)^{b-1} dt.$$

When  $x = 1$ , this is the Beta function, and we use the notation  $B(a, b)$  in that case. For fixed  $a, b$ , the function  $I(a, b) : x \mapsto B(x; a, b)/B(a, b)$  is invertible.

The function  $x \mapsto I_x(a, b)$  and its inverse are useful when we consider the area of a hyperspherical cap, as defined below.

**Lemma A.13 (Hyperspherical Cap).** Let  $n > 1$ ,  $\eta > 1$  and  $\mathbf{x} \in \mathcal{S}_n(1)$ . The set  $\{\mathbf{y} \in \mathcal{B}_n(1) \mid \langle \mathbf{y}, \mathbf{x} \rangle \geq 1/\eta\}$  is the intersection of a half-space and the unit hyperball. It is called a hyperspherical cap and has volume  $V_\eta$  and area  $A_\eta$  satisfying

$$V_\eta = \frac{V_n(1)}{2} \cdot I_{1-\frac{1}{\eta^2}}\left(\frac{n+1}{2}, \frac{1}{2}\right) \quad \text{and} \quad A_\eta = \frac{S_n}{2} \cdot I_{1-\frac{1}{\eta^2}}\left(\frac{n-1}{2}, \frac{1}{2}\right).$$

By placing an appropriate cone inside the hyperspherical cap (see [MV10, Lemma 4.1]), we have:

$$I_{1-\frac{1}{\eta^2}}\left(\frac{n+1}{2}, \frac{1}{2}\right) > \left(1 - \frac{1}{\eta^2}\right)^{n-\frac{1}{2}} \cdot \frac{1 - \frac{1}{\eta}}{n}.$$

By placing the hyperspherical cap into a cylinder of 1-dimensional height  $1 - 1/\eta$ , we have:

$$I_{1-\frac{1}{\eta^2}}\left(\frac{n+1}{2}, \frac{1}{2}\right) < \left(1 - \frac{1}{\eta^2}\right)^{n-1} \cdot n \cdot \left(1 - \frac{1}{\eta}\right).$$

Letting  $\varepsilon$  denote  $I_{1-1/\eta^2}(\frac{n+1}{2}, \frac{1}{2})$ , we obtain the following consequence of the above two inequalities, which we use to estimate the smooth Rényi divergence between uniform distributions in hyperballs:

$$1 - (2n\varepsilon)^{\frac{1}{n+1/2}} < \frac{1}{\eta^2} < 1 - \left(\frac{\varepsilon}{n}\right)^{\frac{1}{n-1}}.$$

For  $\varepsilon = 2^{-c \cdot n}$  for a constant  $c > 0$ , we obtain that  $1/\eta^2$  tends to  $1 - 2^{-c}$  when  $n$  goes to infinity. For  $\varepsilon$  satisfying  $\varepsilon \geq 2^{-o(n)}$  and  $\varepsilon = o(1/n)$  with  $n$  going to infinity, we obtain that  $1/\eta^2 \sim -\ln(\varepsilon)/n$ .

## A.7 Gaussian Distributions

**Definition A.14 (Gaussian Distribution).** Let  $m \geq 1, \sigma > 0$  and  $\mathbf{v} \in \mathbb{R}^m$ . Define  $\rho_\sigma : \mathbf{x} \mapsto \exp(-\|\mathbf{x}\|^2/(2\sigma^2))$ . The discrete Gaussian distribution with standard deviation parameter  $\sigma$  and center parameter  $\mathbf{v}$  is defined as

$$D_{\mathbb{Z}^m, \sigma, \mathbf{v}} : \mathbf{z} \mapsto \frac{\rho_\sigma(\mathbf{z} - \mathbf{v})}{\rho_{\sigma, \mathbf{v}}(\mathbb{Z}^m)},$$

where we let  $\rho_{\sigma, \mathbf{v}}(\mathbb{Z}^m)$  denote  $\sum_{\mathbf{x} \in \mathbb{Z}^m} \rho_\sigma(\mathbf{x} - \mathbf{v})$ . If  $\mathbf{v} = \mathbf{0}$ , we omit it in the subscript.

The divergence between two discrete Gaussian distributions is well-known (see, e.g., [LSS14, Lemma 4.2] for  $a = 2$ ). We give a formulation that includes every order  $\geq 1$ , while restricting our case to the  $\mathbb{Z}^m$  lattice. It may be proved by adapted the proof of [LSS14].

**Lemma A.15.** Let  $m \geq 1, \sigma > 0$  and  $\mathbf{v} \in \mathbb{Z}^m$ . Then for any  $a \in [1, +\infty)$ :

$$R_a(D_{\mathbb{Z}^m, \sigma} \| D_{\mathbb{Z}^m, \sigma, \mathbf{v}}) = \exp\left(a \frac{\|\mathbf{v}\|^2}{2\sigma^2}\right).$$

We also have  $R_\infty(D_{\mathbb{Z}^m, \sigma} \| D_{\mathbb{Z}^m, \sigma, \mathbf{v}}) = +\infty$  if  $\mathbf{v} \neq \mathbf{0}$ .

We also consider bimodal Gaussian distributions.

**Definition A.16 (Bimodal Gaussian Distribution).** Let  $m \geq 1$ . The bimodal Gaussian distribution  $BD_{\mathbb{Z}^m, \sigma, \mathbf{v}}$  with parameters  $\sigma > 0$  and  $\mathbf{v} \in \mathbb{R}^m$  is the distribution obtained by sampling  $b \leftarrow U(\{-1, 1\})$ , and returning  $\mathbf{x} \leftarrow D_{\mathbb{Z}^m, \sigma, b\mathbf{v}}$ . It can be expressed as

$$BD_{\mathbb{Z}^m, \sigma, \mathbf{v}} : \mathbf{z} \mapsto \frac{1}{2} (D_{\mathbb{Z}^m, \sigma, \mathbf{v}}(\mathbf{z}) + D_{\mathbb{Z}^m, \sigma, -\mathbf{v}}(\mathbf{z})).$$

In particular, since  $\rho_{\sigma, \mathbf{v}}(\mathbb{Z}^m) = \rho_{\sigma, -\mathbf{v}}(\mathbb{Z}^m)$  (which can be seen by reordering the sum), we can write

$$BD_{\mathbb{Z}^m, \sigma, \mathbf{v}}(\mathbf{z}) = \frac{1}{\rho_{\sigma, \mathbf{v}}(\mathbb{Z}^m)} \exp\left(\frac{-\|\mathbf{z}\|^2 - \|\mathbf{v}\|^2}{2\sigma^2}\right) \cosh\left(\frac{|\langle \mathbf{z}, \mathbf{v} \rangle|}{\sigma^2}\right).$$

## B Missing proofs

### B.1 Proof of Lemma 2.2

Let  $S = \text{Supp}(p_t) \cup \text{Supp}(p_s)$ . Let us write for any  $x \in \text{Supp}(p_s)$ :

$$\min\left(\frac{p_t(x)}{M p_s(x)}, 1\right) = \frac{\frac{1}{C} \cdot \min(p_t(x), M \cdot p_s(x))}{\frac{M}{C} \cdot p_s(x)},$$

where  $C$  is normalization constant defined as

$$C = \int_{\text{Supp}(p_s)} \min(p_t(x), M \cdot p_s(x)) \, dx.$$

Notably, we have  $1 \geq C \geq 1 - \varepsilon$ , by giving  $p_t(x)$  as an upper bound of the integrand in the first inequality, and by keeping only the set of  $x$ 's such that  $p_t(x) \leq Mp_s(x)$  in the second inequality. We have that the function  $p'_t : x \mapsto \min(p_t(x), Mp_s(x))/C$  is a probability density satisfying  $R_\infty(p'_t \| p_s) \leq M/C$ . Then algorithm  $\mathcal{A}^{\text{real}}$  is a perfect rejection sampling algorithm with target density  $p'_t$  and source density  $p_s$ . The output density of  $\mathcal{B}^{\text{real}}$  is exactly  $p'_t$ , as explained in [Dev86, Chapter II.3] (see in particular [Dev86, Theorems 3.1 and 3.2]). Moreover, the probability that  $\mathcal{A}^{\text{real}}$  outputs nothing is

$$\mathcal{A}^{\text{real}}(\perp) = 1 - \frac{C}{M} \in \left[ \frac{M-1}{M}, \frac{M-1+\varepsilon}{M} \right],$$

and the density of  $\mathcal{A}^{\text{real}}$  is  $x \mapsto (1 - \mathcal{A}^{\text{real}}(\perp)) \cdot p'_t(x) = \min(p_t(x)/M, p_s(x))$ .

Let us then bound the statistical distance.

$$\begin{aligned} \Delta(\mathcal{A}^{\text{real}}, \mathcal{A}^{\text{ideal}}) &= \frac{1}{2} \int_S \left| \frac{p_t(x)}{M} - \min\left(\frac{p_t(x)}{M}, p_s(x)\right) \right| dx + \frac{1}{2} \left| \mathcal{A}^{\text{real}}(\perp) - \frac{M-1}{M} \right| \\ &\leq \frac{1}{2} \int_S \left| \max\left(0, \frac{p_t(x)}{M} - p_s(x)\right) \right| dx + \frac{\varepsilon}{2M} \\ &\leq \frac{1}{2} \int_{\{x \in S \mid p_s(x) \leq p_t(x)/M\}} \left( \frac{p_t(x)}{M} - p_s(x) \right) dx + \frac{\varepsilon}{2M} \\ &\leq \frac{\varepsilon}{2M} + \frac{\varepsilon}{2M}, \end{aligned}$$

by assumption on  $p_s$  and  $p_t$ .

To bound the statistical distance between  $\mathcal{B}_\infty^{\text{real}}$  and  $\mathcal{B}_\infty^{\text{ideal}}$ , we first note that their distributions actually correspond to the distributions of  $\mathcal{A}^{\text{real}}$  and  $\mathcal{A}^{\text{ideal}}$  conditioned on the fact that they do not abort. We have

$$\begin{aligned} \Delta(\mathcal{B}_\infty^{\text{real}}, \mathcal{B}_\infty^{\text{ideal}}) &= \frac{1}{2} \int_{\text{Supp}(p_t)} \left| p_t(x) - \frac{1}{1 - \mathcal{A}^{\text{real}}(\perp)} \min\left(\frac{p_t(x)}{M}, p_s(x)\right) \right| dx \\ &= \frac{1}{a} \int_{\text{Supp}(p_t)} \left| (1 - \mathcal{A}^{\text{real}}(\perp)) p_t(x) - \min\left(\frac{p_t(x)}{M}, p_s(x)\right) \right| dx \\ &= \frac{1}{a} \left[ \int_{\substack{\text{Supp}(p_t) \\ p_s(x) \geq p_t(x)/M}} \left| 1 - \mathcal{A}^{\text{real}}(\perp) - \frac{1}{M} \right| p_t(x) \, dx \right. \\ &\quad \left. + \int_{\substack{\text{Supp}(p_t) \\ p_s(x) < p_t(x)/M}} \left| (1 - \mathcal{A}^{\text{real}}(\perp)) p_t(x) - p_s(x) \right| dx \right], \end{aligned}$$

where  $a = 2(1 - \mathcal{A}^{\text{real}}(\perp))$ .

Recalling the upper and lower bounds on  $\mathcal{A}^{\text{real}}(\perp)$ , the first integral can be bounded as follows:

$$\int_{\substack{x \in \text{Supp}(p_t) \\ p_s(x) \geq p_t(x)/M}} \left| 1 - \mathcal{A}^{\text{real}}(\perp) - \frac{1}{M} p_t(x) \right| dx \leq \frac{\varepsilon}{M} \int_{\substack{x \in \text{Supp}(p_t) \\ p_s(x) \geq p_t(x)/M}} p_t(x) dx.$$

We now observe that:

$$1 - \mathcal{A}^{\text{real}}(\perp) \geq \frac{\int_{\{x \in \text{Supp}(P_t) | P_s(x) \geq P_t(x)/M\}} p_t(x) dx}{M}.$$

Then, when we multiply the left integral by  $1/a$ , we obtain:

$$\frac{1}{a} \int_{\substack{x \in \text{Supp}(p_t) \\ p_s(x) \geq p_t(x)/M}} \left| 1 - \mathcal{A}^{\text{real}}(\perp) - \frac{1}{M} p_t(x) \right| dx \leq \frac{\varepsilon}{2}.$$

Next, we study the right integral. Note that since  $\varepsilon \leq 1/2$ , it holds that

$$0 \leq p_s(x) \leq \frac{p_t(x)}{M} \leq 2(1 - \mathcal{A}^{\text{real}}(\perp))p_t(x),$$

as  $1 - \mathcal{A}^{\text{real}}(\perp) \geq (1 - \varepsilon)/M \geq 1/(2M)$ . Hence the right integral satisfies

$$\int_{\substack{x \in \text{Supp}(p_t) \\ p_s(x) < p_t(x)/M}} |(1 - \mathcal{A}^{\text{real}}(\perp))p_t(x) - p_s(x)| dx \leq (1 - \mathcal{A}^{\text{real}}(\perp))\varepsilon.$$

Finally, when divided by  $2(1 - \mathcal{A}^{\text{real}}(\perp))$ , we get  $\varepsilon/2$  as an upper bound. This provides the result.  $\square$

## B.2 Proof of Lemma 3.2

*Proof.* Let us first take care of the  $m = 1$  case. We define  $g^* : x \mapsto (g(x) + g(-x))/2$  as well as  $f^* : x \mapsto (f(x) + f(-x))/2$ . First, for any  $x \in \mathbb{R}$  and  $v \in [-t, t]$ , we have  $f(x) \leq M \cdot g(x - v)$  as well as  $f(-x) \leq M \cdot g(-x + v)$ . This implies that  $R_\infty(f^* \| g^*) \leq M$ . Now, by construction, these two functions are even. Moreover, they are normalized and are thus probability densities. Finally, we have  $\mathbb{E}_{x \leftarrow f^*}(|x|) = (\mathbb{E}_{x \leftarrow f}(|x|) + \mathbb{E}_{x \leftarrow f}(|-x|))/2 = \mathbb{E}_{x \leftarrow f}(|x|)$ .

In the following, we assume that  $m \geq 2$ . To define  $f^*$  and  $g^*$ , we will switch from Cartesian to hyperpsheral coordinates. This is done the following way. Let  $(x_1, \dots, x_m)$  and  $(\rho, \theta_1, \dots, \theta_{m-1})$  both representing  $\mathbf{x}$  in respectively Carte-

sian and hyperspherical coordinates. They satisfy the relations

$$\begin{aligned}\|\mathbf{x}\| &= \rho \\ x_1 &= \rho \cos(\theta_1) \\ x_2 &= \rho \sin(\theta_1) \cos(\theta_2) \\ &\vdots \\ x_{m-1} &= \rho \left( \prod_{i \leq m-2} \sin(\theta_i) \right) \cos(\theta_{m-1}) \\ x_m &= \rho \left( \prod_{i \leq m-1} \sin(\theta_i) \right).\end{aligned}$$

Let  $\vec{\theta} = (\theta_1, \dots, \theta_{m-1})$  and  $\mathbf{x}(\rho, \vec{\theta})$  be the vector whose coordinates are defined as above. Notice that the absolute value of the determinant of the variable change Jacobian is of the form  $\rho^{m-1} D(\vec{\theta})$  for some  $D : [0, \pi]^{m-2} \times [0, 2\pi] \rightarrow \mathbb{R}_{\geq 0}$ , as all columns except the first one are of the form  $\rho \cdot \mathbf{y}_i(\vec{\theta})$ , and the first column does not depend on  $\rho$ . It then holds that

$$1 = \int_{\mathbb{R}^m} f(\mathbf{x}) \, d\mathbf{x} = \int_0^\infty \rho^{m-1} \int_{[0, \pi]^{m-2} \times [0, 2\pi]} f(\mathbf{x}(\rho, \vec{\theta})) D(\vec{\theta}) \, d\vec{\theta} \, d\rho.$$

We then define:

$$f^* : \mathbf{z} \mapsto \frac{\int_{[0, \pi]^{m-2} \times [0, 2\pi]} f(\mathbf{x}(\|\mathbf{z}\|, \vec{\theta})) D(\vec{\theta}) \, d\vec{\theta}}{\int_{[0, \pi]^{m-2} \times [0, 2\pi]} D(\vec{\theta}) \, d\vec{\theta}}.$$

This is a probability density, as it is integrable and  $\int_{\mathbb{R}^m} f^*(\mathbf{x}) \, d\mathbf{x} = 1$ . The latter can be seen by switching once more to hyperspherical coordinates. By construction, it is isotropic. We also have  $\mathbb{E}_{\mathbf{z} \leftarrow f}(\|\mathbf{z}\|) = \mathbb{E}_{\mathbf{z} \leftarrow f^*}(\|\mathbf{z}\|)$ , which is also seen by applying the same change of variables. We define  $g^*$  in the same way. It remains to prove that  $\sup_{\mathbf{v} \in V} R_\infty(f^* \|g_{+\mathbf{v}}^*) \leq M$ .

Let  $\mathbf{z} \in \mathbb{R}^m$  and  $\mathbf{v} \in V$ . Let  $\lambda = \|\mathbf{z} - \mathbf{v}\|/\|\mathbf{z}\|$ . We consider the scaling  $s : \mathbf{y} \mapsto \lambda \cdot \mathbf{y}$ . For any  $\mathbf{y} \in \mathcal{S}_m(\|\mathbf{z}\|)$  (i.e., any  $\mathbf{y} \in \mathbb{R}^m$  with  $\|\mathbf{y}\| = \|\mathbf{z}\|$ ), we have  $f(\mathbf{y}) \leq M \cdot g(s(\mathbf{y}))$  as  $s(\mathbf{y})$  can be written as  $\mathbf{y} - \mathbf{v}'$ , where  $\mathbf{v}' \in V$ . Indeed, we have, using the triangle inequality:

$$\|s(\mathbf{y}) - \mathbf{y}\| = |\lambda - 1| \|\mathbf{z}\| = \|\|\mathbf{z} - \mathbf{v}\| - \|\mathbf{z}\|\| \leq \|\mathbf{v}\| \leq t.$$

Decomposing every element  $\mathbf{y} \in \mathcal{S}_m(\|\mathbf{z}\|)$  in hyperspherical coordinates as above with  $\rho = \|\mathbf{z}\|$  for unique  $(\theta_1, \dots, \theta_{m-1}) \in [0, \pi]^{m-2} \times [0, 2\pi]$ , we multiply both sides by  $D(\vec{\theta})$ , which is nonnegative, and we integrate over  $[0, \pi]^{m-2} \times [0, 2\pi]$  to get:

$$\int_{[0, \pi]^{m-2} \times [0, 2\pi]} f(\mathbf{x}(\|\mathbf{z}\|, \vec{\theta})) D(\vec{\theta}) \, d\vec{\theta} \leq M \int_{[0, \pi]^{m-2} \times [0, 2\pi]} g(\lambda \mathbf{x}(\|\mathbf{z}\|, \vec{\theta})) D(\vec{\theta}) \, d\vec{\theta}.$$

Recalling the definition of  $\lambda$  and dividing both sides by  $\int_{[0, \pi]^{m-2} \times [0, 2\pi]} D(\vec{\theta}) \, d\vec{\theta}$ , we obtain the result.  $\square$

### B.3 Proof of Theorem 3.4

As in the unimodal case, we first radialize the densities.

**Lemma B.1.** *Let  $m \geq 1, t > 0$  and  $V = \mathcal{B}_m(t)$ . Let  $f, g : \mathbb{R}^m \rightarrow [0, 1]$  be two probability densities over  $\mathbb{R}^m$  and define  $M = \sup_{\mathbf{v} \in V} R_\infty(f \| g_{\pm \mathbf{v}})$ , where  $g_{\pm \mathbf{v}}$  is as in Theorem 3.4. Then there exist two probability densities  $f^*, g^*$  that satisfy*

- $\sup_{\mathbf{v} \in V} R_\infty(f^* \| g_{\pm \mathbf{v}}^*) \leq M$ ,
- $\forall \mathbf{x}, \mathbf{y} \in \mathbb{R}^m, \|\mathbf{x}\| = \|\mathbf{y}\| \implies f^*(\mathbf{x}) = f^*(\mathbf{y})$  and  $g^*(\mathbf{x}) = g^*(\mathbf{y})$ ,
- $\mathbb{E}_{\mathbf{z} \leftarrow f}(\|\mathbf{z}\|) = \mathbb{E}_{\mathbf{z} \leftarrow f^*}(\|\mathbf{z}\|)$ .

*Proof.* We proceed as in the proof of Lemma 3.2 to define  $f^*$  and  $g^*$ , and use the same notations. Let

$$f^* : \mathbf{z} \mapsto \frac{\int_{[0, \pi]^{m-2} \times [0, 2\pi]} f(\mathbf{x}(\|\mathbf{z}\|, \vec{\theta})) D(\vec{\theta}) d\vec{\theta}}{\int_{[0, \pi]^{m-2} \times [0, 2\pi]} D(\vec{\theta}) d\vec{\theta}}.$$

We define  $g^*$  in the same way, with  $f$  replaced by  $g$ . As shown for Lemma 3.2, the first two claims hold.

Let  $\mathbf{z} \in \mathbb{R}^m$  and  $\mathbf{v} \in V$ . For any  $\mathbf{y} \in \mathcal{S}_m(\|\mathbf{z}\|)$ , let  $s_{\mathbf{y}}$  be the rotation that maps  $\mathbf{z}$  to  $\mathbf{y}$ . It is an isometry, so for any  $b \in \{0, 1\}$ , it holds  $\|\mathbf{y} + (-1)^b s_{\mathbf{y}}(\mathbf{v})\| = \|\mathbf{z} + (-1)^b s_{\mathbf{y}}(\mathbf{v})\|$ . We also have  $\|s_{\mathbf{y}}(\mathbf{v})\| = \|\mathbf{v}\|$ , implying that  $s_{\mathbf{y}}(\mathbf{v}) \in V$ . It then holds for any  $\mathbf{y} \in \mathcal{S}_m(\|\mathbf{z}\|)$ :

$$f(s_{\mathbf{y}}(\mathbf{z})) \leq \frac{M}{2} (g(s_{\mathbf{y}}(\mathbf{z} - \mathbf{v})) + g(s_{\mathbf{y}}(\mathbf{z} + \mathbf{v}))).$$

By construction, there exist  $\vec{\theta}_0$  such that  $s_{\mathbf{x}(\|\mathbf{z}\|, \vec{\theta}_0)}(\mathbf{z}) = \mathbf{x}(\|\mathbf{z}\|, \vec{\theta}_0)$ . This lets us multiply both sides with  $D(\vec{\theta}_0)$ , which is nonnegative, and integrate over  $[0, \pi]^{m-2} \times [0, 2\pi]$ . It yields the first claim, up to dividing by the normalisation constant.  $\square$

*Proof (Theorem 3.4).* Thanks to Lemma B.1, we assume without loss of generality that both  $f$  and  $g$  are isotropic. Let us define the moments of  $f, g$  by  $\mu_k^{(\phi)} := \int_0^\infty x^k \cdot \phi(x) dx$  for  $k \geq 0$  and  $\phi \in \{f, g\}$ . We have  $\mu_m^{(\phi)} = \mathbb{E}_{\mathbf{x} \leftarrow \phi}(\|\mathbf{x}\|) / S_m$  and  $\mu_{m-1}^{(\phi)} = 1 / S_m$  as noted in the proof of Theorem 3.1. For any  $r \geq 0, u \in [0, t]$  and  $\theta \in [0, 2\pi)$ , it holds that:

$$f(r) \leq \frac{M}{2} \left( g\left(\sqrt{r^2 + u^2 - 2ru \cos(\theta)}\right) + g\left(\sqrt{r^2 + u^2 + 2ru \cos(\theta)}\right) \right).$$

We will only consider  $\theta = \pi/2$  as it will suffice to obtain the bound. This gives for any  $r \in \mathbb{R}_{\geq 0}$  and  $u \in [0, t]$ :

$$f(r) \leq Mg\left(\sqrt{r^2 + u^2}\right).$$

Let us then multiply both sides by  $r(\sqrt{r^2+t^2})^{m-2}$  and integrate over  $\mathbb{R}_{\geq 0}$ . On the right-hand side, we use the change of variable  $y = \sqrt{r^2+t^2}$ , which yields  $dy = r/\sqrt{r^2+t^2}dr$ , to obtain:

$$\int_0^\infty r(\sqrt{r^2+t^2})^{m-2} f(r) dr \leq M \cdot \int_t^\infty y^{m-1} g(y) dy.$$

Since  $y \mapsto y^{m-1}g(y)$  takes values in  $\mathbb{R}_{\geq 0}$ , we have that  $M/S_m = M\mu_{m-1}^{(g)}$  is an upper bound for the right-hand side. By reordering terms, we obtain:

$$0 \leq \int_0^\infty r \left[ \left( M^{\frac{2}{m-2}} r^2 \right)^{\frac{m-2}{2}} - (r^2+t^2)^{\frac{m-2}{2}} \right] f(r) dr.$$

Now, note that for  $m \geq 4$ , we have:

$$\begin{aligned} & \left( M^{\frac{2}{m-2}} r^2 \right)^{\frac{m-2}{2}} - (r^2+t^2)^{\frac{m-2}{2}} \\ &= \frac{M^{\frac{2}{m-2}} r^2 - r^2 - t^2}{M^{\frac{1}{m-1}} r + \sqrt{r^2+t^2}} \sum_{k=0}^{m-3} \left( M^{\frac{2}{m-2}} r^2 \right)^{\frac{k}{2}} (r^2+t^2)^{\frac{m-3-k}{2}}. \end{aligned}$$

This also holds for  $m = 3$  if replacing the sum by 1. Note that for  $r \geq 0$ , we have  $M^{\frac{1}{m-1}} r + \sqrt{r^2+t^2} \geq t$ . Let  $C = t/(M^{\frac{2}{m-2}} - 1)^{1/2}$ . The inequality  $r^2+t^2 \leq M^{\frac{2}{m-2}} r^2$  holds if and only if  $r \geq C$ . Then for  $r \geq 0$  and  $m \geq 4$ , we have

$$\left( M^{\frac{2}{m-2}} r^2 \right)^{\frac{m-2}{2}} - (r^2+t^2)^{\frac{m-2}{2}} \leq (M^{\frac{2}{m-2}} - 1)(r^2 - C^2) \frac{m-2}{t} \cdot M^{\frac{m-3}{m-2}} r^{m-3}.$$

Since all constants are positive, we obtain (including for  $m \geq 3$ ):

$$0 \leq \int_0^\infty (r^m - C^2 r^{m-2}) f(r) dr.$$

Equivalently, we have that  $\mu_m^{(f)} \geq C^2 \mu_{m-2}^{(f)}$ , which we rewrite this as  $(\mu_m^{(f)}/\mu_{m-1}^{(f)}) \cdot (\mu_{m-1}^{(f)}/\mu_{m-2}^{(f)}) \geq C^2$ . As we have seen in the proof of Theorem 3.1, the Cauchy-Schwarz inequality implies that  $\mu_{m-1}^{(f)}/\mu_{m-2}^{(f)} \leq \mu_m^{(f)}/\mu_{m-1}^{(f)}$ . This leads to the desired lower bound.  $\square$

#### B.4 Proof of Lemma 5.3

*Proof.* In this proof, we omit the  $U(\mathcal{B}_m(r))$  subscripts for the min-entropies. Note that  $H_\infty(\lceil \mathbf{x}_0 \rceil | \lceil \mathbf{x}_1 \rceil) \geq H_\infty(\lceil \mathbf{x}_0 \rceil | \mathbf{x}_1)$ . By definition of the conditional min-entropy, we have

$$2^{-H_\infty(\lceil \mathbf{x}_0 \rceil | \mathbf{x}_1)} = \int_{\mathbf{x}_1 \in \mathcal{B}_{m-n}(r)} \max_{\mathbf{x}_0^{\text{int}} \in \mathbb{Z}^n} \left( \int_{\mathbf{x}_0 \in \mathcal{B}_n^\infty(1/2, \mathbf{x}_0^{\text{int}})} p_{\mathbf{x}_0, \mathbf{x}_1}(\mathbf{x}_0, \mathbf{x}_1) d\mu(\mathbf{x}_0) \right) d\mu(\mathbf{x}_1),$$

where the density satisfies

$$p_{(\mathbf{x}_0, \mathbf{x}_1)}(\mathbf{x}_0, \mathbf{x}_1) = \frac{1}{V_m(r)} \chi_{<r^2}(\|\mathbf{x}_0\|^2 + \|\mathbf{x}_1\|^2).$$

Recall that  $\chi_{<r^2}(y) = 1$  if  $y \leq r^2$  and 0 otherwise and that  $V_m(r)$  is the volume of the Euclidean ball of radius  $r$  in dimension  $m$ .

The maximum is achieved when  $\mathbf{x}_0^{\text{int}} = \mathbf{0}$ . Indeed, for any  $\mathbf{x}_0^{\text{int}} \in \mathbb{Z}^m$ , we have

$$\begin{aligned} & \int_{\mathbf{x}_0 \in \mathcal{B}_n^\infty(1/2, \mathbf{x}_0^{\text{int}})} \chi_{<r^2}(\|\mathbf{x}_0\|^2 + \|\mathbf{x}_1\|^2) d\mu(\mathbf{x}_0) \\ &= \int_{\mathbf{x}_0 \in \mathcal{B}_n^\infty(1/2, 0)} \chi_{<r^2}(\|\mathbf{x}_0 + \mathbf{x}_0^{\text{int}}\|^2 + \|\mathbf{x}_1\|^2) d\mu(\mathbf{x}_0) \\ &\leq \int_{\mathbf{x}_0 \in \mathcal{B}_n^\infty(1/2, 0)} \chi_{<r^2}(\|\mathbf{x}_0\|^2 + \|\mathbf{x}_1\|^2) d\mu(\mathbf{x}_0), \end{aligned}$$

where we used the fact that if  $\|\mathbf{x}_0\|_\infty \leq \frac{1}{2}$  and  $\mathbf{x}_0^{\text{int}} \in \mathbb{Z}^n$ , then we have  $\|\mathbf{x}_0 + \mathbf{x}_0^{\text{int}}\| \geq \|\mathbf{x}_0\|$ . As a result, we can write

$$2^{-H_\infty(\lceil \mathbf{x}_0 \rceil | \mathbf{x}_1)} = \Pr \left[ \|\mathbf{x}_0\|_\infty \leq \frac{1}{2} \right].$$

Now we use the sub-independence of the coordinates slabs in the Euclidean ball [BP98], i.e., denoting  $\mathbf{x}_0 = (x_{01}, x_{02}, \dots, x_{0n})^T$ , we have

$$\Pr \left[ \|\mathbf{x}_0\|_\infty \leq \frac{1}{2} \right] \leq \prod_{i=1}^n \Pr \left[ |x_{0i}| \leq \frac{1}{2} \right].$$

We now use Lemma B.2 below to get

$$\begin{aligned} \Pr \left[ |x_{01}| \leq \frac{1}{2} \right] &= \Pr \left[ \frac{|x_{01}|}{r} \leq \frac{1}{2r} \right] \\ &\leq \Pr \left[ \frac{|x_{01}|}{r} \leq \frac{1}{4\sqrt{m}} \right] \\ &\leq 0.843 + 2 \exp(-m) \leq 0.85, \end{aligned}$$

for  $r \geq 2\sqrt{m}$  and  $m \geq 6$ . □

**Lemma B.2.** *Let  $(x_1, \dots, x_m)^T$  be uniformly chosen in the  $m$ -dimensional Euclidean ball of radius 1. Then*

$$\Pr \left[ |x_1| \leq \frac{1}{4\sqrt{m}} \right] \leq 0.843 + 2 \exp(-m).$$

*Proof.* To show this, we use the fact (see [BGMN05]) that we can obtain samples  $(x_1, \dots, x_m)^T$  by first sampling  $m$  independent Gaussian variables  $g_1, \dots, g_m$  with densities  $t \mapsto \rho_{1/\sqrt{2}}(t)/\sqrt{\pi}$  and then setting  $(x_1, \dots, x_m)^T = \frac{(g_1, \dots, g_m)^T}{\sqrt{\sum_{i=1}^m g_i^2 + z}}$ ,

where  $z$  is independent and has an exponential distribution (i.e., density  $t \mapsto \exp(-t)$ ). With this notation, we have, for all  $\delta > 0$ :

$$\begin{aligned}
\Pr \left[ x_1^2 > \frac{\delta^2}{m} \right] &= \Pr \left[ \frac{g_1^2}{\sum_{i \leq m} g_i^2 + z} > \frac{\delta^2}{m} \right] \\
&\geq \Pr \left[ g_1^2 > 4\delta^2 \text{ and } \sum_{i \leq m} g_i^2 \leq 3m \text{ and } z \leq m \right] \\
&\geq \Pr [|g_1| > 2\delta] - \Pr \left[ \sum_{i \leq m} g_i^2 > 3m \right] - \Pr [z > m] \\
&\geq \Pr [|g_1| > 2\delta] - \exp(-m) - \exp(-m) .
\end{aligned}$$

For the last inequality, note that the distribution of  $2 \sum_{i \leq m} g_i^2$  is the chi-squared distribution of parameter  $m$ . If  $F$  denotes its cumulative density function, then we have the tail bound  $1 - F(x) \leq ((x/m) \exp(1 - x/m))^{m/2}$  for  $x > m$ , which we use with  $x = 6m$ . Taking  $\delta = 1/2$  and numerically evaluating the first term allows to complete the proof of the lemma.  $\square$

## C Complements to Section 5

In this appendix, we bound the (smooth) Rényi divergence for distributions classically used in Lyubashevsky's signatures. This lets us compare these choices with our choice of continuous uniform distributions over hyperballs. We also provide a more detailed description of the adaptation of Lyubashevsky's signature to continuous distributions.

### C.1 Uniform Distribution in Hypercubes

For simplicity and ease of implementation, some applications rely on uniform distributions in hypercubes. The following result is implicit in [Lyu09].

**Lemma C.1 (Rényi Divergence).** *Let  $m \geq 1$  and  $\mathbf{v} \in \mathbb{Z}^m$ . Let  $r, r' \geq 1/2$  such that  $r' \geq r + \|\mathbf{v}\|_\infty$ . Then it holds that*

$$R_\infty \left( U(\mathcal{B}_m^\infty(r) \cap \mathbb{Z}^m) \| U(\mathcal{B}_m^\infty(r', \mathbf{v}) \cap \mathbb{Z}^m) \right) \leq \left( \frac{2r' + 1}{2r - 1} \right)^m .$$

Let  $M > 1$ . The above is  $\leq M$  if  $r \geq \frac{\|\mathbf{v}\|_\infty + (M^{1/m} + 1)/2}{M^{1/m} - 1}$  and  $r' = r + \|\mathbf{v}\|_\infty$ .

*Proof.* For the divergence to be defined, we need  $\mathcal{B}_m^\infty(r) \cap \mathbb{Z}^m \subseteq \mathcal{B}_m^\infty(r', \mathbf{v}) \cap \mathbb{Z}^m$ . This is ensured by the constraint  $r' \geq r + \|\mathbf{v}\|_\infty$ . In that case, the divergence is the ratio of the number of elements in each support, leading to the upper bound. The second claim follows by elementary calculations.  $\square$

The downside of the infinite norm is its lack of geometry: as we detail later, the use of the scalar product induced by the Euclidean norm is crucial to improve the bounds for the smooth divergence and the divergence with a bimodal version of the distribution, both for Gaussian distributions and uniforms in hyperballs. Oppositely, the smooth divergence and considering a bimodal version of the uniform distribution in a hypercube do not bring significant improvements to the radius condition from Lemma C.1.

## C.2 Gaussian Distributions

By Lemma A.15 and the fact that  $\lim_{a \rightarrow +\infty} R_a(P\|Q) = R_\infty(P\|Q)$ , we see that the Rényi divergence of infinite order between two Gaussian distributions with same standard deviation but different centers is infinite. Using Lemma A.7, we are however able to obtain a finite upper bound on the smooth divergence. The result is of the same flavour as [Lyu12, Lemma 4.5], but the proof significantly differs and leads to a smaller upper bound (in [Lyu12, Lemma 4.5], the quantity  $\|\mathbf{v}\|^2$  is divided by  $4\sigma^2$  instead of  $2\sigma^2$  here).

**Lemma C.2 (Smooth Rényi Divergence).** *Let  $m > 0, \mathbf{v} \in \mathbb{R}^m, \varepsilon \in (0, 1)$  and  $\sigma > 0$ . We have:*

$$R_\infty^\varepsilon(D_{\mathbb{Z}^m, \sigma} \| D_{\mathbb{Z}^m, \sigma, \mathbf{v}}) \leq \exp\left(\frac{\|\mathbf{v}\|^2}{2\sigma^2} + \frac{\|\mathbf{v}\| \sqrt{2 \log \frac{1}{\varepsilon}}}{\sigma}\right),$$

Let  $M > 1$ . The above is  $\leq M$  if

$$\sigma \geq \frac{\|\mathbf{v}\|}{\sqrt{2 \log(M)}} \left( \sqrt{\log \frac{1}{\varepsilon}} + \sqrt{\log \frac{1}{\varepsilon} + \log M} \right).$$

*Proof.* Combining Lemmas A.15 and A.7, we obtain that for any  $a \in (1, +\infty)$ :

$$R_\infty^\varepsilon(P\|Q) \leq \exp\left(\frac{a\|\mathbf{v}\|^2}{2\sigma^2} + \frac{1}{a-1} \log \frac{1}{\varepsilon}\right).$$

Let us instantiate this for  $a = 1 + \frac{\sigma}{\|\mathbf{v}\|} \sqrt{2 \log \frac{1}{\varepsilon}}$ , which minimizes the above quantity. This gives us the bound

$$R_\infty^\varepsilon(P\|Q) \leq \exp\left(\frac{\|\mathbf{v}\|^2}{2\sigma^2} + \sqrt{2} \frac{\|\mathbf{v}\|}{\sigma} \sqrt{\log \frac{1}{\varepsilon}}\right).$$

To find when this is  $\leq M$ , we take the logarithm and multiply by  $\sigma^2$  on both sides. Solving a degree-2 equation in  $\sigma$  leads to the second claim.  $\square$

The following is borrowed from [DDLL13], and its proof is provided here for the sake of completeness.

**Lemma C.3 (Rényi Divergence with a Bimodal Gaussian).** *Let  $m \geq 1$ ,  $\mathbf{v} \in \mathbb{R}^m$  and  $\sigma > 0$ . Then the following holds:*

$$R_\infty(D_{\mathbb{Z}^m, \sigma} \| BD_{\mathbb{Z}^m, \sigma, \mathbf{v}}) \leq \exp\left(\frac{\|\mathbf{v}\|^2}{2\sigma^2}\right).$$

*It is an equality if  $\mathbf{v} \in \mathbb{Z}^m$ . Let  $M \geq 1$ . The bound is  $\leq M$  if  $\sigma \geq \|\mathbf{v}\|/(2\sqrt{\log M})$ .*

*Proof.* Let  $\mathbf{z} \in \mathbb{Z}^m$ . We have:

$$\begin{aligned} \frac{D_{\mathbb{Z}^m, \sigma}(\mathbf{z})}{BD_{\mathbb{Z}^m, \sigma, \mathbf{v}}(\mathbf{z})} &= \frac{\rho_{\sigma, \mathbf{v}}(\mathbb{Z}^m)}{\rho_\sigma(\mathbb{Z}^m)} \cdot \frac{\exp\left(\frac{-\|\mathbf{z}\|^2}{2\sigma^2}\right)}{\exp\left(\frac{-\|\mathbf{z}\|^2 - \|\mathbf{v}\|^2}{2\sigma^2}\right) \cosh\left(\frac{|\langle \mathbf{z}, \mathbf{v} \rangle|}{\sigma^2}\right)} \\ &= \frac{\rho_{\sigma, \mathbf{v}}(\mathbb{Z}^m)}{\rho_\sigma(\mathbb{Z}^m)} \cdot \frac{\exp\left(\frac{\|\mathbf{v}\|^2}{2\sigma^2}\right)}{\cosh\left(\frac{|\langle \mathbf{z}, \mathbf{v} \rangle|}{\sigma^2}\right)} \\ &\leq \frac{\rho_{\sigma, \mathbf{v}}(\mathbb{Z}^m)}{\rho_\sigma(\mathbb{Z}^m)} \cdot \exp\left(\frac{\|\mathbf{v}\|^2}{2\sigma^2}\right), \end{aligned}$$

where the last inequality comes from the fact that  $\cosh(x) \geq 1$  for any  $x \in \mathbb{R}$ . Note that for  $\mathbf{z} \in \mathbb{Z}^m$  orthogonal to  $\mathbf{v}$ , this upper bound is reached. Finally, using [MR07, Lemma 2.9], we have that  $\frac{\rho_{\sigma, \mathbf{v}}(\mathbb{Z}^m)}{\rho_\sigma(\mathbb{Z}^m)} \leq 1$ . If  $\mathbf{v} \in \mathbb{Z}^m$  this is actually an equality.  $\square$

As a side note, we observe that this result can be extended to any order and compared to standard results between two Gaussian distributions.

**Corollary C.4.** *Let  $m \geq 1$ ,  $\mathbf{v} \in \mathbb{R}^m$  and  $\sigma > 0$ . Then the following holds:*

$$\forall a \in [1, +\infty], R_a(D_{\mathbb{Z}^m, \sigma} \| BD_{\mathbb{Z}^m, \sigma, \mathbf{v}}) \leq \exp\left(\frac{\|\mathbf{v}\|^2}{2\sigma^2}\right) = (R_a(D_{\mathbb{Z}^m, \sigma} \| D_{\mathbb{Z}^m, \sigma, \mathbf{v}}))^{\frac{1}{a}}.$$

*Proof.* The Rényi divergence is increasing in its order. Thus the upper bound from Lemma C.3 is also an upper bound for any order  $a \in [1, +\infty]$ .  $\square$

### C.3 Lyubashevsky's Signature with Continuous Distributions

We use the same notations as in Section 2.2, with the source density  $g$  and the target density  $f$  being with supports over  $\mathbb{R}^m$  rather than  $\mathbb{Z}^m$ . The modified signature scheme handling continuous source and target distributions is presented in Figure 7. The key generation algorithm is unchanged from Figure 2. Note that if  $f$  and  $g$  were actually discrete densities, then we would exactly recover the scheme from Figure 2.

For correctness, note that

$$\mathbf{A}[\mathbf{z}] - \mathbf{Tc} = \mathbf{A}[\mathbf{y} + \mathbf{Sc}] - \mathbf{Tc} = \mathbf{A}([\mathbf{y}] + \mathbf{Sc}) - \mathbf{Tc} = \mathbf{A}[\mathbf{y}],$$

<b>Sign(<math>\mu, \mathbf{A}, \mathbf{S}</math>) :</b> 1: $\mathbf{y} \leftarrow g$ 2: $\mathbf{c} \leftarrow H(\mathbf{A}[\mathbf{y}], \mu)$ 3: $\mathbf{z} \leftarrow \mathbf{y} + \mathbf{S}\mathbf{c}$ 4: $u \leftarrow U([0, 1])$ 5: <b>if</b> $u \leq \min\left(\frac{f(\mathbf{z})}{M \cdot g(\mathbf{y})}, 1\right)$ <b>then</b> 6:     return $(\lceil \mathbf{z} \rceil, \mathbf{c})$ 7: <b>else</b> 8:     go to Step 1 9: <b>end if</b>	<b>Verify(<math>\mu, \mathbf{z}, \mathbf{c}, \mathbf{A}, \mathbf{T} = \mathbf{A}\mathbf{S}</math>) :</b> 1: <b>if</b> $\ \mathbf{z}\  \leq \gamma$ and $\mathbf{c} = H(\mathbf{A}\mathbf{z} - \mathbf{T}\mathbf{c}, \mu)$ <b>then</b> 2:     return 1 3: <b>else</b> 4:     return 0 5: <b>end if</b>
---	---

**Fig. 7.** Lyubashevsky's signature scheme with continuous distributions.

where the second equality holds because  $\mathbf{S}\mathbf{c}$  is an integer vector.

The main modification towards analyzing the security of the scheme from Figure 7, compared to the one from Figure 2, resides in the observation that the distributions of the pair  $(\mathbf{z}, \mathbf{c})$  obtained by the two processes from Figure 8 have statistical distance  $\leq \varepsilon/M$ , where  $\varepsilon$  is such that  $R_\infty^\varepsilon(f||g) \leq M$ . This is obtained by Lemma 2.2 and the (statistical distance) data processing inequality.

The only other analysis modification is related to the min-entropy of the commitments: the condition  $H_\infty(\mathbf{y}_0|\mathbf{y}_1)_Q = \Omega(\lambda)$  appearing in Section 2.2 now becomes  $H_\infty(\lceil \mathbf{y}_0 \rceil || \lceil \mathbf{y}_1 \rceil)_g = \Omega(\lambda)$ .

1: $\mathbf{y} \leftarrow g$ 2: $\mathbf{c} \leftarrow U(\mathcal{C})$ 3: $\mathbf{z} \leftarrow \mathbf{y} + \mathbf{S}\mathbf{c}$ 4: $u \leftarrow U([0, 1])$ 5: <b>if</b> $u \leq \min\left(\frac{f(\mathbf{z})}{M \cdot g(\mathbf{y})}, 1\right)$ <b>then</b> 6:     return $(\lceil \mathbf{z} \rceil, \mathbf{c})$ 7: <b>else</b> 8:     return $(\perp, \perp)$ 9: <b>end if</b>	1: $\mathbf{c} \leftarrow U(\mathcal{C})$ 2: $\mathbf{z} \leftarrow f$ 3: $u \leftarrow U([0, 1])$ 4: <b>if</b> $u \leq \frac{1}{M}$ <b>then</b> 5:     return $(\lceil \mathbf{z} \rceil, \mathbf{c})$ 6: <b>else</b> 7:     return $(\perp, \perp)$ 8: <b>end if</b>
---	---

**Fig. 8.** Simulating signatures.

## D Another Rejection Sampling Algorithm

In this section, we study the rejection sampling procedure described in [HJMR07] and how to apply it to the Fiat-Shamir with aborts framework.

Let  $m > 0$  and  $V \subset \mathbb{Z}^m$  as well as  $P$  and  $Q$  two probability distributions over  $\mathbb{Z}^m$  such that  $\sup_{\mathbf{v} \in V} \log(R_1(P||Q_{+\mathbf{v}})) < \infty$ . Let  $P_H$  be a distribution

over  $V$ . We first define  $P_t : (\mathbf{z}, \mathbf{v}) \mapsto P_H(\mathbf{v})P(\mathbf{z})$  and  $P_s : (\mathbf{z}, \mathbf{v}) \mapsto P_H(\mathbf{v})Q(\mathbf{z} - \mathbf{v})$ . Let  $p_0(\mathbf{z}, \mathbf{v}) = 0$  for any  $(\mathbf{z}, \mathbf{v}) \in \mathbb{Z}^m \times V$  and recursively define the following:

$$\begin{cases} \alpha_i(\mathbf{z}, \mathbf{v}) &= \min(P_t(\mathbf{z}, \mathbf{v}) - p_{i-1}(\mathbf{z}, \mathbf{v}), (1 - p_{i-1}^*)P_s(\mathbf{z}, \mathbf{v})), \\ p_i(\mathbf{z}, \mathbf{v}) &= p_{i-1}(\mathbf{z}, \mathbf{v}) + \alpha_i(\mathbf{z}, \mathbf{v}), \\ p_i^* &= \sum_{(\mathbf{z}, \mathbf{v}) \in \mathbb{Z}^m \times V} p_i(\mathbf{z}, \mathbf{v}). \end{cases}$$

Finally, define  $\beta_i(\mathbf{z}, \mathbf{v}) = \min\left(\frac{P_t(\mathbf{z}, \mathbf{v}) - p_{i-1}(\mathbf{z}, \mathbf{v})}{(1 - p_{i-1}^*)P_s(\mathbf{z}, \mathbf{v})}, 1\right)$ .

$\mathcal{A}$	$\mathcal{A}'$
1: $i \leftarrow 1$ .	1: Sample $\mathbf{z} \leftarrow P$ .
2: Sample $\mathbf{y} \leftarrow Q$ .	2: Sample $\mathbf{v} \leftarrow P_H$ .
3: Sample $\mathbf{v} \leftarrow P_H$ .	3: Return $(\mathbf{z}, \mathbf{v})$ .
4: Set $\mathbf{z} \leftarrow \mathbf{y} + \mathbf{v}$ .	
5: Sample $u \leftarrow [0, 1]$ .	
6: <b>if</b> $u \leq \beta_i(\mathbf{z}, \mathbf{v})$ <b>then</b>	
7:     Return $(\mathbf{z}, \mathbf{v})$ .	
8: <b>else</b>	
9: $i \leftarrow i + 1$ .	
10:    Go to 2.	
11: <b>end if</b>	

**Fig. 9.** Greedy rejection sampling

**Lemma D.1 (Correctness).** *Let  $r(i, \mathbf{z}, \mathbf{v})$  be the probability that the procedure returns a pair  $(\mathbf{z}, \mathbf{v}) \in \mathbb{Z}^m \times V$  after exactly  $i$  iterations. Then for any  $\mathbf{z} \in \mathbb{Z}^m$  and  $\mathbf{v} \in V$  it holds that  $r(i, \mathbf{z}, \mathbf{v}) = \alpha_i(\mathbf{z}, \mathbf{v})$  and  $\sum_{j=1}^{\infty} r(j, \mathbf{z}, \mathbf{v}) = P_t(\mathbf{z}, \mathbf{v})$ . Put differently, the statistical distance between the distribution of the output of  $\mathcal{A}$  and  $\mathcal{A}'$  is 0.*

*Proof.* The probability that  $(i, \mathbf{z}, \mathbf{v})$  is output is exactly  $P_s(\mathbf{z}, \mathbf{v})\bar{p}_{i-1} \cdot \beta_i(\mathbf{z}, \mathbf{v})$ , where  $\bar{p}_{i-1}$  denotes the probability that the  $i - 1$  first values are rejected. We then show by induction that  $\bar{p}_i = 1 - p_i^*$  for any  $i \in \mathbb{N}$ . In the case  $i = 0$ , we have  $\bar{p}_0 = 1 = 1 - p_0^*$ .

Let us now assume that this holds for some  $i \in \mathbb{N}$ . By induction, let us compute  $\bar{p}_{i+1} = \bar{p}_i \cdot \sum_{(\mathbf{z}, \mathbf{v}) \in \mathbb{Z}^m \times V} (1 - \beta_{i+1}(\mathbf{z}, \mathbf{v}))P_s(\mathbf{z}, \mathbf{v})$ . We have:

$$\begin{aligned} \bar{p}_{i+1} &= (1 - p_i^*) \sum_{(\mathbf{z}, \mathbf{v}) \in \mathbb{Z}^m \times V} (1 - \beta_{i+1}(\mathbf{z}, \mathbf{v}))P_s(\mathbf{z}, \mathbf{v}) \\ &= 1 - p_i^* - \sum_{(\mathbf{z}, \mathbf{v}) \in \mathbb{Z}^m \times V} \min(P_t(\mathbf{z}, \mathbf{v}) - p_i(\mathbf{z}, \mathbf{v}), P_s(\mathbf{z}, \mathbf{v})(1 - p_i^*)) \\ &= 1 - p_i^* - \sum_{(\mathbf{z}, \mathbf{v}) \in \mathbb{Z}^m \times V} \alpha_{i+1}(\mathbf{z}, \mathbf{v}) \\ &= 1 - p_{i+1}^*. \end{aligned}$$

Then  $\forall(\mathbf{z}, \mathbf{v}) \in \mathbb{Z}^m \times V, \forall i \in \mathbb{N}, r(i, \mathbf{z}, \mathbf{v}) = \alpha_i(\mathbf{z}, \mathbf{v})$ . Let us now show that  $\sum_{j=1}^{\infty} r(j, \mathbf{z}, \mathbf{v}) = P_t(\mathbf{z}, \mathbf{v})$ . Note that  $p_i(\mathbf{z}, \mathbf{v}) = \sum_{j=1}^i \alpha_j(\mathbf{z}, \mathbf{v})$  so we can study these partial sums and show that they indeed converge to  $P_t(\mathbf{z}, \mathbf{v})$ .

To conclude the proof, we recall the proof from [HJMR07, Claim IV.1]. We reproduce it here for completeness.

Let us first show that  $\alpha_i(\mathbf{z}, \mathbf{v}) \geq (P_t(\mathbf{z}, \mathbf{v}) - p_{i-1}(\mathbf{z}, \mathbf{v}))P_s(\mathbf{z}, \mathbf{v})$ . We have

$$\begin{aligned} 1 - p_{i-1}^* &= \sum_{(\mathbf{z}, \mathbf{v}) \in \mathbb{Z}^m \times V} (P_t(\mathbf{z}, \mathbf{v}) - p_{i-1}(\mathbf{z}, \mathbf{v})) \\ &\geq P_t(\mathbf{z}, \mathbf{v}) - p_{i-1}(\mathbf{z}, \mathbf{v}). \end{aligned}$$

Then the above inequality holds by definition of  $\alpha_i(\mathbf{z}, \mathbf{v})$ : both  $P_t(\mathbf{z}, \mathbf{v}) - p_{i-1}(\mathbf{z}, \mathbf{v})$  and  $(1 - p_{i-1}^*)P_s(\mathbf{z}, \mathbf{v})$  are  $\geq (P_t(\mathbf{z}, \mathbf{v}) - p_{i-1}(\mathbf{z}, \mathbf{v}))P_s(\mathbf{z}, \mathbf{v})$ .

From that, we find that

$$P_t(\mathbf{z}, \mathbf{v}) - p_i(\mathbf{z}, \mathbf{v}) \leq (P_t(\mathbf{z}, \mathbf{v}) - p_{i-1}(\mathbf{z}, \mathbf{v}))(1 - P_s(\mathbf{z}, \mathbf{v})),$$

and a straightforward induction show that this is  $\leq P_t(\mathbf{z}, \mathbf{v})(1 - P_s(\mathbf{z}, \mathbf{v}))^i$ . Finally, by definition of  $p_{i-1}(\mathbf{z}, \mathbf{v})$ , it holds that  $P_t(\mathbf{z}, \mathbf{v}) - p_i(\mathbf{z}, \mathbf{v}) \geq 0$ .  $\square$